arXiv:gr-qc/0302084v1 20 Feb 2003

Hamiltonian Linearization of the Rest-Frame Instant Form of Tetrad Gravity in a Completely Fixed 3-Orthogonal Gauge: a Radiation Gauge for Background-Independent Gravitational Waves in a Post-Minkowskian Einstein Spacetime.

Juri Agresti
Dipartimento di Fisica
Universita' di Firenze
Polo Scientifico, via Sansone 1
50019 Sesto Fiorentino, Italy
E-mail agresti@fi.infn.it

and

Roberto De Pietri
Dipartimento di Fisica
Campus Universitario, Universita' di Parma
Viale delle Scienze, 43100 Parma, Italy
depietri@pr.infn.it

and

Luca Lusanna
Sezione INFN di Firenze
Polo Scientifico, via Sansone 1
50019 Sesto Fiorentino, Italy
E-mail lusanna@fi.infn.it

and

Luca Martucci
Dipartimento di Fisica and INFN Sezione di Milano
Universita' di Milano I
via G.Celoria 16
20133 Milano
E-mail luca.martucci@mi.infn.it

Abstract

In the framework of the rest-frame instant form of tetrad gravity, where the Hamiltonian is the weak ADM energy $\hat{E}_{ADM}$, we define a special completely fixed 3-orthogonal Hamiltonian gauge, corresponding to a choice of *non-harmonic* 4-coordinates, in which the independent degrees of freedom of the gravitational field are described by two pairs of canonically conjugate Dirac observables (DO) $r_{\bar{a}}(\tau, \vec{\sigma})$, $\pi_{\bar{a}}(\tau, \vec{\sigma})$, $\bar{a} = 1, 2$. We define a Hamiltonian linearization of the theory, i.e. gravitational waves, *without introducing any background 4-metric*, by retaining only the linear terms in the DO's in the super-hamiltonian constraint (the Lichnerowicz equation for the conformal factor of the 3-metric) and the quadratic terms in the DO's in $\hat{E}_{ADM}$. *We solve all the constraints* of the linearized theory: this amounts to work in a well defined post-Minkowskian Christodoulou-Klainermann space-time. The Hamilton equations imply the wave equation for the DO's $r_{\bar{a}}(\tau, \vec{\sigma})$, which replace the two polarizations of the TT harmonic gauge, and that *linearized Einstein's equations are satisfied* . Finally we study the geodesic equation, both for time-like and null geodesics, and the geodesic deviation equation.

February 7, 2008

Typeset using REVTEX

I. INTRODUCTION

In a series of papers [1–3] the rest-frame instant form of the Hamiltonian ADM formulation of both metric and tetrad gravity was given. The aim of these papers was to arrive at a formulation of general relativity with matter such that the switching off of the Newton constant G would produce the description of the same matter in the rest-frame instant form of parametrized Minkowski theories [4,5,1], with the general relativistic covariance deparametrizing to the special relativistic one of these theories, and to study its Hamiltonian formulation consistently with Dirac-Bergmann theory of constraints.

To implement this program we must select the following family of non-compact space-times:

i) *globally hyperbolic*, so that the ADM Hamiltonian formulation [6] is well defined if we start from the ADM action;

ii) *topologically trivial*, so that they can be foliated with space-like hyper-surfaces Σ_τ diffeomorphic to R^3 (3+1 splitting of space-time with τ, the scalar parameter labeling the leaves, as a *mathematical time*);

iii) *asymptotically flat at spatial infinity* and with boundary conditions at spatial infinity independent from the direction, so that the *spi group* of asymptotic symmetries is reduced to the Poincare' group with the ADM Poincare' charges as generators [1]. In this way we can eliminate the *super-translations*, namely the obstruction to define angular momentum in general relativity, and we have the same type of boundary conditions which are needed to get well defined non-Abelian charges in Yang-Mills theory, opening the possibility of a unified description of the four interactions with all the fields belonging to same function space [5]. All these requirements imply that the *allowed foliations* of space-time must have the space-like hyper-planes tending in a direction-independent way to Minkowski space-like hyper-planes at spatial infinity, which moreover must be orthogonal there to the ADM 4-momentum. But these are the conditions satisfied by the singularity-free Christodoulou-Klainermann space-times [7], in which, in presence of matter, the allowed hyper-surfaces define the *rest frame of the universe* and naturally become the Wigner hyper-planes of the rest-frame instant form of the parametrized Minkowski theories describing the same matter when $G \mapsto 0$. Therefore there are *preferred asymptotic inertial observers*, which can be identified with the *fixed stars*. These allowed hyper-surfaces are called *Wigner-Sen-Witten (WSW) hyper-surfaces*, because it can be shown that the Frauendiener re-formulation [8] of Sen-Witten equations [9] for triads allows (after the restriction to the solutions of Einstein's equations) to transport the asymptotic tetrads of the asymptotic inertial observers in each point of the hyper-surface, generating a *local compass of inertia* to be used to define *rotations with respect to the fixed stars* [2]. Besides the existence of a realization of the Poincare' group, the only other needed property is that the admissible WSW hyper-surfaces must admit an involution [10] allowing the definition of a generalized Fourier transform with its associated concepts of positive and negative energy, so to avoid the claimed impossibility to define particles in curved space-times [11].

---

[1] When we switch off the Newton constant G, the ADM Poincare' charges, expressed in Σ_τ-adapted 4-coordinates, become the generators of the *internal Poincare' group* of parametrized Minkowski theories [1].

[2] Instead the standard Fermi-Walker transport of the tetrads of a time-like observer is a standard

iv) All the fields have to belong to suitable *weighted Sobolev spaces* so that the allowed space-like hyper-surfaces are Riemannian 3-manifolds without Killing vectors: in this way we avoid the analogue of the Gribov ambiguity in general relativity and we can get a unification of the function spaces of gravity and particle physics.

After all these preliminaries it is possible to study the Hamiltonian formulation of both ADM metric [1] and tetrad [2,3] gravity [3] with their (8 and 14 respectively) first class constraints as generators of the Hamiltonian gauge transformations. Then it is possible to look, at least at a heuristic level, for Shanmugadhasan canonical transformations [12,4,5] implementing *the separation between the gauge variables and the Dirac observables* (DO) for the gravitational field [4]. A complete exposition of these topics is in Refs. [1–3], where it is shown that it is possible to define a *rest frame instant form of gravity* in which the effective Hamiltonian for the evolution is the *weak ADM energy* E_{ADM} [5] [17].

Let us consider tetrad gravity. In Refs. [2,3] there is a new parametrization of tetrad gravity, still utilizing the ADM action, which allows to solve 13 of the 14 first class constraints. After an allowed 3+1 splitting of space-time with space-like hyper-surfaces Σ_τ, we introduce adapted coordinates [6]. The arbitrary cotetrads ${}^4E^{(\alpha)}_\mu$, appearing in the 4-metric

---

of non-rotation with respect to a local observer in free fall.

[3] More natural for the coupling to the fermions of the standard model of particles. Moreover tetrad gravity is naturally a theory of time-like accelerated observers, generalizing the ones of parametrized Minkowski theories.

[4] See Refs. [13–15] for the interpretation of the gauge variables as *generalized inertial effects* and of the DO's (the non-local deterministically predictable physical degrees of freedom of the gravitational field) as *generalized tidal effects*. The non-locality of DO's (all 3-space is needed for their determination) may be interpreted as a form of Mach principle, even if our boundary conditions are not compatible with the standard interpretation of such principle given by Einstein and Wheeler, who choose spatially compact space-times without boundary [16].

[5] Therefore the formulations with a frozen reduced phase space are avoided. The super-hamiltonian constraint generates *normal* deformations of the space-like hyper-surfaces, which are *not* interpreted as a time evolution (like in the Wheeler-DeWitt approach) but as the Hamiltonian gauge transformations ensuring that the description of gravity is independent from the 3+1 splittings of space-time like it happens in parametrized Minkowski theories.

[6] Instead of local coordinates x^μ for M^4, we use local coordinates σ^A on $R \times \Sigma \approx M^4$ with $\Sigma \approx R^3$ $[x^\mu = z^\mu(\sigma)$ with inverse $\sigma^A = \sigma^A(x)]$, i.e. a Σ_τ-*adapted holonomic coordinate basis* for vector fields $\partial_A = \frac{\partial}{\partial \sigma^A} \in T(R \times \Sigma) \mapsto b^\mu_A(\sigma)\partial_\mu = \frac{\partial z^\mu(\sigma)}{\partial \sigma^A}\partial_\mu \in TM^4$, and for differential

of the ADM action principle, are rewritten in adapted coordinates, ${}^4E^{(\alpha)}_A$, and replaced at the Hamiltonian level by the lapse N and shift $N_{(a)} = {}^3e_{(a)r}\, N^r$ functions, cotriads ${}^3e_{(a)r}$ and boost parameters $\varphi^{(a)}$

$$\begin{pmatrix} {}^4E^{(o)}_A \\ {}^4E^{(a)}_A \end{pmatrix} = \begin{pmatrix} \sqrt{1+\sum_{(c)}\varphi^{(c)2}} & -\epsilon\varphi_{(b)} \\ \varphi^{(a)} & \delta^{(a)}_{(b)} - \epsilon \frac{\varphi^{(a)}\varphi_{(b)}}{1+\sqrt{1+\sum_{(c)}\varphi^{(c)2}}} \end{pmatrix} \times$$
$$\begin{pmatrix} {}^4_{(\Sigma)}\check{\tilde{E}}^{(o)}_A = (N; \vec{0}) \\ {}^4_{(\Sigma)}\check{\tilde{E}}^{(b)}_A = (N^{(b)} = {}^3e^{(b)}_r\, N^r; {}^3e^{(b)}_r) \end{pmatrix},$$

$$\Rightarrow {}^4g_{AB} = {}^4E^{(\alpha)}_A\, {}^4\eta_{(\alpha)(\beta)}\, {}^4E^{(\beta)}_B = {}^4_{(\Sigma)}\check{E}^{(\alpha)}_A\, {}^4\eta_{(\alpha)(\beta)}\, {}^4_{(\Sigma)}\check{E}^{(\beta)}_B =$$
$$= \epsilon \begin{pmatrix} (N^2 - {}^3g_{rs}N^rN^s) & -{}^3g_{st}N^t \\ -{}^3g_{rt}N^t & -{}^3g_{rs} \end{pmatrix},$$

$$\begin{pmatrix} {}^4E^{\mu}_{(o)} \\ {}^4E^{\mu}_{(a)} \end{pmatrix} = \begin{pmatrix} \sqrt{1+\sum_{(c)}\varphi^{(c)2}} & -\varphi^{(b)} \\ \epsilon\varphi_{(a)} & \delta^{(b)}_{(a)} - \epsilon \frac{\varphi_{(a)}\varphi^{(b)}}{1+\sqrt{1+\sum_{(c)}\varphi^{(c)2}}} \end{pmatrix} \begin{pmatrix} l^\mu \\ b^\mu_s\, {}^3e^s_{(b)} \end{pmatrix},$$

$$\begin{pmatrix} {}^4E^{A}_{(o)} \\ {}^4E^{A}_{(a)} \end{pmatrix} = \begin{pmatrix} \sqrt{1+\sum_{(c)}\varphi^{(c)2}} & -\varphi^{(b)} \\ \epsilon\varphi^{(a)} & \delta^{(b)}_{(a)} - \epsilon \frac{\varphi_{(a)}\varphi^{(b)}}{1+\sqrt{1+\sum_{(c)}\varphi^{(c)2}}} \end{pmatrix} \begin{pmatrix} {}^4_{(\Sigma)}\check{\tilde{E}}^A_{(o)} = (1/N; -N^r/N) \\ {}^4_{(\Sigma)}\check{\tilde{E}}^A_{(b)} = (0; {}^3e^r_{(b)}) \end{pmatrix},$$

---

one-forms $dx^\mu \in T^*M^4 \mapsto d\sigma^A = b^A_\mu(\sigma)dx^\mu = \frac{\partial\sigma^A(z)}{\partial z^\mu}dx^\mu \in T^*(R\times\Sigma)$. The 4-metric has Lorentz signature $\epsilon\,(+,-,-,-)$ with $\epsilon = \pm 1$ according to particle physics and general relativity conventions respectively. The induced 4-metric and inverse 4-metric become in the new basis

$${}^4g_{AB} = \{{}^4g_{\tau\tau} = \epsilon(N^2 - {}^3g_{rs}N^rN^s); {}^4g_{\tau r} = -\epsilon\, {}^3g_{rs}N^s; {}^4g_{rs} = -\epsilon\, {}^3g_{rs}\} =$$
$$= \epsilon\, [l_Al_B - {}^3g_{rs}\,(\delta^r_A + N^r\,\delta^\tau_A)(\delta^s_B + N^s\delta^\tau_B)],$$
$${}^4g^{AB} = \{{}^4g^{\tau\tau} = \frac{\epsilon}{N^2}; {}^4g^{\tau r} = -\frac{\epsilon\, N^r}{N^2}; {}^4g^{rs} = -\epsilon({}^3g^{rs} - \frac{N^rN^s}{N^2})\} = \epsilon[l^Al^B - {}^3g^{rs}\delta^A_r\delta^B_s].$$

For the unit normals to Σ_τ we have $l^A = \frac{\epsilon}{N}(1; -N^r)$ and $l_A = N\partial_A\tau = N\delta^\tau_A = (N; \vec{0})$. We introduce the 3-metric of Σ_τ: ${}^3g_{rs} = -\epsilon\, {}^4g_{rs}$ with signature $(+++)$. If ${}^4\gamma^{rs}$ is the inverse of the spatial part of the 4-metric (${}^4\gamma^{ru}\, {}^4g_{us} = \delta^r_s$), the inverse of the 3-metric is ${}^3g^{rs} = -\epsilon\, {}^4\gamma^{rs}$ (${}^3g^{ru}\, {}^3g_{us} = \delta^r_s$). We have the following form for the line element in M^4:

$$ds^2 = \epsilon(N^2 - {}^3g_{rs}N^rN^s)(d\tau)^2 - 2\epsilon\, {}^3g_{rs}N^sd\tau d\sigma^r - \epsilon\, {}^3g_{rs}d\sigma^rd\sigma^s =$$
$$= \epsilon\Big[N^2(d\tau)^2 - {}^3g_{rs}(d\sigma^r + N^rd\tau)(d\sigma^s + N^sd\tau)\Big].$$

$$\begin{aligned}\Rightarrow {}^4g^{AB} &= {}^4E^A_{(\alpha)}\, {}^4\eta^{(\alpha)(\beta)}\, {}^4E^B_{(\beta)} = {}^4_{(\Sigma)}\check{E}^A_{(\alpha)}\, {}^4\eta_{(\alpha)(\beta)}\, {}^4_{(\Sigma)}\check{E}^B_{(\beta)} = \\ &= \epsilon \begin{pmatrix} \frac{1}{N^2} & -\frac{N^s}{N^2} \\ -\frac{N^r}{N^2} & -({}^3g^{rs} - \frac{N^r N^s}{N^2}) \end{pmatrix}. \end{aligned} \tag{1.1}$$

The conjugate momenta are $\tilde{\pi}^N(\tau,\vec{\sigma})$, $\tilde{\pi}^{\vec{N}}_{(a)}(\tau,\vec{\sigma})$, ${}^3\tilde{\pi}^r_{(a)}(\tau,\vec{\sigma})$, $\tilde{\pi}^{\vec{\varphi}}_{(a)}(\tau,\vec{\sigma})$, respectively. There are ten primary constraints $\tilde{\pi}^N(\tau,\vec{\sigma}) \approx 0$, $\tilde{\pi}^{\vec{N}}_{(a)}(\tau,\vec{\sigma}) \approx 0$, $\tilde{\pi}^{\vec{\varphi}}_{(a)}(\tau,\vec{\sigma}) \approx 0$, ${}^3\tilde{M}_{(a)}(\tau,\vec{\sigma}) = \epsilon_{(a)(b)(c)}\, {}^3e_{(b)r}(\tau,\vec{\sigma})\, {}^3\tilde{\pi}^r_{(c)}(\tau,\vec{\sigma}) \approx 0$ and four secondary ones: the super-hamiltonian constraint $\mathcal{H}(\tau,\vec{\sigma}) \approx 0$ and the super-momentum constraints $\mathcal{H}^r(\tau,\vec{\sigma}) \approx 0$. Therefore there are *14 arbitrary gauge variables*, four of which are the lapse and shift functions. All the constraints are first class and the Dirac Hamiltonian depends on 10 Dirac multipliers.

As shown in Ref. [1], a consistent treatment of the boundary conditions at spatial infinity requires the explicit separation of the *asymptotic* part of the lapse and shift functions from their *bulk* part: $N(\tau,\vec{\sigma}) = N_{(as)}(\tau,\vec{\sigma}) + n(\tau,\vec{\sigma})$, $N_r(\tau,\vec{\sigma}) = N_{(as)r}(\tau,\vec{\sigma}) + n_r(\tau,\vec{\sigma})$, with $n(\tau,\vec{\sigma})$ and $n_r(\tau,\vec{\sigma})$ tending to zero at spatial infinity in a direction-independent way. On the contrary, $N_{(as)}(\tau,\vec{\sigma}) = -\lambda_\tau(\tau) - \frac{1}{2}\,\lambda_{\tau u}(\tau)\,\sigma^u$ and $N_{(as)r}(\tau,\vec{\sigma}) = -\lambda_r(\tau) - \frac{1}{2}\,\lambda_{ru}(\tau)\,\sigma^u$. The *Christodoulou-Klainermann space-times* [7], with their *rest-frame* condition of zero ADM 3-momentum and absence of super-translations, are singled out by these considerations. As already said the allowed foliations of these space-times tend asymptotically to Minkowski hyper-planes in a direction-independent way and are asymptotically orthogonal to the ADM four-momentum. The leaves Σ_τ are the WSW hyper-surfaces. They have $N_{(as)}(\tau,\vec{\sigma}) = -\epsilon$, $N_{(as)r}(\tau,\vec{\sigma}) = 0$. As in Refs. [1–3], from now on the lapse and shift functions N and N_r will be replaced by $-\epsilon + n$ and n_r.

It is pointed out in the papers [1,3] that in order to have well defined asymptotic *weak and strong ADM Poincare' charges* (generators of the asymptotic Poincare' group) all fields must have a suitable direction-independent limit at spatial infinity. Recall that the *strong* ADM energy is the flux through the surface at spatial infinity of a function of the 3-metric only, and it is weakly equal to the *weak* ADM energy (volume form) which contains all the dependence on the ADM momenta. This implies [1] that the super-hamiltonian constraint must be interpreted as the equation (*Lichnerowicz equation*) that uniquely determines the *conformal factor* $\phi(\tau,\vec{\sigma}) = (det\, {}^3g(\tau,\vec{\sigma}))^{1/12}$ of the 3-metric as a functional of the other variables. This means that the associated gauge variable is the *canonical momentum* $\pi_\phi(\tau,\vec{\sigma})$ *conjugate to the conformal factor*: it carries information about the extrinsic curvature of Σ_τ. It is just this variable, and *not* York's time, which parametrizes the *normal* deformation of the embeddable space-like hyper-surfaces Σ_τ. Since different forms of Σ_τ correspond to different 3+1 splittings of M^4, the gauge transformations generated by the super-hamiltonian constraint correspond to the transition from an allowed 3+1 splitting to another one: this is the gauge orbit in the phase space over super-space. The theory is therefore independent of the choice of the 3+1 splitting like in parametrized Minkowski theories. As a matter of fact, a gauge fixing for the super-hamiltonian constraint is a choice of a particular 3+1 splitting and this is done by fixing the momentum $\pi_\phi(\tau,\vec{\sigma})$ conjugate to the conformal factor. This shows the dominant role of the conformal 3-geometries in the determination of the physical degrees of freedom, just as in the Lichnerowicz-York conformal approach [18,19,16].

As explained in Refs. [1,3], following a suggestion of Dirac [20], we restrict our space-times to *admit asymptotic Minkowski Cartesian coordinates*, namely the admissible 4-coordinate systems $x^\mu = z^\mu(\sigma^A)$ have the property $x^\mu \rightarrow \delta^\mu_{(\mu)}\, z^{(\mu)}_{(\infty)}(\tau,\vec{\sigma})$ at spatial infinity with $z^{(\mu)}_{(\infty)}(\tau,\vec{\sigma}) = x^{(\mu)}_{(\infty)}(\tau) + b^{(\mu)}_{(\infty) r}(\tau)\, \sigma^r$ [(μ) are flat indices]. Therefore the leaves of the admissible 3+1 splittings of our space-times tend to Minkowski space-like hyper-planes at spatial infinity in a direction-independent way. While $x^{(\mu)}_{(\infty)}(\tau)$ denotes the world-line of a point (centroid) arbitrarily chosen as origin of the 3-coordinate systems on each Σ_τ, the $b^{(\mu)}_{(\infty) A}(\tau)$'s are flat asymptotic tetrads with $b^{(\mu)}_{(\infty)\tau}$ equal to the normal $l^{(\mu)}_{(\infty)}$ to the asymptotic Minkowski hyper-planes. To force the existence of these asymptotic coordinates, we must add to tetrad gravity ten extra configuration degrees of freedom, namely the centroid $x^{(\mu)}_{(\infty)}(\tau)$ and the six independent components of the tetrads $b^{(\mu)}_{(\infty) A}(\tau)$, and of the conjugate ten momenta, $p^{(\mu)}_{(\infty)}$ and a spin tensor $S^{(\mu)(\nu)}_{(\infty)}$. As shown in Refs. [1,3], this increase of variables is balanced by adding ten extra first class constraints determining the ten extra momenta: $\chi^A = p^A_{(\infty)} - \hat{P}^A_{ADM} \approx 0$, $\chi^{AB} = J^{AB}_{(\infty)} - \hat{J}^{AB}_{ADM} \approx 0$, where the weak (volume form) ADM Poincare' charges $\hat{P}^A_{ADM}$, $\hat{J}^{AB}_{ADM}$ and the quantities $p^A_{(\infty)} = b^A_{(\infty)(\mu)}\, p^{(\mu)}_{(\infty)}$, $J^{AB}_{(\infty)} = b^A_{(\infty)(\mu)}\, b^B_{(\infty)(\nu)}\, S^{(\mu)(\nu)}_{(\infty)}$ are expressed in Σ_τ-adapted 4-coordinates.

In this way the ten extra configuration degrees of freedom become arbitrary *gauge variables*. After the splitting of the lapse and shift functions in the asymptotic and bulk parts, this approach [1,3] implies the replacement of the canonical Hamiltonian $\int d^3\sigma \Big[n\, \mathcal{H} + n_r\, \mathcal{H}^r\Big](\tau,\vec{\sigma}) + \lambda_A(\tau)\, \hat{P}^A_{ADM} + \frac{1}{2}\, \lambda_{AB}(\tau)\, \hat{J}^{AB}_{ADM}$ with the Hamiltonian $\int d^3\sigma \Big[n\, \mathcal{H} + n_r\, \mathcal{H}^r + \lambda_n\, \pi_n + \lambda_{\vec{n}\, r}\, \pi^r_{\vec{n}} + \mu_{(a)}\, M_{(a)} + \rho_{(a)}\, \pi^{\vec{\varphi}}_{(a)}\Big](\tau,\vec{\sigma}) + \lambda_A(\tau)\, [p^A_{(\infty)} - \hat{P}^A_{ADM}] + \frac{1}{2}\, \lambda_{AB}(\tau)\, [J^{AB}_{(\infty)} - \hat{J}^{AB}_{ADM}]$.

The following boundary conditions (compatible with Christodoulou-Klainermann space-times) ensure that the Hamiltonian gauge transformations preserve these asymptotic properties defining our class of space-times

$$\begin{aligned}
&{}^3e_{(a)r}(\tau,\vec{\sigma}) \rightarrow_{r\rightarrow\infty} (1 + \frac{M}{2r})\delta_{(a)r} + O(r^{-3/2}),\\
&\qquad \Rightarrow {}^3g_{rs}(\tau,\vec{\sigma}) = [{}^3e_{(a)r}\, {}^3e_{(a)s}](\tau,\vec{\sigma}) \rightarrow_{r\rightarrow\infty} (1 + \frac{M}{r})\delta_{rs} + O(r^{-3/2}),\\
&{}^3\tilde{\pi}^r_{(a)}(\tau,\vec{\sigma}) \rightarrow_{r\rightarrow\infty} O(r^{-5/2}),\\
&n(\tau,\vec{\sigma}) \rightarrow_{r\rightarrow\infty} O(r^{-(2+\epsilon)}), \qquad \epsilon > 0,\\
&n_r(\tau,\vec{\sigma}) \rightarrow_{r\rightarrow\infty} O(r^{-\epsilon}),\\
&\alpha_{(a)}(\tau,\vec{\sigma}) \rightarrow_{r\rightarrow\infty} O(r^{-(1+\epsilon)}),\\
&\varphi_{(a)}(\tau,\vec{\sigma}) \rightarrow_{r\rightarrow\infty} O(r^{-(1+\epsilon)}).
\end{aligned} \tag{1.2}$$

As shown in Refs. [1,3], to get the rest-frame instant form of tetrad gravity, with its

WSW hyper-surfaces as leaves of the admissible 3+1 splittings (when $p^{(\mu)}_{(\infty)}$ is time-like) the gauge fixings must be added with the following procedure :

i) Add three gauge fixings on the boost parameters, namely $\varphi^{(a)}(\tau, \vec{\sigma}) \approx 0$: in this way we choose cotetrads adapted to Σ_τ. The time constancy of this gauge fixing determines the 3 Dirac multipliers $\rho_{(a)}(\tau, \vec{\sigma})$.

ii) Add three gauge fixings, $\alpha_{(a)}(\tau, \vec{\sigma}) \approx 0$, to the rotation constraints ${}^3\tilde{M}_{(a)}(\tau, \vec{\sigma}) \approx 0$: in this way we fix a reference orientation of the cotriads. The time constancy of these gauge fixings determine the 3 Dirac multipliers $\mu_{(a)}(\tau, \vec{\sigma})$.

iii) Add three gauge fixings $\chi_r(\tau, \vec{\sigma}) \approx 0$ to the secondary super-momentum constraints: this amounts to a choice of 3-coordinates on Σ_τ. The requirement of time constancy of the constraints $\chi_r(\tau, \vec{\sigma}) \approx 0$ will generate three gauge fixings $\varphi_r(\tau, \vec{\sigma}) \approx 0$ for the primary constraints $\pi^r_n(\tau, \vec{\sigma}) \approx 0$, which determine the shift functions $n_r(\tau, \vec{\sigma})$ (and therefore the coordinate-dependent gravito-magnetic aspects and the eventual anisotropy of light propagation). The time constancy of the $\varphi_r(\tau, \vec{\sigma})$'s will determine the 3 Dirac multipliers $\lambda_{\vec{n}r}(\tau, \vec{\sigma})$.

iv) Add a gauge fixing $\chi(\tau, \vec{\sigma}) \approx 0$ to the secondary super-hamiltonian constraint, which determines the form of the space-like hyper-surface Σ_τ. Its time constancy produces the gauge fixing $\varphi(\tau, \vec{\sigma}) \approx 0$ for the primary constraint $\pi_n(\tau, \vec{\sigma}) \approx 0$, which determines the lapse function $n(\tau, \vec{\sigma})$, i.e. how the surfaces Σ_τ are packed in the foliation. Now the 3+1 splitting of space-time is completely determined and the time constancy of $\varphi(\tau, \vec{\sigma}) \approx 0$ determines the last Dirac multiplier $\lambda_n(\tau, \vec{\sigma})$. A posteriori after having solved the Hamilton equations one could find the embedding $z^\mu(\tau, \vec{\sigma})$ of the WSW hyper-surfaces into the space-time.

v) Add 6 suitable gauge fixings on the gauge variables $b^{(\mu)}_{(\infty)A}(\tau)$, so that, after having gone to Dirac brackets, we get $J^{AB}_{(\infty)} \equiv \hat{J}^{AB}_{ADM}$ and $\lambda_{AB}(\tau) = 0$. As a consequence of this gauge fixing, the gauge variables $x^{(\mu)}_{(\infty)}(\tau)$ must be replaced by the *canonical non-covariant 4-center of mass* $\tilde{x}^{(\mu)}_{(\infty)}(\tau)$ of the universe. All this implies that the asymptotic Minkowski space-like hyper-planes become orthogonal to the weak ADM 4-momentum $\hat{P}^A_{ADM}$ and that the canonical Hamiltonian becomes $\int d^3\sigma \Big[n\, \mathcal{H} + n_r\, \mathcal{H}^r + \lambda_n\, \pi_n + \lambda_{\vec{n}\, r}\, \pi^r_{\vec{n}} + \mu_{(a)}\, M_{(a)} + \rho_{(a)}\, \pi^{\vec{\varphi}}_{(a)}\Big](\tau, \vec{\sigma}) - \lambda_\tau(\tau)\, [\epsilon_{(\infty)} - \hat{P}^\tau_{ADM}] + \lambda_r(\tau)\, \hat{P}^r_{ADM}$ with $\epsilon_{(\infty)} = -\epsilon\, \sqrt{\epsilon\, p^2_{(\infty)}}$. Namely only the four first class constraints $\epsilon_{(\infty)} - \hat{P}^\tau_{ADM} \approx 0$ and $\hat{P}^r_{ADM} \approx 0$ are left, With the constraints $\hat{P}^r_{ADM} \approx 0$ being *the rest frame condition for the universe.*

vi) Add the gauge fixing $\tau - T_{(\infty)} \approx 0$, implying $\lambda_\tau(\tau) = 0$ and identifying the mathematical time τ with the rest frame time $T_{(\infty)} = p_{(\infty)(\mu)}\, \tilde{x}^{(\mu)}_{(\infty)}/\epsilon_{(\infty)} = p_{(\infty)(\mu)}\, x^{(\mu)}_{(\infty)}/\epsilon_{(\infty)}$ of the universe. The canonical 4-center of mass $\tilde{x}^{(\mu)}_{(\infty)}$ becomes a *decoupled point particle clock.*

vii) Either add the natural gauge fixings $\hat{J}^{\tau r}_{ADM} \approx 0$ to the constraints $\hat{P}^r_{ADM} \approx 0$ or work by choosing $\lambda_r(\tau) = 0$ as a pre-gauge-fixing (see the discussion at the beginning of Section V) .

At this stage the canonical reduction is completed by going to Dirac brackets and it can be shown [1,3] that the Dirac Hamiltonian for the rest-frame instant form of tetrad gravity is the *weak* (volume form) ADM energy: $H_D = \hat{E}_{ADM} = -\epsilon\, \hat{P}^\tau_{ADM}$. It becomes the effective Hamiltonian for the gauge invariant observables parametrizing the reduced phase space in the rest-frame instant form.

To find a canonical basis of Dirac observables for the gravitational field in absence of known solutions of the super-hamiltonian constraint, we can perform a quasi-Shanmugadhasan canonical transformation adapted to only 13 of the constraints and utilize the information that this constraint has to be interpreted as the *Lichnerowicz equation* for the conformal factor $\phi(\tau,\vec{\sigma})=\gamma^{1/12}(\tau,\vec{\sigma})=(det\,{}^3g(\tau,\vec{\sigma}))^{1/12}=e^{q(\tau,\vec{\sigma})/2}$ of the 3-metric. The result of this *point* canonical transformation is ($\bar{a}=1,2$)

$$\begin{array}{|c|c|c|c|} \hline \varphi^{(a)} & n & n_r & {}^3e_{(a)r} \\ \hline \approx 0 & \approx 0 & \approx 0 & {}^3\tilde{\pi}^r_{(a)} \\ \hline \end{array} \quad \longrightarrow \quad \begin{array}{|ccccc|c|c|} \hline \varphi^{(a)} & n & n_r & \alpha_{(a)} & \xi^r & \phi & r_{\bar{a}} \\ \hline \approx 0 & \approx 0 & \approx 0 & \approx 0 & \approx 0 & \pi_\phi & \pi_{\bar{a}} \\ \hline \end{array} \tag{1.3}$$

where $\alpha_{(a)}(\tau,\vec{\sigma})$ are angles and $\xi^r(\tau,\vec{\sigma})$ are a parametrization of the group manifold of the passive 3-diffeomorphisms of Σ_τ, describing its changes of 3-coordinates [7]. The entries ≈ 0 are the new momenta corresponding to 13 Abelianized first class constraints: besides the lapse, shift and boost momenta, there are three Abelianized rotation constraints $\tilde{\pi}^{\vec{\alpha}}_{(a)}(\tau,\vec{\sigma})\approx 0$ and three Abelianized super-momentum constraints $\tilde{\pi}^{\vec{\xi}}_r(\tau,\vec{\sigma})\approx 0$.

The Hamiltonian *gauge variables* are the 13 configuration variables $\varphi^{(a)}(\tau,\vec{\sigma})$, $n(\tau,\vec{\sigma})$, $n_r(\tau,\vec{\sigma})$, $\alpha_{(a)}(\tau,\vec{\sigma})$, $\xi^r(\tau,\vec{\sigma})$ (they depend on the cotetrads and its space gradients) and the momentum $\pi_\phi(\tau,\vec{\sigma})$ conjugate to the conformal factor (it depends also on the time derivative of the cotetrads). The variables $\xi^r(\tau,\vec{\sigma})$ and $\pi_\phi(\tau,\vec{\sigma})$ can be thought as a possible 4-coordinate system with the Lorentz signature given by the pattern *3 configuration + 1 momentum* variables.

Even if we do not know the expression of the final variables $\alpha_{(a)}$, ξ^r, π_ϕ, $r_{\bar{a}}$, $\pi_{\bar{a}}$ in terms of the original variables, the point nature of the canonical transformation allows to write the following inverse relations (the form of the cotriad was determined by solving the multi-temporal equations for the gauge transformations generated by the first class constraints [3])

$$\begin{aligned}
{}^3e_{(a)r}(\tau,\vec{\sigma}) &= {}^3R_{(a)(b)}(\alpha_{(e)}(\tau,\vec{\sigma}))\,\frac{\partial\xi^s(\tau,\vec{\sigma})}{\partial\sigma^r}\,{}^3\hat{e}_{(b)s}(\phi(\tau,\vec{\xi}(\tau,\vec{\sigma})),r_{\bar{a}}(\tau,\vec{\xi}(\tau,\vec{\sigma}))\,),\\
{}^3\tilde{\pi}^r_{(a)}(\tau,\vec{\sigma}) &= \sum_s\int d^3\sigma_1\,\mathcal{K}^r_{(a)s}(\vec{\sigma},\vec{\sigma}_1,\tau|\alpha_{(e)},\xi^u,\phi,r_{\bar{a}}]\,\Big(\phi^{-2}\,e^{-\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}s}\,r_{\bar{a}}}\Big)(\tau,\vec{\sigma}_1)\\
&\quad\Big[\frac{\pi_\phi}{6\phi}+\sqrt{3}\sum_{\bar{b}}\gamma_{\bar{b}s}\,\pi_{\bar{b}}\Big](\tau,\vec{\sigma}_1)+\\
&\quad+\int d^3\sigma_1\,F^r_{(a)(b)}(\vec{\sigma},\vec{\sigma}_1,\tau|\alpha_{(e)},\xi^u,\phi,r_{\bar{a}}]\,\tilde{\pi}^{\vec{\alpha}}_{(b)}(\tau,\vec{\sigma}_1)+
\end{aligned}$$

---

[7] Since there is no canonical choice of an origin in the 3-diffeomorphism group manifold, the choice of a 3-coordinate system is done in two steps: i) by adding the gauge fixings $\xi^r(\tau,\vec{\sigma})-\sigma^r\approx 0$, declaring that this 3-coordinate system $\{\sigma^r\}$ on Σ_τ is conventionally chosen as origin of the 3-coordinate systems; ii) by choosing a parametrization of the reduced cotriads, and therefore of the 3-metric associated with the chosen 3-coordinate system, only in terms of the 3 functions $\phi(\tau,\vec{\sigma})$, $r_{\bar{a}}(\tau,\vec{\sigma})$.

$$+\sum_u \int d^3\sigma_1\, G^{ru}_{(a)}(\vec{\sigma},\vec{\sigma}_1,\tau|\alpha_{(e)},\xi^u,\phi,r_{\bar{a}}]\, \tilde{\pi}^{\vec{\xi}}_u(\tau,\vec{\sigma}_1),$$

$$\Downarrow$$

$$\begin{aligned}{}^3g_{rs}(\tau,\vec{\sigma}) &= \sum_a {}^3e_{(a)r}(\tau,\vec{\sigma})\, {}^3e_{(a)s}(\tau,\vec{\sigma}) = \\ &= \frac{\partial \xi^u(\tau,\vec{\sigma})}{\partial\sigma^r}\, \frac{\partial \xi^v(\tau,\vec{\sigma})}{\partial\sigma^s} \sum_a [{}^3e_{(a)u}\, {}^3e_{(a)v}](\phi(\tau,\vec{\xi}(\tau,\vec{\sigma})), r_{\bar{a}}(\tau,\vec{\xi}(\tau,\vec{\sigma}))).\end{aligned} \tag{1.4}$$

Here ${}^3R$ are arbitrary rotation matrices and ${}^3\hat{e}_{(a)r}$ are reduced cotriads depending only on the three functions ϕ and $r_{\bar{a}}$. The kernels $\mathcal{K}^r_{(a)u}$, $F^r_{(a)(b)}$ and $G^{ru}_{(a)}$ are not known explicitly: they are the solution of quasi-linear partial differential equations determined by the canonicity of the point transformation and by the fact that the rotation constraints do not have vanishing Poisson brackets with the super-momentum constraints [3]. Once we have found the general solution of the quasi-linear equations for the kernels and we have done a definite choice of gauge, the six rotation and super-momentum constraints give further restrictions on the kernels, which amount to restrict the general solution to a particular one.

The family of *3-orthogonal gauges* (all having a diagonal 3-metric, ${}^3g_{rs} = {}^3e_{(a)r}\, {}^3e_{(a)s} = 0 \quad for\ r \neq s$), which is parametrized by the last gauge variable $\pi_\phi(\tau,\vec{\sigma})$, is determined by the gauge fixings $\xi^r(\tau,\vec{\sigma}) - \sigma^r \approx 0$ and by the following parametrization of the reduced cotriads

$${}^3\hat{e}_{(a)r} = \delta_{(a)r}\, \phi^2\, e^{\frac{1}{\sqrt{3}}\sum_{\bar{a}} \gamma_{\bar{a}r}\, r_{\bar{a}}},$$

$$\Downarrow$$

$${}^3\hat{g}_{rs} = \sum_a {}^3\hat{e}_{(a)r}\, {}^3\hat{e}_{(a)s} = \delta_{rs}\, \phi^4\, e^{\frac{2}{\sqrt{3}}\sum_{\bar{a}} \gamma_{\bar{a}r}\, r_{\bar{a}}}. \tag{1.5}$$

The numerical constants $\gamma_{\bar{a}r}$, satisfying $\sum_{\bar{a}} \gamma_{\bar{a}r} = 0$, $\sum_u \gamma_{\bar{a}u}\, \gamma_{\bar{b}u} = \delta_{\bar{a}\bar{b}}$, $\sum_{\bar{a}} \gamma_{\bar{u}}\, \gamma_{\bar{a}v} = \delta_{uv} - \frac{1}{3}$, define a one-parameter family of quasi-Shanmugadhasan canonical bases (1.3). In the gauge $\alpha_{(a)}(\tau,\vec{\sigma}) \approx 0$, $\xi^r(\tau,\vec{\sigma}) - \sigma^r \approx 0$, we have ${}^3e_{(a)r}(\tau,\vec{\sigma}) \approx {}^3\hat{e}_{(a)r}(\tau,\vec{\sigma})$.

The physical deterministically predictable degrees of freedom of the gravitational field are the non-local DO's (their expression in terms of the original variables is not known) $r_{\bar{a}}(\tau,\vec{\sigma})$, $\pi_{\bar{a}}(\tau,\vec{\sigma})$, $\bar{a} = 1, 2$. In general they are not Bergmann observables, i.e. coordinate independent quantities, being non-tensorial and coordinate-dependent.

Even if we do not know the solution $\phi = \phi[\xi^r, \pi_\phi, r_{\bar{a}}, \pi_{\bar{a}}]$ of the Lichnerowicz equation, *the class of Hamiltonian gauges defined by the gauge fixing $\chi(\tau,\vec{\sigma}) = \pi_\phi(\tau,\vec{\sigma}) \approx 0$ has the special property that the DO's $r_{\bar{a}}(\tau,\vec{\sigma})$, $\pi_{\bar{a}}(\tau,\vec{\sigma})$ remain canonical also at the level of Dirac brackets*, so that these gauges can be named *radiation gauges*. When in a radiation gauge there is no other residual gauge freedom, we can express every tensor over the space-time only in terms of the two pairs of canonically conjugate DO's.

This allows *for the first time* to arrive at a completely fixed Hamiltonian gauge of the 3-orthogonal type [8], which when restricted to the solutions of Einstein's equations (i.e. *on-shell*) is equivalent to a well defined choice of 4-coordinates for space-time.

It is evident that the Hamiltonian gauge variables of canonical gravity carry an information about observers in space-time, so that they are not inessential variables like in electromagnetism and Yang-Mills theory but take into account the fact that in general relativity global inertial reference frames do not exist [9]. The separation between gauge variables and DO's is an extra piece of (non-local) information [13–15], which has to be added to the equivalence principle, asserting the local impossibility to distinguish gravitational from inertial effects, to visualize which of the local forces acting on test matter are *generalized inertial (or fictitious) forces depending on the Hamiltonian gauge variables* and which are *genuine gravitational forces depending on the DO's, which are absent in Newtonian gravity* [10]. Both types of forces have a *different appearance in different 4-coordinate systems.*

In every 4-coordinate system (on-shell completely fixed Hamiltonian gauge)

i) the *genuine tidal gravitational forces* appearing in the geodesic deviation equation will be well defined gauge-dependent functionals only of the DO's associated to that gauge, so that DO's can be considered *generalized non-local tidal degrees of freedom* ;

ii) the geodesics will have a different geometrical form which again is functionally dependent on the DO's in that gauge;

iii) the description of the relative 3-acceleration of a free particle in free fall given in the local rest frame of an observer will generated various terms identifiable with the general relativistic extension of the non-relativistic inertial accelerations and again these terms will depend on both the DO's and the Hamiltonian gauge variables [11].

Therefore the Hamiltonian gauge variables, which change value from a gauge to another one, *describe the change in the appearance* of both the physical and apparent gravitational forces going (on-shell) from a coordinate system to another one.

---

[8] Namely with ${}^3g_{rs}(\tau,\vec{\sigma})$ diagonal and with ${}^3g_{rr}(\tau,\vec{\sigma}) = f_r(r_{\bar{a}}(\tau,\vec{\sigma}))$ after the solution of the Lichnerowicz equation. The 3-orthogonal class of gauges seems to be the nearest one to the physical laboratories on the Earth. Let us remember that the standards of length and time are *coordinate units* and not Bergmann observables [21].

[9] The equivalence principle only allows the existence of local inertial frames along time-like geodesics describing the world-line of a scalar test particle in free fall.

[10] When dynamical matter will be introduced, this Hamiltonian procedure will lead to distinguish among action-at-a-distance, gravitational and inertial effects.

[11] Note that the coordinate-dependent definition of gravito-magnetism as the effects induced by ${}^4g_{\tau r}$ is a pure inertial effect, because it is determined by the shift gauge variables.

In this paper we shall start from tetrad gravity in the preferred completely fixed 3-orthogonal gauge $\pi_\phi(\tau, \vec{\sigma}) \approx 0$ [12] and we shall define a background-independent Hamiltonian linearization of the theory after this gauge fixation, which, on-shell, corresponds to a definite choice of a 4-coordinate system.

The *standard linearization* [22–24] of Einstein's equations selects those space-times which admit nearly Lorentzian 4-coordinate systems and whose 4-metric is well approximated by the *splitting* ${}^4g_{\mu\nu} = {}^4\eta_{\mu\nu} + {}^4h_{\mu\nu}$, $|{}^4h_{\mu\nu}| << 1$ (*weak field* approximation) with ${}^4\eta_{\mu\nu}$ the *background* Minkowski 4-metric in Cartesian coordinates. Then it is assumed that $\partial_\alpha\, {}^4h_{\mu\nu} = \frac{1}{L} O({}^4h_{\mu\nu})$, where for the length L one can take the *reduced wavelength* $\lambda/2\pi$ *of the resulting gravitational waves* [13]. Namely, one *assumes* the existence of solutions of Einstein's equations admitting nearly Cartesian 4-coordinates and *split them in a background* and *a perturbation*, with a residual gauge freedom in the choice of 4-coordinates. Then, after the restriction to the family of *harmonic gauges*, one *replaces the residual gauge freedom on the harmonic 4-coordinates with the gauge freedom of a manifestly covariant spin-2 theory over Minkowski space-time with Cartesian coordinates.* Gravitational waves are usually analyzed in the special *TT harmonic gauge*, a special case of the Lorentz gauges of the spin-2 theory. Therefore, there is a *discontinuity in the conceptual interpretation* and one gets that a special relativistic theory with its associated absolute space-time chrono-geometric structure and theory of measurement replaces Einstein space-time with its problematic concerning the identification of point-events (general covariance and the Hole Argument) and with a theory of measurement in curved space-times still to be developed (we have only an axiomatic theory employing *test* non-dynamical objects) [13–15].

Instead, by working in the preferred completely fixed Hamiltonian gauge with well determined 3-coordinates on the WSW hyper-surfaces and well determined lapse and shift functions (so that the reconstruction of Einstein space-time and of its 4-coordinates can be done by using the embedding of WSW hyper-surfaces), we have the possibility of making the linearization of the theory on the WSW hyper-surfaces by approximating the Lichnerowicz equation and the Hamiltonian, namely the weak ADM energy, with their linear and quadratic parts in the DO's respectively, *without never introducing a background.* In this way we may evaluate the linearized conformal factor of the 3-metric and the linearized lapse and shift functions and to obtain a linearized Einstein space-time with 3-orthogonal coordinates on the WSW hyper-surfaces (with a linearized embedding into the space-time), namely a *post-Minkowskian Einstein space-time* linearization of a Christodoulou-Klainermann space-time. It turns out that this gauge is *not* a harmonic gauge (and, as a consequence, not a TT harmonic gauge), but rather it is a *radiation gauge* for the gravitational field without any residual gauge freedom. The DO's $r_{\bar a}(\tau, \vec{\sigma})$ replace the two polarizations e_+ and $e_\times$ of the TT harmonic gauge. Moreover, besides the absence of background, there is also *no post-Newtonian expansion*: our approximate solution of Einstein equations in the chosen 4-coordinate system describes a linearized Einstein manifold with Lorentz signature near to the Minkowski space-time. It can be called a *post-Minkowskian space-time* with a linearized gravitational field dynamically modifying special relativity.

---

[12] The other gauge fixings are $\varphi_{(a)}(\tau, \vec{\sigma}) \approx 0$, $\alpha_{(a)}(\tau, \vec{\sigma}) \approx 0$, $\xi^r(\tau, \vec{\sigma}) - \sigma^r \approx 0$ and Eqs.(1.5).

[13] With the flat Minkowski background, the background Riemann tensor vanishes and the background radius of curvature $\mathcal{R}$ is infinite ($\mathcal{R}^{-2}$ is of the order of the Riemann tensor). If $\mathcal{A}$ is the amplitude of the gravitational wave, the *weak field* approximation is valid if $\mathcal{A} << 1$. In the more general *short-wave* approximation [22] the background is a vacuum Einstein space-time with 4-metric ${}^4g^{(B)}_{\mu\nu}$ and typical background radius of curvature $\mathcal{R}$. The splitting ${}^4g_{\mu\nu} = {}^4g^{(B)}_{\mu\nu} + {}^4h_{\mu\nu}$ is done in *steady* 4-coordinates, where, if $\mathcal{A}$ is the amplitude of the perturbation, one has: 1) ${}^4h_{\mu\nu} = O(\mathcal{A})$; 2) $\partial_\alpha\, {}^4g^{(B)}_{\mu\nu} = O(\mathcal{R}^{-1})$; 3) $\partial_\alpha\, {}^4h_{\mu\nu} = O((\frac{\lambda}{2\pi})^{-1})$. The short-wave approximation is valid if $\mathcal{A} << 1$ and $\frac{\lambda}{2\pi} << \mathcal{R}$.

In the special 3-orthogonal gauge with $\pi_\phi(\tau, \vec{\sigma}) = 0$, the canonical variables for the gravitational field are $r_{\bar a}(\tau, \vec{\sigma})$, $\pi_{\bar a}(\tau, \vec{\sigma})$, and $\phi(\tau, \vec{\sigma}) = e^{q(\tau, \vec{\sigma})/2} = \phi[r_{\bar a}(\tau, \vec{\sigma}), \pi_{\bar a}(\tau, \vec{\sigma})]$ is the conformal factor solution of the Lichnerowicz equation in this gauge. Since the linearization consists in taking only the terms quadratic in the DO's inside the weak ADM energy and only linear terms in the Lichnerowicz equation, we will do the following assumptions:

A) We assume $|r_{\bar a}(\tau, \vec{\sigma})| << 1$ on each WSW hyper-surface and $|\partial_u r_{\bar a}(\tau, \vec{\sigma})| \sim \frac{1}{L} O(r_{\bar a})$, $|\partial_u \partial_v r_{\bar a}(\tau, \vec{\sigma})| \sim \frac{1}{L^2} O(r_{\bar a})$, where L is a *big enough characteristic length interpretable as the reduced wavelength* $\lambda/2\pi$ *of the resulting gravitational waves*. Since the conjugate momenta $\pi_{\bar a}(\tau, \vec{\sigma})$ have the dimensions of $\frac{action}{L^3}$, i.e. of $\frac{k}{L}$ with $k = \frac{c^3}{16\pi G}$, we assume $|\pi_{\bar a}(\tau, \vec{\sigma})| \sim \frac{k}{L} O(r_{\bar a})$, $|\partial_u \pi_{\bar a}(\tau, \vec{\sigma})| \sim \frac{k}{L^2} O(r_{\bar a})$, $|\partial_u \partial_v \pi_{\bar a}(\tau, \vec{\sigma})| \sim \frac{k}{L^3} O(r_{\bar a})$. Therefore, $r_{\bar a}(\tau, \vec{\sigma})$ and $\pi_{\bar a}(\tau, \vec{\sigma})$ are *slowly varying over the length* L (for $r_{\bar a}, \pi_{\bar a} \rightarrow 0$ we get the void space-times of Ref. [3]). From Eq.(D7) we get that the Riemann tensor of our space-time is of order $\frac{1}{L\,k}\, O(\pi_{\bar a}) = \frac{1}{L^2}\, O(r_{\bar a}) \approx \mathcal{R}^{-2}$, where $\mathcal{R}$ is the mean radius of curvature. Therefore we get that the *requirements of the weak field approximation are satisfied*: i) $\mathcal{A} = O(r_{\bar a})$, if $\mathcal{A}$ is the amplitude of the gravitational wave; ii) $\frac{L}{\mathcal{R}} = O(r_{\bar a})$, namely $\frac{\lambda}{2\pi} << \mathcal{R}$.

B) We also assume $q(\tau, \vec{\sigma}) \sim O(r_{\bar a})$, $\partial_u q(\tau, \vec{\sigma}) \sim \frac{1}{L} O(r_{\bar a})$, $\partial_u \partial_v q(\tau, \vec{\sigma}) \sim \frac{1}{L^2} O(r_{\bar a})$, so that we get $\phi(\tau, \vec{\sigma}) = e^{q(\tau, \vec{\sigma})/2} \approx 1 + \frac{1}{2} q(\tau, \vec{\sigma}) + O(r^2_{\bar a})$ for the conformal factor. The Lichnerowicz equation becomes a linear partial differential equation for $q(\tau, \vec{\sigma})$. The linearization is done by systematically discarding terms of order $O(r^2_{\bar a})$ in the Lichnerowicz equation and $O(r^3_{\bar a})$ in the weak ADM energy.

With these positions we can, for the first time, *solve all the constraints* of tetrad gravity, Lichnerowicz equation included, and find explicitly the kernels in Eq.(1.4). We can check that our gauge is not a harmonic gauge. Notwithstanding this fact, the Hamilton equations imply the *wave equation* for the DO's $r_{\bar a}(\tau, \vec{\sigma})$. We can check explicitly that the linearized Einstein equations in this 4-coordinate system are satisfied. *Therefore, for the first time we get a definition of gravitational wave in a linearized post-Minkoskian Einstein space-time without introducing any background* and independently from the post-Newtonian approximation.

After a comparison with other approaches, in which it is emphasized the coordinate-dependent nature of effects like gravito-magnetism, we make a study of the time-like geodesics and of the geodesic deviation equation. The study of null geodesics allows to show the post-Minkowskian modification of the light-cone in each point of space-time. Then we analyze the embedding of the WSW hyper-surfaces of our gauge in the post-Minkowskian space-time and the associated congruence of time-like observers. Moreover, anticipating the introduction of matter to be done in a future paper, the comparison of our gauge with the post-Newtonian approach shows that it is the equation determining the bulk part $n(\tau, \vec{\sigma})$ of the lapse function which tends to the Poisson equation for the Newton potential for $c \rightarrow \infty$.

In Section II we solve the linearized Lichnerowicz equation and the rotation and super-momentum constraints in our gauge. In Section III we determine the quadratic part of the weak ADM energy in terms of the DO's and we evaluate the lapse and shift functions of our gauge. Then we find the linearized 4-metric and the tetrads of our gauge and we study the Landau-Lifschiz energy-momentum pseudo-tensor in our 4-coordinates. Section IV contains a comparison of our results with the Lichnerowicz-York conformal approach, with the standard harmonic gauges (and their associated gravito-electric-magnetic analogy) and with the post-Newtonian approximation. The Hamilton equations for the DO's and their solution, included the verification that Einstein's equations are satisfied, are presented in Section V. In Section VI we study the time-like geodesics and the geodesic deviation equation of our space-time. Section VII contains the determination of the embedding of the WSW hyper-surfaces of our gauge into the space-time and the study of the associated congruence of time-like observers and of the null geodesics. Final remarks and the future perspectives for the introduction of matter are drawn in the Conclusions.

In Appendix A we reproduce the results about tetrad gravity [3] needed for this paper. In Appendix B we give the determination of some kernels connected with the Shanmugadhasan transformation. Appendix C contains the material on the Fourier transform on the WSW hyper-surfaces of our gauge needed for the study of the Hamilton equations. Finally in Appendix D there is the linearized form of the relevant 3-tensors on the WSW hyper-surfaces of our gauge and of the relevant 4-tensors over our space-time. In this Appendix there is the explicit verification that Einstein's equations are satisfied.

II. LINEARIZATION IN THE 3-ORTHOGONAL GAUGE WITH $\pi_\phi(\tau, \vec{\sigma}) = 0$ AND SOLUTION OF THE CONSTRAINTS.

In this Section we study the linearization of tetrad gravity and the solution of its constraints in the completely fixed 3-orthogonal gauge with $\pi_\phi(\tau, \vec{\sigma}) = 0$ defined in the Introduction.

A. The Super-Hamiltonian Constraint.

Let us first consider the linearization of the *Lichnerowicz equation*, see Eq.(A1), which is the super-hamiltonian constraint of Eq.(A2), interpreted as an equation for the conformal factor of the 3-metric.

Since the super-hamiltonian constraint of Eq.(A2)becomes $\tilde{\mathcal{H}}_R(\tau, \vec{\sigma}) = \frac{\epsilon c^3}{16\pi G}\Big(-8\triangle\phi(\tau, \vec{\sigma}) + \frac{2}{\sqrt{3}}\sum_{\bar{a}u}\gamma_{\bar{a}u}\partial_u^2 r_{\bar{a}}(\tau, \vec{\sigma})\Big) + \frac{1}{L^2} O(r^2_{\bar{a}}) \approx 0$, where $\triangle = \vec{\partial}^2$ is the flat Laplacian $[\triangle \frac{1}{4\pi|\vec{\sigma}|} = -\delta^3(\vec{\sigma})]$, only the term $(-\tilde{\triangle} + \frac{1}{8}\, {}^3\tilde{R})\phi$ in Eq.(A1) gives a contribution of order $O(r_{\bar{a}})$. As a consequence the linearized Lichnerowicz equation for $\phi = e^{q/2} = 1 + \frac{1}{2} q + O(r^2_{\bar{a}})$ becomes

$$\triangle q(\tau, \vec{\sigma}) = \frac{1}{2\sqrt{3}} \sum_{u\bar{a}} \gamma_{\bar{a}u} \partial_u^2 r_{\bar{a}}(\tau, \vec{\sigma}) + \frac{1}{L^2} O(r^2_{\bar{a}}), \tag{2.1}$$

whose solution vanishing at spatial infinity is

$$q(\tau, \vec{\sigma}) = -\frac{1}{2\sqrt{3}} \sum_{u\bar{a}} \gamma_{\bar{a}u} \int d^3\sigma_1 \frac{\partial^2_{1u} r_{\bar{a}}(\tau, \vec{\sigma}_1)}{4\pi |\vec{\sigma} - \vec{\sigma}_1|} + O(r^2_{\bar{a}}). \tag{2.2}$$

B. The Rotation and Super-Momentum Constraints.

After having solved the super-hamiltonian constraint we have to solve the six rotation and super-momentum constraints. As said in the Introduction, this is equivalent [3] to find the kernels appearing in the linearization of Eqs.(1.4). After putting equal to zero the Abelianized rotation and super-momentum constraints in Eqs.(1.4), the old momenta ${}^3\check{\tilde{\pi}}^r_{(a)}(\tau, \vec{\sigma}) \overset{def}{=} {}^3\tilde{\pi}^r_{(a)}(\tau, \vec{\sigma})|_{3-O, \pi_\phi = 0}$ [14] are given by Eqs.(A4), whose linearization is

$$\begin{aligned}
{}^3\check{\tilde{\pi}}^r_{(a)}(\tau, \vec{\sigma}) &= \sqrt{3} \sum_{s\bar{a}} \gamma_{\bar{a}s} \int d^3\sigma_1 \tilde{\mathcal{K}}^r_{(a)s}(\vec{\sigma}, \vec{\sigma}_1, \tau | \phi, r_{\bar{a}}] \Big(\phi^{-2} e^{-\frac{1}{\sqrt{3}}\sum_{\bar{b}}\gamma_{\bar{b}s} r_{\bar{b}}} \pi_{\bar{a}} \Big)(\tau, \vec{\sigma}_1) = \\
&= \sqrt{3} \sum_{s\bar{a}} \gamma_{\bar{a}s} \int d^3\sigma_1 \mathcal{K}^{(o)r}_{(a)s}(\vec{\sigma}, \vec{\sigma}_1) \pi_{\bar{a}}(\tau, \vec{\sigma}_1) + \frac{k}{L^2} O(r^2_{\bar{a}}) = \\
&= \sqrt{3} \sum_{s\bar{a}} \gamma_{\bar{a}s} \int d^3\sigma_1 \Big[\delta^r_s \delta_{(a)s} \delta^3(\vec{\sigma}, \vec{\sigma}_1) + \mathcal{T}^{(o)r}_{(a)s}(\vec{\sigma}, \vec{\sigma}_1) \Big] \pi_{\bar{a}}(\tau, \vec{\sigma}_1) +
\end{aligned}$$

---

[14] $|_{3-0}$ means in the family of 3-orthogonal gauges.

$$+\frac{k}{L^2} O(r_{\bar{a}}^2),$$

with

$$\begin{aligned}
\mathcal{K}^{(o)r}_{(a)s}(\vec{\sigma},\vec{\sigma}_1) &= \tilde{\mathcal{K}}^{r}_{(a)s}(\vec{\sigma},\vec{\sigma}_1,\tau|1,0] = \delta^r_s \delta_{(a)s} \delta^3(\vec{\sigma},\vec{\sigma}_1) + \mathcal{T}^{(o)r}_{(a)s}(\vec{\sigma},\vec{\sigma}_1) = \\
&= \delta^r_s \delta_{(a)s} \delta^3(\vec{\sigma},\vec{\sigma}_1) - \frac{\partial G^{(o)rs}_{(a)}(\vec{\sigma},\vec{\sigma}_1)}{\partial \sigma^s_1}. \qquad (2.3)
\end{aligned}$$

Therefore, the linearized kernels $\mathcal{K}^{(o)r}_{(a)s}$'s are determined by the linearized kernels $G^{(o)ru}_{(a)}$'s.

As a consequence, the linearization implies that the partial differential equations (A5) for the kernels $G^{ru}_{(a)}$'s have to be restricted to $r_{\bar{a}}(\tau,\vec{\sigma}) = \pi_{\bar{a}}(\tau,\vec{\sigma}) = 0$ (zero curvature limit), namely the linearized kernels $G^{(o)ru}_{(a)}$'s are the same as for void space-times [3]. Their general solutions $G^{(o)ru}_{(a)}(\vec{\sigma},\vec{\sigma}_1)$'s will determine the $\mathcal{K}^{(o)r}_{(a)u}$'s [Eq.(2.3)] and the $F^{(o)r}_{(a)(b)}$'s [see Appendix B]. Moreover, to satisfy simultaneously the rotation and super-momentum constraints , the $\mathcal{K}^{(o)r}_{(a)u}$'s must satisfy the linearized version of Eqs.(A6). These linearized equations, which restrict the general solution, will be given in Eqs.(2.8) and are equivalent to the statement that the old momenta ${}^3\tilde{\pi}^r_{(a)}$ satisfy the linearized form of the six rotation and super-momentum constraints if we have

$$\begin{aligned}
&{}^3\check{\tilde{\pi}}^b_{(a)}(\tau,\vec{\sigma}) = {}^3\check{\tilde{\pi}}^a_{(b)}(\tau,\vec{\sigma}), \qquad a \neq b, \\
&\partial_r\, {}^3\check{\tilde{\pi}}^r_{(a)}(\tau,\vec{\sigma}) = 0. \qquad (2.4)
\end{aligned}$$

Eqs.(2.4) also correspond to the linearization of the three Einstein equations associated with the super-momentum constraints of ADM metric gravity.

The zero curvature limit of Eqs.(A5) implies that the $G^{(o)ru}_{(a)}$'s are determined by the following linear partial differential equations [15]

$$\begin{aligned}
&1)\ s = a \quad \textit{homogeneous equations}: \\
&\frac{\partial G^{(o)a2}_{(a)}(\vec{\sigma},\vec{\sigma}_1)}{\partial \sigma^1_1} + \frac{\partial G^{(o)a1}_{(a)}(\vec{\sigma},\vec{\sigma}_1)}{\partial \sigma^2_1} = \\
&= \frac{\partial G^{(o)a3}_{(a)}(\vec{\sigma},\vec{\sigma}_1)}{\partial \sigma^2_1} + \frac{\partial G^{(o)a2}_{(a)}(\vec{\sigma},\vec{\sigma}_1)}{\partial \sigma^3_1} = \\
&= \frac{\partial G^{(o)a1}_{(a)}(\vec{\sigma},\vec{\sigma}_1)}{\partial \sigma^3_1} + \frac{\partial G^{(o)a3}_{(a)}(\vec{\sigma},\vec{\sigma}_1)}{\partial \sigma^1_1} = 0, \qquad a = 1,2,3;
\end{aligned}$$

---

[15] $r_1, r_2 \neq b,\ r_1 \neq r_2;\ \epsilon_{(a)(r_1)(b)} = -\delta_{(a)r_2}\epsilon_{(r_1)(r_2)(b)},\ \epsilon_{(a)(r_2)(b)} = \delta_{(a)r_1}\epsilon_{(r_1)(r_2)(b)},\ \epsilon_{(u)(r_1)(b)} = -\delta_{(u)r_2}\epsilon_{(r_1)(r_2)(b)},\ \epsilon_{(u)(r_2)(b)} = \delta_{(u)r_1}\epsilon_{(r_1)(r_2)(b)}.$

$$\Downarrow$$

$$\frac{\partial^2 G^{(o)a1}_{(a)}(\vec{\sigma},\vec{\sigma}_1)}{\partial\sigma^2_1\partial\sigma^3_1} = \frac{\partial^2 G^{(o)a2}_{(a)}(\vec{\sigma},\vec{\sigma}_1)}{\partial\sigma^3_1\partial\sigma^1_1} = \frac{\partial^2 G^{(o)a3}_{(a)}(\vec{\sigma},\vec{\sigma}_1)}{\partial\sigma^1_1\partial\sigma^3_1} = 0,$$

$$2)\; s \neq a\;\; s \neq r,\; r \neq a \quad inhomogeneous\, equations:$$

$$\begin{aligned}
&\frac{\partial G^{(o)sr}_{(a)}(\vec{\sigma},\vec{\sigma}_1)}{\partial\sigma^s_1} + \frac{\partial G^{(o)ss}_{(a)}(\vec{\sigma},\vec{\sigma}_1)}{\partial\sigma^r_1} = \\
&= \frac{\partial G^{(o)sa}_{(a)}(\vec{\sigma},\vec{\sigma}_1)}{\partial\sigma^r_1} + \frac{\partial G^{(o)sr}_{(a)}(\vec{\sigma},\vec{\sigma}_1)}{\partial\sigma^a_1} = 0, \\
&\frac{\partial G^{(o)ss}_{(a)}(\vec{\sigma},\vec{\sigma}_1)}{\partial\sigma^a_1} + \frac{\partial G^{(o)sa}_{(a)}(\vec{\sigma},\vec{\sigma}_1)}{\partial\sigma^s_1} = \delta^3(\vec{\sigma};\vec{\sigma}_1).
\end{aligned} \tag{2.5}$$

Each set of homogeneous equations, considered as equations for functions of $\vec{\sigma}$, is of the form $\partial_2\, u_{(1)}(\vec{\sigma}) + \partial_1\, u_{(2)}(\vec{\sigma}) = \partial_3\, u_{(2)}(\vec{\sigma}) + \partial_2\, u_{(3)}(\vec{\sigma}) = \partial_1\, u_{(3)}(\vec{\sigma}) + \partial_3\, u_{(1)}(\vec{\sigma}) = 0$. This is a system of three linear partial differential equations for the three unknown functions $u_{(i)}(\vec{\sigma})$ of *elliptic* type, since the determinant of its characteristic matrix [25] is $2\xi_1\xi_2\xi_3 \neq 0$. Moreover it is integrable, since $u_{(r)}(\vec{\sigma}) = f_{(r)}(\sigma^r)$ with arbitrary $f_{(r)}$ are solutions of the system. We do not know whether they exhaust all the possible solutions. Therefore, $G^{(o)ar}_{(a)}(\vec{\sigma},\vec{\sigma}_1) = h^{(o)ar}_{(a)}(\vec{\sigma},\sigma^r_1)$, with $h^{(o)ar}_{(a)}$ arbitrary functions, are solutions of the homogeneous equations.

As a consequence, if $\bar{G}^{(o)ru}_{(a)}(\vec{\sigma},\vec{\sigma}_1)$, $r \neq a$, is a particular solution of each set of inhomogeneous equations (2.5), then the general solution is $G^{(o)ru}_{(a)}(\vec{\sigma},\vec{\sigma}_1) = \bar{G}^{(o)ru}_{(a)}(\vec{\sigma},\vec{\sigma}_1) + g^{(o)ru}_{(a)}(\vec{\sigma},\vec{\sigma}_1)$, $r \neq a$, with the $g^{(o)ru}_{(a)}$'s arbitrary homogeneous solutions (again with $g^{(o)ru}_{(a)}(\vec{\sigma},\sigma^r_1)$, if this is the most general solution of the associated homogeneous equations).

Therefore the general solution of Eqs.(2.5) for the kernels $G^{(o)ru}_{(a)}(\vec{\sigma},\vec{\sigma}_1)$ can be written in the following form

$$G^{(o)ru}_{(a)}(\vec{\sigma},\vec{\sigma}_1) = \delta^r_{(a)} h^{(o)au}_{(a)}(\vec{\sigma},\vec{\sigma}_1) + (1-\delta^r_{(a)})[\bar{G}^{(o)ru}_{(a)}(\vec{\sigma},\vec{\sigma}_1) + g^{(o)ru}_{(a)}(\vec{\sigma},\vec{\sigma}_1)], \tag{2.6}$$

with arbitrary $h^{(o)au}_{(a)}$'s and $g^{(o)ru}_{(a)}$'s. Then, Eq(2.3) gives the following expression for the kernels $\mathcal{K}^{(o)r}_{(a)u}$'s

$$\begin{aligned}
&\mathcal{K}^{(o)r}_{(a)u}(\vec{\sigma},\vec{\sigma}_1) = \delta^r_{(a)}\delta_{(a)u}\delta^3(\vec{\sigma},\vec{\sigma}_1) - \\
&- \Big(\delta^r_{(a)}\frac{\partial h^{(o)au}_{(a)}(\vec{\sigma},\vec{\sigma}_1)}{\partial\sigma^u_1} + (1-\delta^r_{(a)})[\frac{\partial\bar{G}^{(o)ru}_{(a)}(\vec{\sigma},\vec{\sigma}_1)}{\partial\sigma^u_1} + \frac{\partial g^{(o)ru}_{(a)}(\vec{\sigma},\vec{\sigma}_1)}{\partial\sigma^u_1}]\Big).
\end{aligned} \tag{2.7}$$

The solutions of Eqs.(2.5) for the $G^{(o)ru}_{(a)}$'s are restricted by the requirement that the $\mathcal{K}^{(o)r}_{(a)u}$'s of Eqs.(2.7) satisfy the zero curvature limit of Eqs. (A6), which in the 3-orthogonal gauges become [16]

$$\sum_r \Big[\delta_{(a)r}\mathcal{K}^{(o)r}_{(b)u} - \delta_{(b)r}\mathcal{K}^{(o)r}_{(a)u}\Big](\vec{\sigma},\vec{\sigma}_1) = \Big[\frac{\partial G^{(o)au}_{(b)}(\vec{\sigma},\vec{\sigma}_1;\tau)}{\partial \sigma_1^u} - \frac{\partial G^{(o)bu}_{(a)}(\vec{\sigma},\vec{\sigma}_1;\tau)}{\partial \sigma_1^u}\Big] = 0, \qquad a \neq b,$$

$$\partial_r\, \mathcal{K}^{(o)r}_{(a)u}(\vec{\sigma},\vec{\sigma}_1) = \partial_r\, \Big[\delta^r_{(a)}\delta_{(a)u}\delta^3(\vec{\sigma},\vec{\sigma}_1) - \frac{\partial G^{(o)ru}_{(a)}(\vec{\sigma},\vec{\sigma}_1;\tau)}{\partial \sigma_1^u}\Big] = 0. \tag{2.8}$$

The first set of Eqs.(2.8) becomes the following set of three linear partial differential equations to get the $g^{(o)ru}_{(a)}$'s of Eq.(2.6) with $a \neq b$ in terms of the $\bar{G}^{(o)ru}_{(a)}$'s

$$\begin{aligned}\frac{\partial f^u_{ab}(\vec{\sigma},\vec{\sigma}_1)}{\partial \sigma^u} &= \frac{\partial}{\partial \sigma_1^u}\Big(g^{(o)au}_{(b)}(\vec{\sigma},\vec{\sigma}_1) - g^{(o)bu}_{(a)}(\vec{\sigma},\vec{\sigma}_1)\Big) = \\ &= -\Big(\frac{\partial \bar{G}^{(o)au}_{(b)}(\vec{\sigma},\vec{\sigma}_1)}{\partial \sigma_1^u} - \frac{\partial \bar{G}^{(o)bu}_{(a)}(\vec{\sigma},\vec{\sigma}_1)}{\partial \sigma_1^u}\Big) \overset{def}{=} m^u_{ab}(\vec{\sigma}_1,\vec{\sigma}).\end{aligned} \tag{2.9}$$

For each pair $a \neq b$, this is a system of three elliptic linear partial differential equations for the f^u_{ab}'s. Each choice of the $g^{(o)au}_{(b)}$'s, $a \neq b$, which gives a solution of this system, implies that the associated kernels $\mathcal{K}^{(o)r}_{(a)u}$'s satisfy the rotation constraints.

Having found a solution for the $g^{(o)au}_{(b)}$'s, $a \neq b$, the second set of Eqs.(2.8) becomes the following set of equations for the $h^{(o)au}_{(a)}$'s of Eq.(2.6) in terms of the $\bar{G}^{(o)ru}_{(a)}$'s and $g^{(o)ru}_{(a)}$'s

$$\begin{aligned}\sum_r \partial_r \delta^r_{(a)}\Big[\frac{\partial h^{(o)au}_{(a)}(\vec{\sigma},\vec{\sigma}_1)}{\partial \sigma_1^u}\Big] &= \sum_r \partial_r\, \Big[\delta^r_{(a)}\delta_{(a)u}\delta^3(\vec{\sigma},\vec{\sigma}_1) - \\ -(1-\delta^r_{(a)})\Big[&\frac{\partial \bar{G}^{(o)ru}_{(a)}(\vec{\sigma},\vec{\sigma}_1)}{\partial \sigma_1^u} + \frac{\partial g^{(o)ru}_{(a)}(\vec{\sigma},\vec{\sigma}_1)}{\partial \sigma_1^u}\Big]\Big)\Big].\end{aligned} \tag{2.10}$$

By using the linearization of the Green function of the covariant divergence given in Eqs.(A8) and (A9), we get ($f^{su}_{(T)}$ are solutions of the homogeneous equation)

$$\begin{aligned}\frac{\partial h^{(o)su}_{(s)}(\vec{\sigma},\vec{\sigma}_1)}{\partial \sigma_1^u} &= f^{su}_{(T)}(\vec{\sigma},\vec{\sigma}_1) - \\ &- \int d^3\sigma_2 \sum_{(a)} \delta_{(s)(a)} c^s(\vec{\sigma},\vec{\sigma}_2) \sum_r \frac{\partial}{\partial \sigma_2^r}\Big[\delta^r_{(a)}\delta_{(a)u}\delta^3(\vec{\sigma}_2,\vec{\sigma}_1) - \\ &-(1-\delta^r_{(a)})\Big[\frac{\partial \bar{G}^{(o)ru}_{(a)}(\vec{\sigma}_2,\vec{\sigma}_1)}{\partial \sigma_1^u} + \frac{\partial g^{(o)ru}_{(a)}(\vec{\sigma}_2,\vec{\sigma}_1)}{\partial \sigma_1^u}\Big]\Big].\end{aligned} \tag{2.11}$$

Again this is a system of elliptic linear partial differential equations for the $h^{(o)au}_{(a)}$'s with fixed a.

---

[16] At zeroth order ($\phi(\tau,\vec{\sigma}) = 1$) in the 3-orthogonal gauges the spin connection vanishes, ${}^3\hat{\omega}_{r(a)}(\tau,\vec{\sigma}) = 0 + O(r_{\bar{a}})$, see Eq.(A7).

After having found the solutions of Eqs. (2.5), (2.9) and (2.11) for the kernels $G^{(o)ru}_{(a)}$'s, $\mathcal{K}^{(o)r}_{(a)u}$'s and $F^{(o)r}_{(a)(b)}$'s, every remaining arbitrariness will be fixed by the boundary conditions at spatial infinity for the momenta ${}^3\tilde{\pi}^r_{(a)}(\tau,\vec{\sigma})$, given in Eqs.(1.2). The final solutions are equivalent not only to the solution of the rotation and diffeomorphisms constraints, but also to their Abelianization in the 3-orthogonal gauges with $\alpha_{(a)}(\tau,\vec{\sigma})=0$.

C. A Solution of the Rotation and Super-Momentum Constraints.

We have found the following particular solution $G^{(o)ru}_{(a)} = \bar{G}^{(o)ru}_{(a)}$ of Eqs.(2.5) with $g^{(o)au}_{(b)} = h^{(o)au}_{(a)} = 0$ and vanishing for $|\vec{\sigma}| \rightarrow \infty$ (we use $\delta(\sigma^r, \sigma^r_1 \rightarrow -\infty) = 0$ for finite σ^r),

$$G^{(o)ar}_{(a)}(\vec{\sigma},\vec{\sigma}_1) = 0, \qquad a = 1,2,3,$$

$$\begin{aligned}
G^{(o)21}_{(1)}(\vec{\sigma},\vec{\sigma}_1) &= G^{(o)11}_{(2)}(\vec{\sigma},\vec{\sigma}_1) = G^{(o)33}_{(2)}(\vec{\sigma},\vec{\sigma}_1) = G^{(o)23}_{(3)}(\vec{\sigma},\vec{\sigma}_1) = \\
&= \frac{1}{2}\int_{-\infty}^{\sigma^2_1} dw^2_1\, \delta^3(\vec{\sigma}, \sigma^1_1 w^2_1 \sigma^3_1) = \frac{1}{2}\, \delta(\sigma^1,\sigma^1_1)\, \theta(\sigma^2_1,\sigma^2)\, \delta(\sigma^3,\sigma^3_1),
\end{aligned}$$

$$\begin{aligned}
G^{(o)22}_{(1)}(\vec{\sigma},\vec{\sigma}_1) &= G^{(o)12}_{(2)}(\vec{\sigma},\vec{\sigma}_1) = G^{(o)33}_{(1)}(\vec{\sigma},\vec{\sigma}_1) = G^{(o)13}_{(3)}(\vec{\sigma},\vec{\sigma}_1) = \\
&= \frac{1}{2}\int_{-\infty}^{\sigma^1_1} dw^1_1\, \delta^3(\vec{\sigma}, w^1_1 \sigma^2_1 \sigma^3_1) = \frac{1}{2}\, \theta(\sigma^1_1,\sigma^1)\, \delta(\sigma^2,\sigma^2_1)\, \delta(\sigma^3,\sigma^3_1),
\end{aligned}$$

$$\begin{aligned}
G^{(o)31}_{(1)}(\vec{\sigma},\vec{\sigma}_1) &= G^{(o)11}_{(3)}(\vec{\sigma},\vec{\sigma}_1) = G^{(o)32}_{(2)}(\vec{\sigma},\vec{\sigma}_1) = G^{(o)22}_{(3)}(\vec{\sigma},\vec{\sigma}_1) = \\
&= \frac{1}{2}\int_{-\infty}^{\sigma^3_1} dw^3_1\, \delta^3(\vec{\sigma}, \sigma^1_1 \sigma^2_1 w^3_1) = \frac{1}{2}\, \delta(\sigma^1,\sigma^1_1)\, \delta(\sigma^2,\sigma^2_1)\, \theta(\sigma^3_1,\sigma^3),
\end{aligned}$$

$$\begin{aligned}
G^{(o)23}_{(1)}(\vec{\sigma},\vec{\sigma}_1) &= G^{(o)13}_{(2)}(\vec{\sigma},\vec{\sigma}_1) = -\frac{1}{2}\frac{\partial}{\partial \sigma^3_1}\int_{-\infty}^{\sigma^1_1} dw^1_1 \int_{-\infty}^{\sigma^2_1} dw^2_1\, \delta^3(\vec{\sigma}, w^1_1 w^2_1 \sigma^3_1) = \\
&= -\frac{1}{2}\, \theta(\sigma^1_1,\sigma^1)\, \theta(\sigma^2_1,\sigma^2)\, \frac{\partial \delta(\sigma^3,\sigma^3_1)}{\partial \sigma^3_1},
\end{aligned}$$

$$\begin{aligned}
G^{(o)32}_{(1)}(\vec{\sigma},\vec{\sigma}_1) &= G^{(o)12}_{(3)}(\vec{\sigma},\vec{\sigma}_1) = -\frac{1}{2}\,\frac{\partial}{\partial \sigma^2_1}\int_{-\infty}^{\sigma^1_1} dw^1_1 \int_{-\infty}^{\sigma^3_1} dw^3_1\, \delta^3(\vec{\sigma}, w^1_1 \sigma^2_1 w^3_1) = \\
&= -\frac{1}{2}\, \theta(\sigma^1_1,\sigma^1)\, \frac{\partial \delta(\sigma^2,\sigma^2_1)}{\partial \sigma^2_1}\, \theta(\sigma^3_1,\sigma^3),
\end{aligned}$$

$$\begin{aligned}
G^{(o)31}_{(2)}(\vec{\sigma},\vec{\sigma}_1) &= G^{(o)21}_{(3)}(\vec{\sigma},\vec{\sigma}_1) = -\frac{1}{2}\,\frac{\partial}{\partial \sigma^1_1}\int_{-\infty}^{\sigma^2_1} dw^2_1 \int_{-\infty}^{\sigma^3_1} dw^3_1\, \delta^3(\vec{\sigma}, \sigma^1_1 w^2_1 w^3_1) = \\
&= -\frac{1}{2}\, \frac{\partial \delta(\sigma^1,\sigma^1_1)}{\partial \sigma^1_1}\, \theta(\sigma^2_1,\sigma^2)\, \theta(\sigma^3_1,\sigma^3),
\end{aligned}$$

or

$$
\begin{aligned}
G^{(o)ru}_{(a)}(\vec{\sigma},\vec{\sigma}_1) &= -\frac{1}{2}\,[1-\delta^r_{(a)}]\,[1-2(\delta_{ru}+\delta_{au})]\\
&\quad \frac{\partial}{\partial\sigma_1^u}\int_{-\infty}^{\sigma_1^r} dw_1^r \int_{-\infty}^{\sigma_1^a} dw_1^a\, \delta^3(\vec{\sigma}, w_1^r w_1^a \sigma_1^{k\neq(r,a)}) =\\
&= -\frac{1}{2}\,[1-\delta^r_{(a)}]\,[1-2(\delta_{ru}+\delta_{au})]\\
&\quad \frac{\partial}{\partial\sigma_1^u}\Big[\theta(\sigma_1^r,\sigma^r)\,\theta(\sigma_1^a,\sigma^a)\,\delta(\sigma^{k\neq(r,a)},\sigma_1^{k\neq(r,a)})\Big].
\end{aligned}
\tag{2.12}
$$

where $\theta(x)$ is the step function $[\frac{d}{dx}\theta(x)=\delta(x)]$. This implies the following expression for the kernels $\mathcal{K}^{(o)r}_{(a)u}$'s

$$
\mathcal{K}^{(o)r}_{(a)u}(\vec{\sigma},\vec{\sigma}_1) = \delta^r_u\delta_{(a)u}\delta^3(\vec{\sigma},\vec{\sigma}_1) + \mathcal{T}^{(o)r}_{(a)u}(\vec{\sigma},\vec{\sigma}_1) = \delta^r_{(a)}\delta_{(a)u}\delta^3(\vec{\sigma},\vec{\sigma}_1) - \frac{\partial G^{(o)ru}_{(a)}(\vec{\sigma},\vec{\sigma}_1)}{\partial\sigma_1^u},
$$

$$
with
$$

$$
\mathcal{T}^{(o)a}_{(a)u}(\vec{\sigma},\vec{\sigma}_1) = 0, \qquad a=1,2,3,
$$

$$
\begin{aligned}
\mathcal{T}^{(o)2}_{(1)1}(\vec{\sigma},\vec{\sigma}_1) &= \mathcal{T}^{(o)1}_{(2)1}(\vec{\sigma},\vec{\sigma}_1) = -\frac{1}{2}\,\frac{\partial}{\partial\sigma_1^1}\int_{-\infty}^{\sigma_1^2} dw_1^2\,\delta^3(\vec{\sigma},\sigma_1^1 w_1^2 \sigma_1^3) =\\
&= -\frac{1}{2}\,\frac{\partial\delta(\sigma^1,\sigma_1^1)}{\partial\sigma_1^1}\,\theta(\sigma_1^2,\sigma^2)\,\delta(\sigma^3,\sigma_1^3) =\\
&= -\frac{1}{2}\,\frac{\partial^2}{(\partial\sigma_1^1)^2}\int_{-\infty}^{\sigma_1^1} dw_1^1 \int_{-\infty}^{\sigma_1^2} dw_1^2\,\delta^3(\vec{\sigma},w_1^1 w_1^2 \sigma_1^3),\\
\mathcal{T}^{(o)2}_{(1)2}(\vec{\sigma},\vec{\sigma}_1) &= \mathcal{T}^{(o)1}_{(2)2}(\vec{\sigma},\vec{\sigma}_1) = -\frac{1}{2}\,\frac{\partial}{\partial\sigma_1^2}\int_{-\infty}^{\sigma_1^1} dw_1^1\,\delta^3(\vec{\sigma},w_1^1 \sigma_1^2 \sigma_1^3) =\\
&= -\frac{1}{2}\,\theta(\sigma_1^1,\sigma^1)\,\frac{\partial\delta(\sigma^2,\sigma_1^2)}{\partial\sigma_1^2}\,\delta(\sigma^3,\sigma_1^3) =\\
&= -\frac{1}{2}\,\frac{\partial^2}{(\partial\sigma_1^2)^2}\int_{-\infty}^{\sigma_1^1} dw_1^1 \int_{-\infty}^{\sigma_1^2} dw_1^2\,\delta^3(\vec{\sigma},w_1^1 w_1^2 \sigma_1^3),\\
\mathcal{T}^{(o)2}_{(1)3}(\vec{\sigma},\vec{\sigma}_1) &= \mathcal{T}^{(o)1}_{(2)3}(\vec{\sigma},\vec{\sigma}_1) = \frac{1}{2}\,\frac{\partial^2}{(\partial\sigma_1^3)^2}\int_{-\infty}^{\sigma_1^1} dw_1^1 \int_{-\infty}^{\sigma_1^2} dw_1^2\,\delta^3(w_1^1 w_1^2 \sigma_1^3) =\\
&= \frac{1}{2}\,\theta(\sigma_1^1,\sigma^1)\,\theta(\sigma_1^2,\sigma^2)\,\frac{\partial^2\delta(\sigma^3,\sigma_1^3)}{(\partial\sigma_1^3)^2},\\
\mathcal{T}^{(o)3}_{(1)1}(\vec{\sigma},\vec{\sigma}_1) &= \mathcal{T}^{(o)1}_{(3)1}(\vec{\sigma},\vec{\sigma}_1) = -\frac{1}{2}\,\frac{\partial}{\partial\sigma_1^1}\int_{-\infty}^{\sigma_1^3} dw_1^3\,\delta^3(\vec{\sigma},\sigma_1^1 \sigma_1^2 w_1^3) =\\
&= -\frac{1}{2}\,\frac{\partial\delta(\sigma^1,\sigma_1^1)}{\partial\sigma_1^1}\,\delta(\sigma^2,\sigma_1^2)\,\theta(\sigma_1^3,\sigma^3) =\\
&= -\frac{1}{2}\,\frac{\partial^2}{(\partial\sigma_1^1)^2}\int_{-\infty}^{\sigma_1^1} dw_1^1 \int_{-\infty}^{\sigma_1^3} dw_1^3\,\delta^3(\vec{\sigma},w_1^1 \sigma_1^2 w_1^3),
\end{aligned}
$$

$$
\begin{aligned}
\mathcal{T}^{(o)3}_{(1)2}(\vec{\sigma},\vec{\sigma}_1) &= \mathcal{T}^{(o)1}_{(3)2}(\vec{\sigma},\vec{\sigma}_1) = \frac{1}{2}\,\frac{\partial^2}{(\partial\sigma_1^2)^2}\,\int_{-\infty}^{\sigma_1^1} dw_1^1 \int_{-\infty}^{\sigma_1^3} dw_1^3\,\delta^3(w_1^1\sigma_1^2w_1^3) = \\
&= \frac{1}{2}\,\theta(\sigma_1^1,\sigma^1)\,\frac{\partial^2\delta(\sigma^2,\sigma_1^2)}{(\partial\sigma_1^2)^2}\,\theta(\sigma_1^3,\sigma^3),\\
\\
\mathcal{T}^{(o)3}_{(1)3}(\vec{\sigma},\vec{\sigma}_1) &= \mathcal{T}^{(o)1}_{(3)3}(\vec{\sigma},\vec{\sigma}_1) = -\frac{1}{2}\,\frac{\partial}{\partial\sigma_1^3}\,\int_{-\infty}^{\sigma_1^1} dw_1^1\,\delta^3(\vec{\sigma},w_1^1\sigma_1^2\sigma_1^3) = \\
&= -\frac{1}{2}\,\theta(\sigma_1^1,\sigma^1)\,\delta(\sigma^2,\sigma_1^2)\,\frac{\partial\delta(\sigma^3,\sigma_1^3)}{\partial\sigma_1^3} = \\
&= -\frac{1}{2}\,\frac{\partial^2}{(\partial\sigma_1^3)^2}\,\int_{-\infty}^{\sigma_1^1} dw_1^1 \int_{-\infty}^{\sigma_1^3} dw_1^3\,\delta^3(\vec{\sigma},w_1^1\sigma_1^2w_1^3),\\
\\
\mathcal{T}^{(o)2}_{(3)1}(\vec{\sigma},\vec{\sigma}_1) &= \mathcal{T}^{(o)3}_{(2)1}(\vec{\sigma},\vec{\sigma}_1) = \frac{1}{2}\,\frac{\partial^2}{(\partial\sigma_1^1)^2}\int_{-\infty}^{\sigma_1^2} dw_1^2 \int_{-\infty}^{\sigma_1^3} dw_1^3\,\delta^3(\vec{\sigma},\sigma_1^1w_1^2w_1^3) = \\
&= \frac{1}{2}\,\frac{\partial^2\delta(\sigma^1,\sigma_1^1)}{(\partial\sigma_1^1)^2}\,\theta(\sigma_1^2,\sigma^2)\,\theta(\sigma_1^3,\sigma^3),\\
\\
\mathcal{T}^{(o)2}_{(3)2}(\vec{\sigma},\vec{\sigma}_1) &= \mathcal{T}^{(o)3}_{(2)2}(\vec{\sigma},\vec{\sigma}_1) = -\frac{1}{2}\,\frac{\partial}{\partial\sigma_1^2}\,\int_{-\infty}^{\sigma_1^3} dw_1^3\,\delta^3(\vec{\sigma},\sigma_1^1\sigma_1^2w_1^3) = \\
&= -\frac{1}{2}\,\delta(\sigma^1,\sigma_1^1)\,\frac{\partial\delta(\sigma^2,\sigma_1^2)}{\partial\sigma_1^2}\,\theta(\sigma_1^3,\sigma^3) = \\
&= -\frac{1}{2}\,\frac{\partial^2}{(\partial\sigma_1^2)^2}\,\int_{-\infty}^{\sigma_1^2} dw_1^2 \int_{-\infty}^{\sigma_1^3} dw_1^3\,\delta^3(\vec{\sigma},\sigma_1^1w_1^2w_1^3),\\
\\
\mathcal{T}^{(o)2}_{(3)3}(\vec{\sigma},\vec{\sigma}_1) &= \mathcal{T}^{(o)3}_{(2)3}(\vec{\sigma},\vec{\sigma}_1) = -\frac{1}{2}\,\frac{\partial}{\partial\sigma_1^3}\int_{-\infty}^{\sigma_1^2} dw_1^2\,\delta^3(\vec{\sigma},\sigma_1^1w_1^2\sigma_1^3) = \\
&= -\frac{1}{2}\,\delta(\sigma^1,\sigma_1^1)\,\theta(\sigma_1^2,\sigma^2)\,\frac{\partial\delta(\sigma^3,\sigma_1^3)}{\partial\sigma_1^3} = \\
&= -\frac{1}{2}\,\frac{\partial^2}{(\partial\sigma_1^3)^2}\,\int_{-\infty}^{\sigma_1^2} dw_1^2 \int_{-\infty}^{\sigma_1^3} dw_1^2\,\delta^3(\vec{\sigma},\sigma_1^1w_1^2w_1^3),\\
\\
&\quad or\\
\\
\mathcal{T}^{(o)r}_{(a)u}(\vec{\sigma},\vec{\sigma}_1) &= \frac{1}{2}\,[1-\delta^r_{(a)}]\,[1-2(\delta_{ru}+\delta_{au})]\\
&\quad \frac{\partial^2}{(\partial\sigma_1^u)^2}\,\int_{-\infty}^{\sigma_1^r} dw_1^r \int_{-\infty}^{\sigma_1^a} dw_1^a\,\delta^3(\vec{\sigma},w_1^rw_1^a\sigma_1^{k\neq(r,a)}) = \\
&= \frac{1}{2}\,[1-\delta^r_{(a)}]\,[1-2(\delta_{ru}+\delta_{au})]\\
&\quad \frac{\partial^2}{(\partial\sigma_1^u)^2}\,\Big[\theta(\sigma_1^r,\sigma^r)\,\theta(\sigma_1^a,\sigma^a)\,\delta(\sigma^{k\neq(r,a)},\sigma_1^{k\neq(r,a)})\Big]. \qquad (2.13)
\end{aligned}
$$

The kernels $F^{(o)r}_{(a)(b)}$'s are given in Appendix B.

D. The Old Cotriad Momenta from the Solution of the Constraints.

Eqs.(2.3) imply that the cotriad momenta, solution of both the linearized rotation and super-momentum constraints, have the following expression in terms of the DO momenta $\pi_{\bar a}(\tau, \vec \sigma)$

$$
\begin{aligned}
{}^3\check{\tilde\pi}^1_{(1)}(\tau, \vec \sigma) &= \sqrt{3} \sum_{\bar a} \gamma_{\bar a 1} \pi_{\bar a}(\tau, \vec \sigma),\\
{}^3\check{\tilde\pi}^2_{(1)}(\tau, \vec \sigma) &= -\frac{\sqrt{3}}{2} \sum_{\bar a} \Big[\gamma_{\bar a 1} \int d^3\sigma_1 \frac{\partial \delta(\sigma^1, \sigma^1_1)}{\partial \sigma^1_1} \theta(\sigma^2_1, \sigma^2) \delta(\sigma^3, \sigma^3_1) \pi_{\bar a}(\tau, \vec \sigma_1) +\\
&+ \gamma_{\bar a 2} \int d^3\sigma_1 \theta(\sigma^1_1, \sigma^1) \frac{\partial \delta(\sigma^2, \sigma^2_1)}{\partial \sigma^2_1} \delta(\sigma^3, \sigma^3_1) \pi_{\bar a}(\tau, \vec \sigma_1) -\\
&- \gamma_{\bar a 3} \int d^3\sigma_1 \theta(\sigma^1_1, \sigma^1) \theta(\sigma^2_1, \sigma^2) \frac{\partial^2 \delta(\sigma^3, \sigma^3_1)}{(\partial \sigma^3_1)^2} \pi_{\bar a}(\tau, \vec \sigma_1)\Big] =\\
&= \frac{\sqrt{3}}{2} \sum_{\bar a} \Big[\gamma_{\bar a 1} \int_{\sigma^2}^{\infty} d\sigma^2_1 \frac{\partial \pi_{\bar a}(\tau, \sigma^1 \sigma^2_1 \sigma^3)}{\partial \sigma^1} +\\
&+ \gamma_{\bar a 2} \int_{\sigma^1}^{\infty} d\sigma^1_1 \frac{\partial \pi_{\bar a}(\tau, \sigma^1_1 \sigma^2 \sigma^3)}{\partial \sigma^2} +\\
&- \gamma_{\bar a 3} \int_{\sigma^1}^{\infty} d\sigma^1_1 \int_{\sigma^2}^{\infty} d\sigma^2_1 \frac{\partial^2 \pi_{\bar a}(\tau, \sigma^1_1 \sigma^2_1 \sigma^3)}{(\partial \sigma^3)^2}\Big],\\
{}^3\check{\tilde\pi}^3_{(1)}(\tau, \vec \sigma) &= -\frac{\sqrt{3}}{2} \sum_{\bar a} \Big[\gamma_{\bar a 1} \int d^3\sigma_1 \frac{\partial \delta(\sigma^1, \sigma^1_1)}{\partial \sigma^1_1} \delta(\sigma^2, \sigma^2_1) \theta(\sigma^3_1, \sigma^3) \pi_{\bar a}(\tau, \vec \sigma_1) -\\
&- \gamma_{\bar a 2} \int d^3\sigma_1 \theta(\sigma^1_1, \sigma^1) \delta(\sigma^2, \sigma^2_1) \frac{\partial \delta(\sigma^3, \sigma^3_1)}{\partial \sigma^3_1} \pi_{\bar a}(\tau, \vec \sigma_1)\Big] =\\
&= -\frac{\sqrt{3}}{2} \sum_{\bar a} \Big[\gamma_{\bar a 1} \int_{\sigma^3}^{\infty} d\sigma^3_1 \frac{\partial \pi_{\bar a}(\tau, \sigma^1 \sigma^2 \sigma^3_1)}{\partial \sigma^1} +\\
&+ \gamma_{\bar a 2} \int_{\sigma^1}^{\infty} d\sigma^1_1 \int_{\sigma^3}^{\infty} d\sigma^3_1 \frac{\partial^2 \pi_{\bar a}(\tau, \sigma^1_1 \sigma^2 \sigma^3_1)}{(\partial \sigma^2)^2} +\\
&+ \gamma_{\bar a 3} \int_{\sigma^1}^{\infty} d\sigma^1_1 \frac{\partial \pi_{\bar a}(\tau, \sigma^1_1 \sigma^2 \sigma^3)}{\partial \sigma^3}\Big],
\end{aligned}
$$

$$
\begin{aligned}
{}^3\check{\tilde\pi}^1_{(2)}(\tau, \vec \sigma) &= {}^3\check{\tilde\pi}^2_{(1)}(\tau, \vec \sigma),\\
{}^3\check{\tilde\pi}^2_{(2)}(\tau, \vec \sigma) &= \sqrt{3} \sum_{\bar a} \gamma_{\bar a 2} \pi_{\bar a}(\tau, \vec \sigma),\\
{}^3\check{\tilde\pi}^3_{(2)}(\tau, \vec \sigma) &= -\frac{\sqrt{3}}{2} \sum_{\bar a} \Big[\gamma_{\bar a 1} \int d^3\sigma_1 \frac{\partial^2 \delta(\sigma^1, \sigma^1_1)}{(\partial \sigma^1_1)^2} \theta(\sigma^2_1, \sigma^2) \theta(\sigma^3_1, \sigma^3) \pi_{\bar a}(\tau, \vec \sigma_1) +\\
&+ \gamma_{\bar a 2} \int d^3\sigma_1 \delta(\sigma^1, \sigma^1_1) \frac{\partial \delta(\sigma^2, \sigma^2_1)}{\partial \sigma^2_1} \theta(\sigma^3_1, \sigma^3) \pi_{\bar a}(\tau, \vec \sigma_1) +\\
&+ \gamma_{\bar a 3} \int d^3\sigma_1 \theta(\sigma^1_1, \sigma^1) \delta(\sigma^2, \sigma^2_1) \frac{\partial \delta(\sigma^3, \sigma^3_1)}{\partial \sigma^3_1} \pi_{\bar a}(\tau, \vec \sigma_1)\Big] =\\
&= \frac{\sqrt{3}}{2} \sum_{\bar a} \Big[\gamma_{\bar a 1} \int_{\sigma^2}^{\infty} d\sigma^2_1 \int_{\sigma^3}^{\infty} d\sigma^3_1 \frac{\partial^2 \pi_{\bar a}(\tau, \sigma^1 \sigma^2_1 \sigma^3_1)}{(\partial \sigma^1)^2} +
\end{aligned}
$$

$$
\begin{aligned}
&+ \gamma_{\bar a 2} \int_{\sigma^3}^{\infty} d\sigma_1^3 \frac{\partial \pi_{\bar a}(\tau, \sigma^1\sigma^2\sigma^3_1)}{\partial \sigma^2} + \\
&+ \gamma_{\bar a 3} \int_{\sigma^2}^{\infty} d\sigma_1^2 \frac{\partial \pi_{\bar a}(\tau, \sigma^1\sigma^2_1\sigma^3)}{\partial \sigma^3} \Big],
\end{aligned}
$$

$$
\begin{aligned}
{}^3\check{\tilde\pi}^1_{(3)}(\tau,\vec\sigma) &= {}^3\check{\tilde\pi}^3_{(1)}(\tau,\vec\sigma),\\
{}^3\check{\tilde\pi}^2_{(3)}(\tau,\vec\sigma) &= {}^3\check{\tilde\pi}^3_{(2)}(\tau,\vec\sigma),\\
{}^3\check{\tilde\pi}^3_{(3)}(\tau,\vec\sigma) &= \sqrt{3}\sum_{\bar a}\gamma_{\bar a 3}\pi_{\bar a}(\tau,\vec\sigma),
\end{aligned}
$$

or

$$
\begin{aligned}
{}^3\check{\tilde\pi}^r_{(a)}(\tau,\vec\sigma) &= \sqrt{3}\sum_{\bar a}\gamma_{\bar a r}\delta^r_{(a)}\pi_{\bar a}(\tau,\vec\sigma) + \frac{\sqrt{3}}{2}[1-\delta^r_{(a)}]\sum_{\bar a u}\gamma_{\bar a u}[1-2(\delta_{ru}+\delta_{au})]\\
&\quad \frac{\partial^2}{(\partial\sigma^u)^2}\int_{\sigma^r}^{\infty} d\sigma^r_1 \int_{\sigma^a}^{\infty} d\sigma^a_1\, \pi_{\bar a}(\tau, \sigma^r_1\sigma^a_1\sigma^{k\neq r,a}).
\end{aligned}
\qquad (2.14)
$$

Clearly Eqs.(2.4) are satisfied.

The solution (2.12) for the $G^{(o)ru}_{(a)}$'s is such that the momenta ${}^3\check{\tilde\pi}^r_{(a)}(\tau,\vec\sigma)$ of Eq.(2.14) tend to zero for $|\vec\sigma| \rightarrow \infty$, as required by Eqs.(1.2), *if the momenta* $\pi_{\bar a}(\tau,\vec\sigma)$ *satisfy the restrictions*

$$
\int_{-\infty}^{+\infty} d\sigma^r\, \pi_{\bar a}(\tau,\vec\sigma) = 0. \qquad (2.15)
$$

For instance these restrictions are satisfied if $\pi_{\bar a}(\tau,\vec\sigma) = \frac{\partial^3\tilde\pi_{\bar a}(\tau,\vec\sigma)}{\partial\sigma^1\partial\sigma^2\partial\sigma^3}$ with $\frac{\partial^2\tilde\pi_{\bar a}(\tau,\vec\sigma)}{\partial\sigma^i\partial\sigma^j} \rightarrow_{\sigma^k\rightarrow\infty} 0$ $[i,j,k$ cyclic$]$ in a direction-independent way.

We have not succeeded in finding a solution without these restrictions. Eqs.(2.15) can be thought of as 6 additional constraints defined on 2-dimensional surfaces. As we shall see, the consistency of these restrictions with the final Hamilton equations will impose the following restrictions on the DO's $r_{\bar a}(\tau,\vec\sigma)$

$$
\int_{-\infty}^{+\infty} d\sigma^r\, r_{\bar a}(\tau,\vec\sigma) = 0. \qquad (2.16)
$$

Therefore Eqs.(2.15) and (2.16) are 6 pairs of second class constraints and we could think to find a Shanmugadhasan canonical transformation from the DO's $r_{\bar a}(\tau,\vec\sigma)$, $\pi_{\bar a}(\tau,\vec\sigma)$ to a new basis in which 6 pairs of conjugate variables vanish due to Eqs.(2.15),(2.16) and the physics is concentrated in the remaining pairs. However, we shall not look for such a transformation, because it is highly non-trivial due to the fact that these constraints are defined only on 2-dimensional surfaces. We shall go on to work with the DO's $r_{\bar a}(\tau,\vec\sigma)$, $\pi_{\bar a}(\tau,\vec\sigma)$ even if this will imply formal complications.

In conclusion, we have been able to solve *all* the Hamiltonian constraints of tetrad gravity on the linearized WSW hyper-surfaces of our gauge.

III. THE WEAK ADM ENERGY, THE LAPSE AND SHIFT FUNCTIONS AND THE 4-METRIC.

After the solution of all the constraints, in this Section we determine the weak ADM energy, the lapse and shift functions and the 4-metric of our linearized theory. Since we need the weak ADM energy, namely the Hamiltonian in the rest-frame instant form of tetrad gravity, see Eq.(A11), we need the following results of Appendix A: Eqs.(A10), where there is the expression of the weak and strong ADM Poincare' generators, and Eqs.(A12), where there is the expression of the weak ADM Poincare' generators in our gauge in the canonical basis (1.3).

In Eqs.(A11) we put: i) $\phi = 1 + \frac{1}{2} q + O(r^2_{\bar a})$ with $q(\tau, \vec \sigma)$ given by the solution (2.2) of the linearized Lichnerowicz equation; ii) the expression (2.14) for the cotriad momenta. In this way we get the form $\hat E_{ADM}[r_{\bar a}, \pi_{\bar a}]$ of the weak ADM energy only in terms of the DO's of our completely fixed gauge.

A. The ADM Energy and the Lapse and Shift Funcions.

The *Hamiltonian linearization* of tetrad gravity in our completely fixed gauge is defined by *approximating the weak ADM energy* $\hat E_{ADM}[r_{\bar a}, \pi_{\bar a}]$ with the quadratic functional of $r_{\bar a}(\tau, \vec \sigma)$ and of $\pi_{\bar a}(\tau, \vec \sigma)$ contained in it

$$
\begin{aligned}
\hat E_{ADM} &= - \epsilon \hat P^\tau_{ADM,R} = \frac{12\pi G}{c^3} \int d^3\sigma \sum_{\bar a} \pi^2_{\bar a}(\tau, \vec \sigma) + \\
&+ \frac{24\pi G}{c^3} \sum_{\bar a \bar b} \sum_{uv} \delta^u_{(a)} \gamma_{\bar a u} \gamma_{\bar b v} \int d^3\sigma_1 d^3\sigma_2 \mathcal{T}^{(o)u}_{(a)v}(\vec \sigma_1, \vec \sigma_2) \pi_{\bar a}(\tau, \vec \sigma_1) \pi_{\bar b}(\tau, \vec \sigma_2) + \\
&+ \frac{6\pi G}{c^3} \sum_{\bar a \bar b} \sum_{rs} \gamma_{\bar a r} \gamma_{\bar b s} \int d^3\sigma d^3\sigma_1 d^3\sigma_2 \Big[\sum_u \mathcal{T}^{(o)u}_{(a)r}(\vec \sigma, \vec \sigma_1) \mathcal{T}^{(o)u}_{(a)s}(\vec \sigma, \vec \sigma_2) + \\
&+ \sum_{uv} (\delta^u_{(b)} \delta^v_{(a)} - \delta^u_{(a)} \delta^v_{(b)}) \mathcal{T}^{(o)u}_{(a)r}(\vec \sigma, \vec \sigma_1) \mathcal{T}^{(o)v}_{(b)s}(\vec \sigma, \vec \sigma_2) \Big] \pi_{\bar a}(\tau, \vec \sigma_1)\, \pi_{\bar b}(\tau, \vec \sigma_2) - \\
&- \frac{c^3}{16\pi G} \sum_r \int d^3\sigma \Big[\frac{1}{6} \Big(\sum_{\bar a u} \gamma_{\bar a u} \frac{\partial}{\partial \sigma^r} \int d^3\sigma_1 \frac{\partial^2_{1u} r_{\bar a}(\tau, \vec \sigma_1)}{4\pi |\vec \sigma - \vec \sigma_1|} \Big)^2 - \\
&- \frac{1}{3} \sum_{\bar a} \Big(\partial_r r_{\bar a}(\tau, \vec \sigma) \Big)^2 + \frac{2}{3} \Big(\sum_{\bar a} \gamma_{\bar a r} \partial_r r_{\bar a}(\tau, \vec \sigma) \Big)^2 - \\
&- \frac{1}{3} \sum_{\bar a \bar b} \sum_u \gamma_{\bar a r} \partial_r r_{\bar a}(\tau, \vec \sigma) \gamma_{\bar b u} \frac{\partial}{\partial \sigma^r} \int d^3\sigma_1 \frac{\partial^2_{1u} r_{\bar b}(\tau, \vec \sigma_1)}{4\pi |\vec \sigma - \vec \sigma_1|} \Big] = \\
&= \frac{12\pi G}{c^3} \int d^3\sigma \sum_{\bar a} \pi^2_{\bar a}(\tau, \vec \sigma) + \frac{12\, \pi G}{c^3} \sum_{\bar a \bar b} \sum_{rs} \gamma_{\bar a r} \gamma_{\bar b s} \\
&\int d^3\sigma d^3\sigma_1 d^3\sigma_2\, \sum_u \mathcal{T}^{(o)u}_{(a)r}(\vec \sigma, \vec \sigma_1) \mathcal{T}^{(o)u}_{(a)s}(\vec \sigma, \vec \sigma_2)\, \pi_{\bar a}(\tau, \vec \sigma_1)\, \pi_{\bar b}(\tau, \vec \sigma_2) - \\
&- \frac{c^3}{16\pi G} \sum_r \int d^3\sigma \Big[\frac{1}{6} \Big(\sum_{\bar a u} \gamma_{\bar a u} \frac{\partial}{\partial \sigma^r} \int d^3\sigma_1 \frac{\partial^2_{1u} r_{\bar a}(\tau, \vec \sigma_1)}{4\pi |\vec \sigma - \vec \sigma_1|} \Big)^2 - \\
&- \frac{1}{3} \sum_{\bar a} \Big(\partial_r r_{\bar a}(\tau, \vec \sigma) \Big)^2 + \frac{2}{3} \Big(\sum_{\bar a} \gamma_{\bar a r} \partial_r r_{\bar a}(\tau, \vec \sigma) \Big)^2 -
\end{aligned}
$$

$$- \frac{1}{3} \sum_{\bar{a}\bar{b}} \sum_u \gamma_{\bar{a}r} \partial_r r_{\bar{a}}(\tau, \vec{\sigma}) \gamma_{\bar{b}u} \frac{\partial}{\partial \sigma^r} \int d^3\sigma_1 \frac{\partial^2_{1u} r_{\bar{b}}(\tau, \vec{\sigma}_1)}{4\pi |\vec{\sigma} - \vec{\sigma}_1|} \Big] + O(r^3_{\bar{a}}). \tag{3.1}$$

See Eq.(C13) of Appendix C for the Fourier transform of the ADM energy.

The determination of the *bulk lapse function* $n(\tau, \vec{\sigma})$ is done by solving the integral equation (A13) written in our completely fixed gauge. Therefore we must evaluate $\frac{\delta \hat{\mathcal{H}}_R(\tau, \vec{\sigma}_1)}{\delta \phi(\tau, \vec{\sigma})}$ and $\frac{\delta \hat{E}_{ADM}}{\delta \phi(\tau, \vec{\sigma})}$. From the linearized version of the super-hamiltonian constraint given before Eq.(2.1) we get

$$\frac{\delta \hat{\mathcal{H}}_R(\tau, \vec{\sigma}_1)}{\delta \phi(\tau, \vec{\sigma})} = -\epsilon \frac{c^3}{2\pi G} \triangle_1 \delta^3(\vec{\sigma}, \vec{\sigma}_1) + O(r^2_{\bar{a}}), \tag{3.2}$$

while from Eqs.(A12) and (2.1) we get

$$\begin{aligned} \frac{\delta \hat{E}_{ADM}}{\delta \phi(\tau, \vec{\sigma})} &= \epsilon \frac{c^3}{4\pi G} \int d^3\sigma_1 \sum_r \Big[2\partial_{1r} q(\tau, \vec{\sigma}_1) - \frac{1}{\sqrt{3}} \sum_{\bar{a}} \gamma_{\bar{a}r} \partial_{1r} r_{\bar{a}}(\tau, \vec{\sigma}_1) \Big] \partial_{1r} \delta^3(\vec{\sigma}, \vec{\sigma}_1) + O(r^2_{\bar{a}}) = \\ &= -\epsilon \frac{c^3}{2\pi G} \Big[\triangle q(\tau, \vec{\sigma}) - \frac{1}{2\sqrt{3}} \sum_{\bar{a}r} \gamma_{\bar{a}r} \partial^2_r r_{\bar{a}}(\tau, \vec{\sigma}) \Big] + O(r^2_{\bar{a}}) = \\ &= 0 + O(r^2_{\bar{a}}). \end{aligned} \tag{3.3}$$

Then the integral equation (A13) for the lapse function becomes the following partial differential equation

$$-\epsilon \frac{c^3}{2\pi G} \int d^3\sigma_1 n(\tau, \vec{\sigma}_1) \triangle_1 \delta^3(\vec{\sigma}, \vec{\sigma}_1) = -\epsilon \frac{c^3}{2\pi G} \triangle n(\tau, \vec{\sigma}) = 0 + O(r^2_{\bar{a}}),$$

$$\Downarrow$$

$$n(\tau, \vec{\sigma}) = 0 + O(r^2_{\bar{a}}). \tag{3.4}$$

The determination of the *bulk shift functions* uses this result and the linearized form of Eq.(A14) (see Eq.(C14) of Appendix C for the Fourier transform),

$$\begin{aligned} n_r(\tau, \vec{\sigma}) &= -\sqrt{3} \frac{4\pi G}{c^3} \int d^3\sigma_1 \sum_{wu} (\delta_{wu} \delta_{(a)(b)} + \delta_{(a)u} \delta_{(b)w} - \delta_{(a)w} \delta_{(b)u}) \\ &\quad \sum_v \int d^3\sigma_2 \Big[\delta^w_v \delta_{(a)v} \delta^3(\vec{\sigma}_1, \vec{\sigma}_2) + \mathcal{T}^{(o)w}_{(a)v}(\vec{\sigma}_1, \vec{\sigma}_2) \Big] \\ &\quad \sum_{\bar{a}} \gamma_{\bar{a}v} \pi_{\bar{a}}(\tau, \vec{\sigma}_2) G^{(o)ur}_{(b)}(\vec{\sigma}_1, \vec{\sigma}) + O(r^2_{\bar{a}}) = \\ &= \frac{\partial}{\partial \sigma^r} \Big(\sqrt{3} \frac{2\pi G}{c^3} \sum_{\bar{a}v} \gamma_{\bar{a}v} \\ &\quad \Big[\sum_{ua, u \neq a} [1 - 2(\delta_{uv} + \delta_{av})] \int_{-\infty}^{\sigma^u} d\sigma^u_1 \int_{-\infty}^{\sigma^a} d\sigma^a_1 \int_{\sigma^u_1}^{\infty} d\sigma^u_2 \int_{\sigma^a_1}^{\infty} d\sigma^a_2 \\ &\quad \frac{\partial^2 \pi_{\bar{a}}(\tau, \sigma^u_2 \sigma^a_2 \sigma^{k \neq u,a}_2)}{(\partial \sigma^v_2)^2} |_{\sigma^k_2 = \sigma^k} - \end{aligned}$$

$$
- 2 \sum_{u \neq r} [1 - 2(\delta_{uv} + \delta_{rv})] \int_{-\infty}^{\sigma^r} d\sigma_1^r \int_{-\infty}^{\sigma^u} d\sigma_1^u \int_{\sigma_1^r}^{\infty} d\sigma_2^r \int_{\sigma_1^u}^{\infty} d\sigma_2^u
\frac{\partial^2 \pi_{\bar{a}}(\tau, \sigma_2^r \sigma_2^u \sigma_2^{k \neq r,u})}{(\partial \sigma_2^v)^2} |_{\sigma_2^k = \sigma^k} \Big] \Big) + O(r_{\bar{a}}^2). \qquad (3.5)
$$

In conclusion in our completely fixed gauge the lapse and shift functions are

$$
\begin{aligned}
N(\tau, \vec{\sigma}) &= -\epsilon + n(\tau, \vec{\sigma}) = -\epsilon + O(r_{\bar{a}}^2), \\
N_r(\tau, \vec{\sigma}) &= n_r(\tau, \vec{\sigma}) = -\epsilon\, {}^4\hat{g}_{\tau r}(\tau, \vec{\sigma}).
\end{aligned} \qquad (3.6)
$$

Since the shift functions are of order $O(r_{\bar{a}})$, we get the following results :

i) the 4-coordinates associated to our Hamiltonian gauge are *not synchronous*, so that we are using non-time-orthogonal reference frames and we cannot use Einstein's convention for the synchronization of clocks [26]; a non-standard definition of simultaneity of distant time-like observers is needed, consistent with the Hamiltonian description based on the Cauchy simultaneity WSW space-like hyper-surfaces Σ_τ;

ii) there may be coordinate-dependent gravito-magnetic effects;

iii) the *velocity of light becomes direction-dependent* (see Ref. [26]) : if u^i is a unit 3-vector with respect to the 3-metric ${}^3\tilde{\gamma}_{rs} = -\epsilon({}^3g_{rs} + \frac{n_r n_s}{\epsilon\, {}^4g_{\tau\tau}})$, i.e. ${}^3\tilde{\gamma}_{rs} u^r u^s = 1$, [see Appendix A of Ref. [2], after Eq.(A5)], the light velocity in direction u^i is $w(u^i) = \frac{\sqrt{{}^4g_{\tau\tau}}}{\frac{u^r n_r}{\sqrt{{}^4g_{\tau\tau}}} + 1} = 1 - u^r\, n_r + O(r_{\bar{a}}^2)$ with $\Big(\frac{u^r n_r}{\sqrt{{}^4g_{\tau\tau}}}\Big)^2 = (u^r\, n_r)^2 + O(r_{\bar{a}}^3) < 1$.

B. The Linearized 4-Metric.

After the solution of all the constraints and the determination of the lapse and shift functions, the *4-metric* of our linearized space-time with $\Sigma_\tau^{(WSW)}$-adapted coordinates in the 3-orthogonal gauge with $\pi_\phi(\tau, \vec{\sigma}) = 0$ becomes (we write it in the form of a perturbation of the Minkowski metric in Cartesian coordinates *only to visualize* the deviations from special relativity of this background-independent post-Minkowskian space-time in the 4-coordinates associated with our preferred 3-orthogonal gauge)

$$
{}^4\hat{g}_{AB}(\tau, \vec{\sigma}) = {}^4\eta_{AB} + {}^4h_{AB}(\tau, \vec{\sigma}),
$$

$$
\begin{aligned}
{}^4h_{\tau\tau}(\tau, \vec{\sigma}) &= 0 + O(r_{\bar{a}}^2), \\
{}^4h_{\tau r}(\tau, \vec{\sigma}) &= -\epsilon n_r(\tau, \vec{\sigma}) \overset{def}{=} -\epsilon \frac{2}{c^2} A_{GEM,3-0,r}(\tau, \vec{\sigma}) = \frac{\partial}{\partial \sigma^r} \Big(\epsilon \sqrt{3} \frac{2\pi G}{c^3} \sum_{\bar{a}v} \gamma_{\bar{a}v} \\
&\Big[\sum_{ua, u \neq a} [1 - 2(\delta_{uv} + \delta_{av})] \\
&\int_{-\infty}^{\sigma^u} d\sigma_1^u \int_{-\infty}^{\sigma^a} d\sigma_1^a \int_{\sigma_1^u}^{\infty} d\sigma_2^u \int_{\sigma_1^a}^{\infty} d\sigma_2^a \frac{\partial^2 \pi_{\bar{a}}(\tau, \sigma_2^u \sigma_2^a \sigma_2^{k \neq u,a})}{(\partial \sigma_2^v)^2} |_{\sigma_2^k = \sigma^k} -
\end{aligned}
$$

$$
\begin{aligned}
&- 2 \sum_{u\neq r} [1 - 2(\delta_{uv} + \delta_{rv})] \\
&\qquad \int_{-\infty}^{\sigma^r} d\sigma_1^r \int_{-\infty}^{\sigma^u} d\sigma_1^u \int_{\sigma_1^r}^{\infty} d\sigma_2^r \int_{\sigma_1^u}^{\infty} d\sigma_2^u \, \frac{\partial^2 \pi_{\bar a}(\tau, \sigma_2^r \sigma_2^u \sigma^{k\neq r,u})}{(\partial \sigma_2^v)^2} |_{\sigma_2^k = \sigma^k} \Big] \Big) + \\
&+ O(r_{\bar a}^2), \\
{}^4h_{rs}(\tau, \vec \sigma) &= -\epsilon [{}^3\hat g_{rs}(\tau, \vec \sigma) - \delta_{rs}] \overset{def}{=} \delta_{rs}\, k_r(\tau, \vec \sigma) = \\
&= -2\epsilon \Big[q(\tau, \vec \sigma) + \frac{2}{\sqrt{3}} \sum_{\bar a} \gamma_{\bar ar} r_{\bar a}(\tau, \vec \sigma) \Big] \delta_{rs} + O(r_{\bar a}^2) = \\
&= - \frac{2\epsilon}{\sqrt{3}} \sum_{\bar a} \Big[\gamma_{\bar ar} r_{\bar a}(\tau, \vec \sigma) - \frac{1}{2} \sum_u \gamma_{\bar au} \int d^3\sigma_1 \frac{\partial^2_{1u} r_{\bar a}(\tau, \vec \sigma_1)}{4\pi |\vec \sigma - \vec \sigma_1|} \Big] \delta_{rs} + O(r_{\bar a}^2),
\end{aligned}
$$

$$
\begin{aligned}
\sqrt{{}^4\hat g(\tau, \vec \sigma)} &= -\epsilon \sqrt{\hat \gamma(\tau, \vec \sigma)} + O(r_{\bar a}^2) = -\epsilon\, \phi^6(\tau, \vec \sigma) + O(r_{\bar a}^2) = -\epsilon\, [1 + 3\, q(\tau, \vec \sigma)] + O(r_{\bar a}^2) = \\
&= -\epsilon\, \Big[1 - \frac{\sqrt{3}}{2} \sum_{u\bar a} \gamma_{\bar au} \int d^3\sigma_1 \frac{\partial^2_{1u} r_{\bar a}(\tau, \vec \sigma_1)}{4\pi |\vec \sigma - \vec \sigma_1|} \Big] + O(r_{\bar a}^2), \\
{}^4\hat g^{\tau\tau}(\tau, \vec \sigma) &= \epsilon + O(r_{\bar a}^2), \\
{}^4\hat g^{\tau r}(\tau, \vec \sigma) &= -\epsilon \delta^{rs} n_s(\tau, \vec \sigma) + O(r_{\bar a}^2), \\
{}^4\hat g^{rs}(\tau, \vec \sigma) &= -\epsilon\, \Big({}^3\hat g^{rs} - n^r\, n^s \Big)(\tau, \vec \sigma) = \\
&= -\epsilon \Big[1 - \frac{2}{\sqrt{3}} \sum_{\bar a} \Big(\gamma_{\bar ar} r_{\bar a}(\tau, \vec \sigma) - \frac{1}{2} \sum_u \gamma_{\bar au} \int d^3\sigma_1 \frac{\partial^2_{1u} r_{\bar a}(\tau, \vec \sigma_1)}{4\pi |\vec \sigma - \vec \sigma_1|} \Big) \Big] \delta^{rs} + O(r_{\bar a}^2).
\end{aligned}
\tag{3.7}
$$

Therefore, the 3-metric on the linearized WSW hyper-surfaces of our gauge is only conformal to the Euclidean 3-metric, namely the linearized $\Sigma_\tau^{(WSW)}$ of our gauge are *conformally flat*.

The linearized cotriads, triads and adapted cotetrads and tetrads become [see Eqs.(A7) and (A16)]

$$
\begin{aligned}
{}^3\hat e_{(a)r}(\tau, \vec \sigma) &= \delta_{(a)r} \Big[1 + \frac{1}{\sqrt{3}} \sum_{\bar a} \Big(\gamma_{\bar ar} r_{\bar a}(\tau, \vec \sigma) - \\
&- \frac{1}{2} \sum_u \gamma_{\bar au} \int d^3\sigma_1 \frac{\partial^2_{1u} r_{\bar a}(\tau, \vec \sigma_1)}{4\pi |\vec \sigma - \vec \sigma_1|} \Big) \Big] + O(r_{\bar a}^2), \\
{}^3\hat e^r_{(a)}(\tau, \vec \sigma) &= \delta^r_{(a)} \Big[1 - \frac{1}{\sqrt{3}} \sum_{\bar a} \Big(\gamma_{\bar ar} r_{\bar a}(\tau, \vec \sigma) + \\
&+ \frac{1}{2} \sum_u \gamma_{\bar au} \int d^3\sigma_1 \frac{\partial^2_{1u} r_{\bar a}(\tau, \vec \sigma_1)}{4\pi |\vec \sigma - \vec \sigma_1|} \Big) \Big] + O(r_{\bar a}^2), \\
{}^3\hat e(\tau, \vec \sigma) &= \sqrt{\hat \gamma(\tau, \vec \sigma)} = 1 + 3 q(\tau, \vec \sigma) + O(r_{\bar a}^2) = \\
&= 1 - \frac{\sqrt{3}}{2} \sum_{\bar au} \gamma_{\bar au} \int d^3\sigma_1 \frac{\partial^2_{1u} r_{\bar a}(\tau, \vec \sigma_1)}{4\pi |\vec \sigma - \vec \sigma_1|} \Big) + O(r_{\bar a}^2),
\end{aligned}
$$

$$
\begin{aligned}
{}^3\hat{g}_{rs}(\tau,\vec{\sigma}) &= -\epsilon\, {}^4\hat{g}_{rs}(\tau,\vec{\sigma}) = {}^3\hat{e}_{(a)r}(\tau,\vec{\sigma})\, {}^3\hat{e}_{(a)s}(\tau,\vec{\sigma}) = \\
&= \Big[1 + \frac{2}{\sqrt{3}} \sum_{\bar{a}} \Big(\gamma_{\bar{a}r} r_{\bar{a}}(\tau,\vec{\sigma}) - \\
&- \frac{1}{2} \sum_{u} \gamma_{\bar{a}u} \int d^3\sigma_1 \frac{\partial^2_{1u} r_{\bar{a}}(\tau,\vec{\sigma}_1)}{4\pi|\vec{\sigma}-\vec{\sigma}_1|}\Big)\Big]\delta^{rs} + O(r^2_{\bar{a}}),
\end{aligned}
\qquad (3.8)
$$

$$
\begin{aligned}
{}^4_{(\Sigma)}\check{\tilde{E}}^A_{(o)}(\tau,\vec{\sigma}) &= l^A(\tau,\vec{\sigma}) = -\epsilon(1; -\delta^{rs} n_s(\tau,\vec{\sigma})) + O(r^2_{\bar{a}}), \\
{}^4_{(\Sigma)}\check{\tilde{E}}^A_{(a)}(\tau,\vec{\sigma}) &= (0; {}^3\hat{e}^r_{(a)}(\tau,\vec{\sigma})) = \\
&= (0; \delta^r_{(a)}\Big[1 - \frac{1}{\sqrt{3}} \sum_{\bar{a}} \Big(\gamma_{\bar{a}r} r_{\bar{a}}(\tau,\vec{\sigma}) + \\
&+ \frac{1}{2} \sum_{u\bar{a}} \gamma_{\bar{a}u} \int d^3\sigma_1 \frac{\partial^2_{1u} r_{\bar{a}}(\tau,\vec{\sigma}_1)}{4\pi|\vec{\sigma}-\vec{\sigma}_1|}\Big)\Big]\ ) + O(r^2_{\bar{a}}), \\
{}^4_{(\Sigma)}\check{\tilde{E}}^{(o)}_A(\tau,\vec{\sigma}) &= l_A(\tau,\vec{\sigma}) = -\epsilon(1; 0) + O(r^2_{\bar{a}}), \\
{}^4_{(\Sigma)}\check{\tilde{E}}^{(a)}_A(\tau,\vec{\sigma}) &= (\delta^r_{(a)} n_r(\tau,\vec{\sigma}); {}^3\hat{e}_{(a)r}) = \\
&= (\delta^r_{(a)} n_r(\tau,\vec{\sigma});\ \delta_{(a)r}\ \Big[1 + \frac{1}{\sqrt{3}} \sum_{\bar{a}} \Big(\gamma_{\bar{a}r} r_{\bar{a}}(\tau,\vec{\sigma}) - \\
&- \frac{1}{2} \sum_{u\bar{a}} \gamma_{\bar{a}u} \int d^3\sigma_1 \frac{\partial^2_{1u} r_{\bar{a}}(\tau,\vec{\sigma}_1)}{4\pi|\vec{\sigma}-\vec{\sigma}_1|}\Big)\Big]\ ) + O(r^2_{\bar{a}}).
\end{aligned}
\qquad (3.9)
$$

Since in our gauge we have $\varphi_{(a)}(\tau,\vec{\sigma}) = 0$, the tetrads and cotetrads ${}^4E^A_{(\alpha)}$ and ${}^4E^{(\alpha)}_A$ of Eqs.(1.1) coincide with those of Eqs.(3.9).

See Appendix D for the associated Christoffel symbols, spin connection, field strength and Riemann tensor of Σ_τ and for the 4-tensors of M^4.

C. The Energy-Momentum Landau-Lifschitz Pseudo-Tensor.

Usually the energy of a gravitational wave on the Minkowski background is evaluated [22,24] as a mean value over various wave-lengths of the coordinate-dependent Landau-Lifschitz energy obtained from the Landau-Lifschitz symmetric pseudo-tensor [27].

The Landau-Lifschitz pseudo-tensor ${}_{(L)}t^{\mu\nu} = {}_{(L)}t^{\nu\mu}$, which contains no second derivatives of the metric and gives meaningful results only in an asymptotically flat Cartesian coordinate system, was found trying to rewrite the consequence ${}^4\nabla_\mu\, {}^4T^{\mu\nu} \circeq 0$ of the Bianchi identities applied to Einstein's equations in the form $\partial_\mu\big[{}_{(L)}t^{\mu\nu} + \sqrt{{}^4g}\, {}^4T^{\mu\nu}\big] \circeq 0$. Starting from Einstein's equations ${}^4T^{\mu\nu} \circeq \frac{c^3}{8\pi G}({}^4R^{\mu\nu} - \frac{1}{2}{}^4g^{\mu\nu}\, {}^4R)$, one rewrites them as

$$
{}_{(L)}t^{\mu\nu} + \sqrt{{}^4g}\, {}^4T^{\mu\nu} \circeq \partial_\rho h^{\mu[\nu\rho]}, \qquad \mathit{with}
$$

$$
\begin{aligned}
h^{\mu[\nu\rho]} &= -h^{\mu[\rho\nu]} = \frac{c^3}{16\pi G}\partial_\sigma[{}^4g({}^4g^{\mu\nu}\,{}^4g^{\rho\sigma} - {}^4g^{\mu\rho}\,{}^4g^{\nu\sigma})] = \frac{c^3}{16\pi G}\,\partial_\sigma\,\mathcal{T}^{\rho\nu\sigma\mu},\\
&\partial_\nu\partial_\rho h^{\mu[\nu\rho]} = \partial_\nu\partial_\rho\partial_\sigma\,\mathcal{T}^{\rho\nu\sigma\mu} = 0,\\
\mathcal{T}^{\alpha\nu\beta\mu} &= {}^4\hat{g}^{\alpha\beta}\,{}^4\hat{g}^{\mu\nu} - {}^4\hat{g}^{\alpha\mu}\,{}^4\hat{g}^{\beta\nu} = -\mathcal{T}^{\nu\alpha\beta\mu} = -\mathcal{T}^{\alpha\nu\mu\beta} = \mathcal{T}^{\beta\mu\alpha\nu},\\
&[\mathcal{T}^{\alpha\nu\beta\mu} + \mathcal{T}^{\alpha\beta\mu\nu} + \mathcal{T}^{\alpha\mu\nu\beta} = 0],\quad {}^4\hat{g}^{\mu\nu} = \sqrt{{}^4g}\,{}^4g^{\mu\nu}.
\end{aligned}
\tag{3.10}
$$

Then one gets

$$
\begin{aligned}
{}_{(L)}t^{\mu\nu} &= \frac{c^3}{16\pi G}[\partial_\rho\,{}^4\hat{g}^{\mu\nu}\,\partial_\gamma\,{}^4\hat{g}^{\gamma\rho} - \partial_\rho\,{}^4\hat{g}^{\mu\rho}\partial_\gamma\,{}^4\hat{g}^{\nu\gamma} + \frac{1}{2}{}^4g^{\mu\nu}\,{}^4g_{\rho\sigma}\,\partial_\gamma\,{}^4\hat{g}^{\rho\delta}\,\partial_\delta\,{}^4\hat{g}^{\sigma\gamma} -\\
&- ({}^4g^{\mu\rho}\,{}^4g_{\gamma\delta}\partial_\sigma\,{}^4\hat{g}^{\nu\delta}\,\partial_\rho\,{}^4\hat{g}^{\gamma\sigma} + {}^4g^{\nu\rho}\,{}^4g_{\gamma\delta}\partial_\sigma\,{}^4\hat{g}^{\mu\delta}\,\partial_\rho\,{}^4\hat{g}^{\gamma\sigma}) + {}^4g_{\rho\sigma}\,{}^4g^{\gamma\delta}\partial_\gamma\,{}^4\hat{g}^{\mu\rho}\partial_\delta\,{}^4\hat{g}^{\nu\sigma} +\\
&+ \frac{1}{8}(2\,{}^4g^{\mu\rho}\,{}^4g^{\nu\sigma} - {}^4g^{\mu\nu}\,{}^4g^{\rho\sigma})(2\,{}^4g_{\gamma\delta}\,{}^4g_{\alpha\beta} - {}^4g_{\delta\alpha}\,{}^4g_{\gamma\beta})\partial_\rho\,{}^4\hat{g}^{\gamma\beta}\partial_\sigma\,{}^4\hat{g}^{\delta\alpha}].
\end{aligned}
\tag{3.11}
$$

However, as noted in Ref. [28] and emphasized in Ref. [29], if the energy flux carried away to infinity by gravitational radiation in a given asymptotically flat space-time is deduced from the Landau-Lifschitz pseudo-tensor, then the result is *reliable only* on an appropriately chosen flat background metric and then a connection between Bondi energy at null infinity and ADM energy at spatial infinity can be established [29]. Since in our approach to Hamiltonian linearization of Einstein's equation we have no background metric, the Landau-Lifschitz pseudo-tensor is not a useful quantity.

If we write the Landau-Lifschitz in $\Sigma_\tau^{(WSW)}$-adapted coordinates, ${}_{(L)}t^{AB}(\tau,\vec{\sigma})$ and then we choose our special 3-orthogonal coordinates, then the Landau-Lifschitz 4-momentum $P^A_{LL} = \int d^3\sigma\,{}_{(L)}t^{A\tau}(\tau,\vec{\sigma})$ has to be contrasted with the weak ADM energy $\hat{E}_{ADM}$, namely the Hamiltonian in the rest-frame instant form of gravity, and with the vanishing weak ADM momentum $\hat{P}^r_{ADM} \approx 0$ evaluated in those coordinates.

¿From an explicit calculation done by using Eqs.(3.7) it turns out that P^A_{LL} does not agree with E_{ADM} and $\hat{P}^r_{ADM}$ [for its expression see Eq.(5.11)]

$$
\begin{aligned}
E_{LL} &= \int d^3\sigma\,{}_{(LL)}t^{\tau\tau}(\tau,\vec{\sigma}) =\\
&= \frac{c^3}{16\pi G}\int d^3\sigma\Big(- \frac{25}{24}(\partial_\tau A)^2 - \frac{175}{72}\sum_i(\partial_i A)^2 - \frac{5}{12}\partial_\tau A\sum_i\partial_\tau f_i +\\
&+ \frac{1}{4}\sum_i(\partial_\tau f_i)^2 - \frac{1}{8}(\sum_i\partial_\tau f_i)^2 - \frac{5}{3}\partial_\tau A\sum_i\partial_i n_i +\\
&- \frac{5}{12}\sum_{i,j}\partial_j A\partial_j f_i - \sum_i\partial_\tau f_i\partial_i n_i - \frac{5}{3}\sum_i\partial_i A\partial_i f_i +\\
&+ \frac{1}{2}\sum_{r,s}(\partial_s n_r)^2 + \frac{1}{2}\sum_{r,s}\partial_s n_r\partial_r n_s - \sum_{r,s}\partial_s n_s\partial_r n_r +\\
&- \frac{1}{2}\sum_i(\partial_i f_i)^2 - \frac{1}{8}\sum_{i,j,r}\partial_i f_j\partial_i f_r + \frac{1}{4}\sum_{i,j}(\partial_j f_i)^2\Big)(\tau,\vec{\sigma}) +\\
&+ O(r^3_{\bar{a}}) \neq \hat{E}_{ADM},
\end{aligned}
$$

$$
\begin{aligned}
P^r_{LL} &= \int d^3\sigma\, {}_{(LL)}t^{r\tau}(\tau,\vec{\sigma}) = \\
&= \frac{c^3}{16\pi G} \int d^3\sigma\, \Big(\frac{115}{36}\, \partial_\tau A \partial_r A + \frac{5}{12}\, \partial_r A \sum_i \partial_\tau f_i + \frac{2}{3}\, \partial_r A \partial_\tau f_r + \\
&+ \frac{5}{3}\, \partial_\tau A \partial_r f_r + \frac{5}{12}\, \partial_\tau A \sum_i \partial_r f_i + \frac{1}{4} \sum_{i,j} \partial_\tau f_i \partial_r f_j - \\
&- \frac{1}{2} \sum_i \partial_\tau f_i \partial_r f_i + \partial_\tau f_r \partial_r f_r - \frac{7}{3} \sum_i \partial_i A \partial_i n_r + \\
&+ \frac{4}{3} \sum_i \partial_r A \partial_i n_i + \sum_i \partial_i A \partial_r n_i - \sum_i \partial_i f_r \partial_i n_r - \\
&- \sum_i \partial_i f_i \partial_i n_r + \sum_i \partial_r f_i \partial_i n_i + \partial_r f_r \sum_i \partial_i n_i \Big)(\tau,\vec{\sigma}) + \\
&+ O(r^3_{\bar{a}}) \neq \hat{P}^r_{ADM},
\end{aligned}
$$

where

$$
\begin{aligned}
A(\tau,\vec{\sigma}) &= \sqrt{3} \sum_{u\bar{a}} \gamma_{\bar{a}u} \int d^3\sigma_1 \frac{\partial^2_{1u} r_{\bar{a}}(\tau,\vec{\sigma}_1)}{4\pi|\vec{\sigma}-\vec{\sigma}_1|} \\
f_i(\tau,\vec{\sigma}) &= \frac{2}{\sqrt{3}} \sum_{\bar{a}} \gamma_{\bar{a}i}\, r_{\bar{a}}(\tau,\vec{\sigma}).
\end{aligned}
\tag{3.12}
$$

Therefore, when in a future paper we will add matter to tetrad gravity, we will have to devise a method independent from the Landau-Lifschitz pseudo-tensor to identify the energy of the matter and its variation due to the emission of gravitational waves.

IV. COMPARISON WITH THE LICHNEROWICZ-YORK CONFORMAL APPROACH, WITH THE STANDARD LINEARIZED THEORY IN HARMONIC GAUGES AND WITH THE POST-NEWTONIAN APPROXIMATION.

Since in the literature there are many *coordinate-dependent* definitions of *gravitomagnetism*, which are a source of ambiguities, in this Section we shall review same of them and we will rephrase them in the language of our linearized post-Minkowskian space-time. Moreover we will show that our completely fixed Hamiltonian gauge does not belong to the family of the harmonic gauges used in the standard background-dependent linearization of Einstein's equations. Finally we will make some comments on the post-Newtonian approximation of our space-time.

A. Comparison with the Lichnerowicz-York Conformal Approach.

To establish the connection with the Lichnerowicz-York conformal approach [18,19,16] we need the extrinsic curvature of the WSW hyper-surfaces of our gauge. From Eq.(A15), we get that the *extrinsic curvature and the ADM momentum* of the WSW hyper-surfaces of our gauge in the linearized theory are (see Eq.(C15) of Appendix C for the Fourier transform)

$$\begin{aligned}
{}^3\hat{K}_{rs}(\tau,\vec{\sigma}) &= \epsilon\frac{4\pi G}{c^3}\sum_u(\delta_{ru}\delta_{(a)s}+\delta_{su}\delta_{(a)r}-\delta_{rs}\delta_{(a)u})\,{}^3\check{\tilde{\pi}}^u_{(a)}(\tau,\vec{\sigma})+O(r^2_{\bar{a}}) =\\
&= \epsilon\sqrt{3}\frac{4\pi G}{c^3}\sum_{\bar{a}}\Big[2\delta_{rs}\gamma_{\bar{a}s}\pi_{\bar{a}}(\tau,\vec{\sigma})+\\
&+[1-\delta_{rs}]\sum_w\gamma_{\bar{a}w}[1-2(\delta_{rw}+\delta_{sw})]\\
&\quad\frac{\partial^2}{\partial(\sigma^w)^2}\int_{\sigma^r}^{\infty}d\sigma_1^r\int_{\sigma^s}^{\infty}d\sigma_1^s\;\pi_{\bar{a}}(\tau,\sigma_1^r\sigma_1^s\sigma^{k\neq r,s})\Big]+O(r^2_{\bar{a}}) =\\
&= {}^3\hat{K}^{rs}(\tau,\vec{\sigma})+O(r^2_{\bar{a}}),
\end{aligned}$$

$$\partial_s\,{}^3\hat{K}^{rs}(\tau,\vec{\sigma}) = 0+O(r^2_{\bar{a}}),$$

$$\begin{aligned}
{}^3\hat{\tilde{\Pi}}^{rs}(\tau,\vec{\sigma}) &= \frac{1}{4}\Big[{}^3\hat{e}^s_{(a)}\,{}^3\check{\tilde{\pi}}^r_{(a)}+{}^3\hat{e}^r_{(a)}\,{}^3\check{\tilde{\pi}}^s_{(a)}\Big](\tau,\vec{\sigma}) = \frac{1}{4}\Big[{}^3\check{\tilde{\pi}}^r_{(s)}+{}^3\check{\tilde{\pi}}^s_{(r)}\Big](\tau,\vec{\sigma}) =\\
&= \frac{\sqrt{3}}{2}\Big[\delta_{rs}\sum_{\bar{a}}\gamma_{\bar{a}r}\pi_{\bar{a}}(\tau,\vec{\sigma})+\\
&+\frac{1}{2}(1-\delta_{rs})\sum_{\bar{a}w}\gamma_{\bar{a}w}[1-2(\delta_{rw}+\delta_{sw})]\,\partial^2_w\int_{\sigma^r}^{\infty}d\sigma_1^r\int_{\sigma^s}^{\infty}d\sigma_1^s\,\pi_{\bar{a}}(\tau,\sigma_1^r\sigma_1^s\sigma^{k\neq r,s})\Big]+\\
&+O(r^2_{\bar{a}}) =\\
&= \frac{\epsilon c^3}{16\pi G}({}^3\hat{K}^{ra}-\delta^{rs}\,{}^3\hat{K})+O(r^2_{\bar{a}}) = \frac{\epsilon c^3}{16\pi G}\,{}^3\hat{K}^{rs}+O(r^2_{\bar{a}}),
\end{aligned}$$

$$\Downarrow$$

$$ {}^3\hat{\tilde{\Pi}}^{rs}{}_{|r} = \partial_r\, {}^3\hat{\tilde{\Pi}}^{rs} + O(r^2_{\bar a}) = 0 + O(r^2_{\bar a}), \tag{4.1} $$

since the trace, proportional to *York's time* $\mathcal{T} = -\epsilon \frac{64\pi G}{3c^3}\, {}^3K$, turns out to be

$$ {}^3\hat{K}(\tau,\vec{\sigma}) = \delta^{rs}\, {}^3\hat{K}_{rs}(\tau,\vec{\sigma}) + O(r^2_{\bar a}) = 0 + O(r^2_{\bar a}). \tag{4.2} $$

This means that *the linearized WSW hyper-surfaces are constant mean extrinsic curvature (CMC) surfaces and satisfy the maximal slicing condition.* Moreover, the ADM super-momentum constraints are satisfied.

Therefore, York's traceless *distorsion tensor* ${}^3A_{rs} = {}^3K_{rs} - \frac{1}{3}\, {}^3g_{rs}\, {}^3K = {}^3\hat{K}_{rs} + O(r^2_{\bar a})$, see Ref. [16], Chapter 4.10 and Appendix C of Ref. [30], coincides with the extrinsic curvature in the linearized theory. After a conformal rescaling ${}^3g_{rs} = \phi^4\, {}^3\breve{g}_{rs}$ [17] we get ${}^3\breve{A}^{rs} = {}^3\breve{A}^{rs}_{TT} + {}^3\breve{A}^{rs}_{L} = \phi^{10}\, {}^3A^{rs} = {}^3\hat{K}^{rs} + O(r^2_{\bar a})$, where ${}^3\breve{A}^{rs}_{TT}$ and ${}^3\breve{A}^{rs}_{L}$ are the *transverse traceless* (TT) and *longitudinal* (L) components respectively. The longitudinal component satisfies ${}^3\breve{A}^{rs}_{L\,|s} = {}^3\breve{A}^{rs}{}_{|s} \overset{def}{=} (\breve{\triangle}_L W)^r = \breve{\triangle} W^r + \frac{1}{3}(W^s{}_{|s})^{|r} + {}^3\breve{R}^r{}_s W^s$, where $W^r(\tau,\vec{\sigma})$ is *York's gravito-magnetic vector potential.* In the linearized theory it satisfies $\triangle\Big(\delta^r_s + \frac{\partial^r\partial_s}{3\triangle}\Big) W^s = \partial_s\, {}^3\hat{K}^{rs} + O(r^2_{\bar a}) = 0 + O(r^2_{\bar a})$, whose solution is

$$ W^r(\tau,\vec{\sigma}) = 0 + O(r^2_{\bar a}), $$

$$ \Downarrow $$

$$ \begin{aligned} {}^3\breve{A}^{rs}_{L} &= (LW)^{rs} \overset{def}{=} W^{s|r} + W^{r|s} - \frac{2}{3}\, {}^3\breve{g}^{rs} W^u{}_{|u} = [\delta^r_u \partial^s + \delta^s_u \partial^r - \frac{2}{3}\delta^{rs}\partial_u] W^u + O(r^2_{\bar a}) = \\ &= 0 + O(r^2_{\bar a}). \end{aligned} \tag{4.3} $$

Therefore, York's transverse traceless physical degrees of freedom of momentum type, namely ${}^3\breve{A}^{rs}_{TT} = {}^3\breve{A}^{rs} - {}^3\breve{A}^{rs}_{L} = {}^3\hat{K}^{rs} + O(r^2_{\bar a})$, are linear functionals of the $\pi_{\bar a}$'s of our linearized theory. The coordinate-type physical degrees of freedom in ${}^3\breve{g}_{rs}$ depend upon the $r_{\bar a}$'s [18].

B. Comparison with the Standard Linearized Theory in Harmonic Gauges and its Associated Gravito-Electro-Magnetic Analogy.

Let us remark that our special 3-orthogonal gauge is not a member of the family of harmonic gauges, because, in coordinates adapted to our gauge, the condition $\partial_A\left(\sqrt{{}^4g}\, {}^4g^{AB}\right) = 0$ becomes

---

[17] All the quantities evaluated with the rescaled metric having $det\, |{}^3\breve{g}| = 1$ will be denoted with a $\breve{}$.

[18] In the 3-orthogonal gauge we have ${}^3\breve{g}_{rs} = {}^3\hat{g}^{diag}_{rs}$, see Eq.(A17).

$$\Big[\partial_A\,\Big(\sqrt{\hat{\gamma}}\,{}^4\hat{g}^{\tau A}\Big)\Big](\tau ,\vec{\sigma}) = \Big[\partial_\tau\,\Big(\sqrt{\hat{\gamma}} - 1\Big) - \sum_s \partial_s\, n_s\Big](\tau ,\vec{\sigma}) + O(r^2_{\bar{a}}) = 0 + O(r^2_{\bar{a}}),$$

$$\Big[\partial_A\,\Big(\sqrt{\hat{\gamma}}\,{}^4\hat{g}^{r A}\Big)\Big](\tau ,\vec{\sigma}) = \Big[\partial_\tau\, n_r - \partial_r\,\Big(\frac{2}{\sqrt{3}}\sum_{\bar{a}}[\gamma_{\bar{a}r} r_{\bar{a}})(\tau ,\vec{\sigma}) +$$

$$+ \frac{1}{2}\sum_u \gamma_{\bar{a}u}\int d^3\sigma_1 \frac{\partial^2_{1u} r_{\bar{a}}(\tau ,\vec{\sigma}_1)}{4\pi |\vec{\sigma} - \vec{\sigma}_1|}\Big)\Big](\tau ,\vec{\sigma}) + O(r^2_{\bar{a}}) \neq 0 + O(r^2_{\bar{a}}). \tag{4.4}$$

Therefore, on the solutions of Einstein's equations [15] our radiation gauge identifies a 4-coordinate system different from those compatible with the standard linearized theory coupled to matter.

This theory, in the post-Newtonian approximation, allows to make the *gravito-electro-magnetic* (GEM) re-formulation of Einstein's theory (see Refs. [16,31]), emphasizing the spin 1 aspects of this spin 2 theory over the Minkowski background in Cartesian coordinates.

By putting ${}^4g_{\mu\nu} = {}^4\eta_{\mu\nu} + h_{\mu\nu}$ with $h_{\mu\nu} = \bar{h}_{\mu\nu} + \frac{1}{2}\,{}^4\eta_{\mu\nu}\, h$ ($h = {}^4\eta^{\mu\nu}\, h_{\mu\nu}$, $\sqrt{{}^4g}\,{}^4g^{\mu\nu} = {}^4\eta^{\mu\nu} - \bar{h}^{\mu\nu}$, $\bar{h}^{\mu\nu} = {}^4\eta^{\mu\alpha}\,{}^4\eta^{\nu\beta}\, h_{\alpha\beta}$) Einstein's equations in presence of matter become $\Box\, \bar{h}_{\mu\nu} = {}^4\eta^{\alpha\beta}\, \partial_\alpha\, \partial_\beta\, \bar{h}_{\mu\nu} \overset{\circ}{=} -\frac{1}{k\,c}\, T_{\mu\nu}$ when the *Lorentz gauge condition* $\partial_\nu\, \bar{h}^{\mu\nu} = \partial_\nu\, \Big(\sqrt{{}^4g}\,{}^4g^{\mu\nu}\Big) = 0$, viz. *the conditions for harmonic coordinates (which include the post-Newtonian ones)*, are imposed. Even if in this approximation there is no back-reaction of the gravitational force on the sources, it allows to make an analogy with electromagnetism [23]. $\bar{h}_{\mu\nu}$ replaces the 4-potential A_μ with the coordinate transformation (changing the Christoffel symbols but not the curvature tensor) $\bar{h}^{'}_{\mu\nu} = \bar{h}_{\mu\nu} - \partial_\mu\, b_\nu - \partial_\nu\, b_\mu + {}^4\eta_{\mu\nu}\, \partial_\alpha\, b^\alpha$ corresponding to the gauge transformation $A^{'}_\mu = A_\mu + \partial_\mu\, \Lambda$. Maxwell equations $\Box\, A_\mu - \partial_\mu\, \partial_\alpha\, A^\alpha \overset{\circ}{=} -\frac{1}{c}\, j_\mu$ correspond to the linearized Einstein equations $\Box\, \bar{h}_{\mu\nu} + {}^4\eta_{\mu\nu}\, \partial_\alpha\, \partial_\beta\, \bar{h}^{\alpha\beta} - \partial_\mu\, \partial_\alpha\, \bar{h}^\alpha{}_\nu - \partial_\nu\, \partial_\alpha\, \bar{h}^\alpha{}_\mu \overset{\circ}{=} -\frac{2}{k}\, T_{\mu\nu}$. The *Lorentz gauge* $\partial_\mu\, A^\mu = 0$ (with residual gauge freedom given by functions Λ's satisfying $\Box\, \Lambda = 0$) with equations $\Box\, A_\mu \overset{\circ}{=} -\frac{1}{c}\, j_\mu$ corresponds to the *harmonic gauge (harmonic coordinates)* $\partial_\nu\, \bar{h}^{\mu\nu} = 0$ with $\Box\, \bar{h}_{\mu\nu} = -\frac{2}{k}\, T_{\mu\nu}$ (with residual freedom in the choice of coordinates having the b_μ's satisfying $\Box\, b^\mu = 0$; in Cartesian coordinates the equations of motion are decoupled one from the other).

The retarded solution $\bar{h}_{\mu\nu}(x^o, \vec{x}) \overset{\circ}{=} \frac{4G}{c^4}\, \int d^3x^{'}\, \frac{T_{\mu\nu}(x^o - |\vec{x} - \vec{x}^{'}|, \vec{x}^{'})}{|\vec{x} - \vec{x}^{'}|}$ allows to define:

i) the *gravito-electric* or Newton potential $\bar{h}_{oo} = -\epsilon\, \frac{4}{c^2}\, \Phi_{GEM}$ determined by $\rho = T^{oo}/c^2$ as the effective gravitational charge density;

ii) the *gravito-magnetic vector potential* (the shift functions) of Ref. [16,31] $\bar{h}_{oi} = \epsilon\, \frac{2}{c^2}\, A_{GEM\, i}$ determined by $j^i = T^{oi}/c$ as the effective gravitational current density;

iii) to disregard at the lowest order the 3-metric $\bar{h}_{ij} = O(c^{-4})$.

As shown in Ref. [31], in the harmonic gauge the *gravito-electro-magnetic 4-potential* $(\Phi_{GEM}, \vec{A}_{GEM})$ *satisfies the Lorentz gauge condition*

$$\partial_o \Phi_{GEM} + \vec{\partial} \cdot (\frac{1}{2} \vec{A}_{GEM}) = 0. \tag{4.5}$$

The factor $\frac{1}{2}$ derives from the fact that the effective gravito-magnetic charge ($Q_B = 2 M_{source}$) is twice the gravito-electric charge ($Q_E = M_{source}$). The *gravito-electro-magnetic fields* are

$\vec{E}_{GEM} = -\vec{\partial}\Phi_{GEM} - \partial_o(\frac{1}{2}\vec{A}_{GEM})$, $\vec{B}_{GEM} = \vec{\partial} \times \vec{A}_{GEM} \overset{def}{=} c\,\vec{\Omega}_{GEM}$: in the harmonic gauge (harmonic coordinates) they satisfy Maxwell equations

$$\begin{aligned}
&\vec{\partial} \cdot \vec{E}_{GEM} = 4\pi G\rho, \qquad \vec{\partial} \cdot (\frac{1}{2}\vec{B}_{GEM}) = 0,\\
&\vec{\partial} \times \vec{E}_{GEM} = -\partial_o(\frac{1}{2}\vec{B}_{GEM}), \qquad \vec{\partial} \times (\frac{1}{2}\vec{B}_{GEM}) = \partial_o\vec{E}_{GEM} + \frac{4\pi G}{c}\vec{j},
\end{aligned} \tag{4.6}$$

and one can consider as electro-magnetic-like gauge transformations the residual coordinate freedom.

In this approximation the GEM fields are determined *only by the matter* and vanish in absence of matter, viz. they are independent from the physical degrees of freedom (like $r_{\bar{a}}$, $\pi_{\bar{a}}$) of the gravitational field.

If we consider solutions of the homogeneous equations $\Box \bar{h}_{\mu\nu} \overset{\circ}{=} 0$, $\partial_\nu \bar{h}^{\mu\nu} = 0$, they are used to describe *gravitational waves* $\bar{h}_{\mu\nu} = a_{\mu\nu} e^{ik_\alpha x^\alpha}$ ($a_{\mu\nu} = const.$, $k_\mu k^\mu = 0$, $a_{\mu\nu} k^\nu = 0$) as the only independent degrees of freedom of the gravitational field. The residual freedom of coordinate transformations in the harmonic gauge may be used to get $a^{'}_{\mu\nu} = a_{\mu\nu} - c_\mu\, k_\nu - c_\nu\, k_\mu + {}^4\eta_{\mu\nu}\, c_\alpha\, c^\alpha$ with $a^{'}_{o\nu} = a^{'}_{\mu o} = 0$ and $a^{'}{}_\mu{}^\mu = a^{'}{}_i{}^i = 0$ through the choice $b^\mu = -i\, c^\mu\, e^{ik_\alpha x^\alpha}$ (solution of $\Box\, b^\mu = 0$). In *this completely fixed harmonic gauge (with non-diagonal 3-metric), named TT gauge,* only two independent degrees of freedom (polarization states) survive: they are the counterpart of $r_{\bar{a}}$, $\pi_{\bar{a}}$ of our completely fixed 3-orthogonal radiation gauge with $\pi_\phi(\tau, \vec{\sigma}) = 0$. If $k^\mu = (0; 0, \frac{\omega}{c}, \frac{\omega}{c})$, the only non-zero $a^{'}_{\mu\nu}$ of the transverse gravitational wave are $a^{'}_{xx}$, $a^{'}_{yy} = -a^{'}_{xx}$, $a^{'}_{xy} = a^{'}_{yx}$, and we get $ds^2 = \epsilon \Big[(dx^o)^2 - (1 + f_{xx})\, dx^2 - 2\, f_{xy}\, dxdy - (1 - f_{xx})\, dy^2 - dz^2\Big]$ with $e_+ = f_{xx} = a^{'}_{xx}\, cos\,(\frac{\omega}{c}\,(z - x^o) + \varphi)$, $e_\times = f_{xy} = a^{'}_{xy}\, cos\,(\frac{\omega}{c}\,(z - x^o) + \psi)$. In this gauge the lapse and shift functions n, n_i are all *zero* and there is *no GEM potential* ($\Phi_{GEM}, \vec{A}_{GEM}$) *coming from the transverse gravitational wave.*

Let us remark that in the electro-magnetic case on Minkowski space-time with Cartesian coordinates the radiation gauge $A_o = \vec{\partial} \cdot \vec{A} = 0$, with the transverse fields $\vec{A}_\perp$, $\vec{E}_\perp$ as the only physical degrees of freedom (DO's), is a particular case (name it *gauge T*) of the Lorentz gauge $\partial^\mu\, A_\mu = 0$, obtainable with a $\Lambda = -i\, c\, e^{i\, k \cdot x}$ with $k^2 = 0$ from $\Box\, \Lambda = 0$. Instead, the 4-coordinate system on the post-Minkowskian space-times connected with our radiation gauge for the gravitational waves is different from the 4-coordinate system which is reinterpreted as the TT harmonic gauge of the spin-2 theory on the background Minkowski space-time with Cartesian coordinates.

Instead in our non-harmonic radiation 3-orthogonal gauge with $\pi_\phi(\tau, \vec{\sigma}) = 0$, even in absence of matter the shift functions do *not* vanish and we can define

i) a *vanishing gravito-electric potential* $\Phi^{(rad)}_{GEM}(\tau, \vec{\sigma}) = 0 + O(r^2_{\bar{a}})$;

ii) a *gravito-magnetic vector potential* $\vec{A}^{(rad)}_{GEM}(\tau, \vec{\sigma}) = \{-\epsilon\, \frac{c^2}{2}\, n_r(\tau, \vec{\sigma})\}$ determined by the shift functions;

iii) a *gravito-electric field* $\vec{E}^{(rad)}_{GEM}(\tau, \vec{\sigma}) = -\partial_\tau\, [\frac{1}{2}\, \vec{A}^{(rad)}_{GEM}(\tau, \vec{\sigma})] = -\epsilon\, \frac{c^2}{4}\, \partial_\tau\, n_r(\tau, \vec{\sigma})$ *also* determined by the shift functions.

iv) a *gravito-magnetic field* $\vec{B}^{(rad)}_{GEM}(\tau, \vec{\sigma}) = \vec{\partial} \times \vec{A}^{(rad)}_{GEM}(\tau, \vec{\sigma}) \overset{def}{=} c\,\vec{\Omega}^{(rad)}_{GEM}(\tau, \vec{\sigma})$, with $\vec{\Omega}^{(rad)}_{GEM}(\tau, \vec{\sigma}) = \{\Omega^{(rad)}_{GEM,r} = c^{-1}\, \epsilon_{ruv}\, \frac{\partial n_v}{\partial \sigma^u}\}$ the gravito-magnetic precession angular velocity (connected with the precessional effects of Lense-Thirring or dragging of the inertial systems) again determined by the shift functions.

Since we are not in a harmonic gauge the *Lorentz condition is not satisfied* $\vec{\partial}\cdot(\frac{1}{2}\,\vec{A}^{(rad)}_{GEM})\neq 0$. *The analogy with Maxwell equations is partially lost in our completely fixed (radiation) gauge*: while the equations deriving from the existence of the potential $\vec{\partial}\cdot(\frac{1}{2}\,\vec{B}^{(rad)}_{GEM})=0$ and $\vec{\partial}\times\vec{E}^{(rad)}_{GEM}=-\partial_\tau\,(\frac{1}{2}\,\vec{B}^{(rad)}_{GEM})$ hold, the other two are not satisfied: $\vec{\partial}\cdot\vec{E}^{(rad)}_{GEM}\neq 0$, $\vec{\partial}\times(\frac{1}{2}\,\vec{B}^{(rad)}_{GEM})\neq\partial_\tau\,\vec{E}^{(rad)}_{GEM}$. This shows how the gravito-electric-magnetic analogy is *coordinate dependent*.

Finally, the coordinate transformation (a passive 4-diffeomorphism) from the $\Sigma^{(WSW)}_\tau$-adapted 4-coordinates $\tau,\vec{\sigma}$ of our completely fixed gauge to the 4-coordinates $x^\mu=x^\mu(\tau,\vec{\sigma})$ of the standard harmonic TT gauge and the relation between the DO's $r_{\bar{a}}(\tau,\vec{\sigma})$ and the TT polarizations e_+, $e_\times$ can be obtained as solution of the following system of partial differential equations:

$$
{}^4g_{\mu\nu}(x)=\frac{\partial x_\mu}{\partial\sigma_A}\,\frac{\partial x_\nu}{\partial\sigma_B}\,{}^4g_{AB}(\tau,\vec{\sigma}),\qquad {}^4g_{\mu\nu}=\begin{pmatrix}\epsilon & 0 & 0 & 0\\ 0 & -\epsilon\,e_+ & -\epsilon\,e_\times & 0\\ 0 & -\epsilon\,e_\times & +\epsilon\,e_+ & 0\\ 0 & 0 & 0 & -\epsilon\end{pmatrix},\tag{4.7}
$$

with ${}^4g_{AB}$ given by Eqs.(3.7).

If we write $x_\mu(\tau,\vec{\sigma})=f_\mu(\tau,\vec{\sigma})+g_\mu(\tau,\vec{\sigma})$ with $f_\mu=O(1)$ and $g_\mu=O(r_{\bar{a}})$ and if we require the asymptotic behaviour $f_\mu(\tau,\vec{\sigma})\rightarrow_{|\vec{\sigma}|\rightarrow\infty}x_{(\infty)\mu}(0)+u_\mu(P_{ADM})\,\tau+\epsilon_{r\,\mu}(u(P_{ADM}))\,\sigma^r$ with $u^\mu(p)=p^\mu/\sqrt{\epsilon\,p^2}$, then we get that

i) the four equations with ${}^4g_{oo}=\epsilon$, ${}^4g_{oi}=0$ can be solved to give $\frac{\partial x_\mu}{\partial\tau}$ in terms of $\frac{\partial x_\mu}{\partial\sigma^r}$;

ii) the system of partial differential equations ${}^4g_{13}={}^4g_{23}=0$, ${}^4g_{33}=-\epsilon$ and ${}^4g_{11}=-{}^4g_{22}$ have to be used for the determination of the $\vec{\sigma}$-dependence of the four functions $x_\mu(\tau,\vec{\sigma})$;

iii) the use of the solution of point ii) inside point i) allows to find $\frac{\partial x_\mu}{\partial\tau}$ and then, by integration, the functions $x_\mu(\tau,\vec{\sigma})$;

iv) finally the equations $e_+=-\epsilon\,{}^4g_{11}$ and $e_\times=-\epsilon\,{}^4g_{12}$ give the two polarizations e_+, $e_\times$ in terms of the Dirac observables $r_{\bar{a}}(\tau,\vec{\sigma})$.

C. Connection with the Post-Newtonian Approximation.

The standard post-Newtonian approximation [32] is applied to the gravitational field created by an isolated compact object, like the Earth, described by an energy-momentum tensor $T^{\mu\nu}$ [$T^{oo}=O(c^2)$, $T^{oi}=O(c)$, $T^{ij}=O(c^o)$]. It is a *weak field* (${}^4g={}^4\eta+{}^4h$) *near zone approximation* giving corrections to Newton gravity.

Following Ref. [33], if $V=\frac{GM}{R}$ is the Newton potential of the compact object (of mass M and radial dimensions $R_o\leq R$), the post-Newtonian (PN) expansion of the 4-metric ${}^4g_{\mu\nu}$ is a series in the dimensionless parameter $\zeta\approx\frac{\sqrt{V}}{c^2}\approx\frac{R}{cT}$ (T is a characteristic time of variation of the source). The 1PN approximation keeps the following terms in ${}^4g_{\mu\nu}$: ${}^4g_{oo}=\epsilon+\frac{a_{oo}}{c^2}+\frac{b_{oo}}{c^4}$, ${}^4g_{oi}=\frac{a_{oi}}{c^3}$, ${}^4g_{ij}=-\epsilon\delta_{ij}+\frac{a_{ij}}{c^2}$.

In Ref. [32] the PN approximation was given in a *PN 4-coordinate system* (adiabatic approximation of a Robertson-Walker line element appropriate to a homogeneous isotropic cosmological model containing the compact object), defined as a *local, quasi-Cartesian* 4-coordinate system at rest respect to the universe rest frame and in a *standard PN gauge* where the 4-metric has the form (we use the notation of $\Sigma_\tau^{(WSW)}$ -adapted coordinates) [19]

$$
\begin{aligned}
{}^4g_{\tau\tau} &= \epsilon e^{-\frac{2\epsilon}{c^2}V} = \epsilon[1 - \frac{2\epsilon}{c^2} V + O(c^{-4})],\\
{}^4g_{\tau r} &= -\frac{4\epsilon}{c^2}\, V_r + O(c^{-4}),\\
{}^4g_{rs} &= -\epsilon\, {}^3g_{rs} = -\epsilon\, \delta_{rs}\, (1 + \frac{U}{c^2}) + O(c^{-4}),
\end{aligned}
\qquad (4.8)
$$

with $U = V$ in the standard PN gauge. Here V is the Newton potential generated by the compact object.

Now our 4-coordinate system, in our completely fixed Hamiltonian gauge, is by definition in the rest frame of the universe (rest-frame instant form). In presence of matter, the linearization condition (as we will show in a future paper on tetrad gravity plus a perfect fluid, where it will be shown that both the lapse and shift functions depend on matter so that $n \neq 0$)) requires $T^{AB} = O(r_{\bar a})$ and we have

$$
\begin{aligned}
{}^4g_{\tau\tau} &= \epsilon\, [(-\epsilon + n)^2 - {}^3g_{rs}\, n^r n^s] = \epsilon(1 - 2\epsilon\, n[matter]) + O(r^2_{\bar a}),\\
{}^4g_{\tau r} &= -\epsilon\, {}^3g_{rs}\, n^s = -\epsilon\, n_r[r_{\bar a}, matter] + O(r^2_{\bar a}),\\
{}^4g_{rs} &= -\epsilon\, {}^3g_{rs} = -\epsilon\, \delta_{rs}\, (1 + k_r[r_{\bar a}, matter]) + O(r^2_{\bar a}),\\
\phi &= [det\, {}^3g]^{1/12} = (1 + \frac{U}{4c^2}) + O(r^2_{\bar a}).
\end{aligned}
\qquad (4.9)
$$

Therefore, in our gauge the *Newton potential is the lapse function*, $V = c^2\, n[matter]$ (and the equation determining the lapse tends to the Poisson equation), the *gravito-magnetic potentials are the shift functions*, $V_r = 4c^2\, n_r[r_{\bar a}, matter]$, and the *Lichnerowicz equation for the conformal factor ϕ amounts to a determination of U*, which may not coincide with the Newton potential like it happens in the standard PN gauge. By solving the linearized equations for the DO's (see next Section), $\Box\, r_{\bar a} = Z_{\bar a}[matter]$, i.e. by considering their

---

[19] In Refs. [34,33] there are various definitions of 4-coordinate systems. The *standard PN gauge* is said to correspond to the conditions $\partial_s\, {}^4g_{\tau s} - \frac{1}{2}\, \partial_\tau\, {}^4g_{ss} = O(c^{-5})$, $\partial_s\, {}^4g_{rs} - \frac{1}{2}\, ({}^4g_{ss} - {}^4g_{\tau\tau}) = O(c^{-4})$. Instead the *algebraic spatial isotropy condition* of Ref. [33], $-{}^4g_{\bar\tau\bar\tau}\, {}^4\tilde g_{rs} = \delta_{rs} + O(c^{-4})$, contains both the harmonic and the standard post-Newtonian gauges. Finally a *ADM Hamiltonian gauge*, needed to include 2.5PN gravitational radiation reaction, is ${}^3K = 0$, $\partial_s\, {}^3g_{rs} - \frac{1}{3}\, \partial_r\, {}^3g_{ss} = 0$.

Lienard-Wiechert retarded solution without incoming free radiation, we can recover a form like the one of Eqs.(4.8), which does not depend explicitly on the DO's $r_{\bar{a}}$ [20].

Let us now consider the relation of our linearized radiation gauge with Ref. [35], where the Galileo generally covariant formulation of Newtonian gravity, depending on 27 fields, was obtained as a limit $c \rightarrow \infty$ on the ADM action of metric gravity.

The final action of Ref. [35] with general Galileo covariance depends on the 26 fields A_o, α_o, A_r, α_r, ${}^3\check{g}_{rs}$, $\check{\gamma}_{rs}$, $\check{\beta}_{rs}$ (after having put the field $\Theta = 1$ by rescaling the absolute time) . There are 18 first class constraints and 8 pairs of second class ones. It turns out that α_o, A_r, α_r, three components of ${}^3\check{g}_{rs}$, one component of the momentum conjugate to ${}^3\check{g}_{rs}$, the trace $\check{\beta}^T$ and the longitudinal $\check{\beta}^L_r$ parts of $\check{\beta}_{rs}$ in its TT decomposition, and the longitudinal $\check{\gamma}^L_r$ part of $\check{\gamma}_{rs}$ are Hamiltonian gauge variables, while A_o (the Newton potential) and the remaining components of ${}^3\check{g}_{rs}$, $\check{\gamma}_{rs}$, $\check{\beta}_{rs}$ are determined, together with their conjugate momenta, by the second class constraints. There are no propagating dynamical degrees of freedom. The gauge variables describe the inertial forces in arbitrary accelerated non-Galilean reference frames.

To make a comparison with the results of Sections 6 and 7 of Ref. [35] the starting point is the following parametrization of the 4-metric of Eq.(3.7) (we show only the 26 terms which appear in the Newtonian action)

$$\begin{aligned}
{}^4g_{\tau\tau} &= \epsilon\,[1 - \frac{2A_o}{c^2} + \frac{2\alpha_o}{c^4} + O(c^{-6})] = \\
&= \epsilon\,(1 - 2\epsilon\, n) + O(r^2_{\bar{a}}),\\
\\
{}^4g_{\tau r} &= \epsilon\,[A_r + \frac{\alpha_r}{c^2} + O(c^{-4})] = -\epsilon\, n_r + O(r^2_{\bar{a}}),\\
\\
{}^4g_{rs} &= -\epsilon\,{}^3g_{rs} = -\epsilon[{}^3\check{g}_{rs} + \frac{\check{\gamma}_{rs}}{c^2} + \frac{\check{\beta}_{rs}}{c^4} + O(c^{-6})] = \\
&= -\epsilon\,\delta_{rs}\,(1 + k_r) + O(r^2_{\bar{a}}),
\end{aligned}$$

$$\Downarrow$$

$$\begin{aligned}
n[matter] &= \epsilon\,[\frac{A_o}{c^2} + \frac{\alpha_o}{c^4} + O(c^{-6}, r^2_{\bar{a}})],\\
n_r[r_{\bar{a}}, matter] &= -A_r - \frac{1}{c^2}\alpha_r + O(c^{-4}, r^2_{\bar{a}}),
\end{aligned}$$

---

[20] Let us also remark that strictly speaking to reach Newton gravity one usually performs a double limit: i) the zero-curvature limit $r_{\bar{a}}, \pi_{\bar{a}} \rightarrow 0$, eliminating the genuine degrees of freedom of the gravitational field, which do not exist in Newtonian gravity; ii) the $1/c$ expansion. However the zero curvature limit is not consistent with Einstein's equations with matter and must be replaced (in a linearized theory) with the restriction of the DO's $r_{\bar{a}}$, $\pi_{\bar{a}}$ to the Lienard-Wiechert solution. Then the $1/c$ expansion will kill the curvature for $c \rightarrow \infty$.

$$\delta_{rs}\,(1+k_r[r_{\bar a}, matter]) = {}^3\check{g}_{rs} + \frac{\check{\gamma}_{rs}}{c^2} + \frac{\check{\beta}_{rs}}{c^4} + O(c^{-6}, r^2_{\bar a}). \tag{4.10}$$

Instead the PN approximation (4.8) implies

$$A_o = \epsilon V, \quad A_r = 0, \quad \alpha_r = -4V_r, \quad {}^3\check{g}_{rs} = \delta_{rs}, \quad \check{\gamma}_{rs} = U\,\delta_{rs}, \tag{4.11}$$

which are consistent with the gauge freedom of Ref. [35] (it is a possible gauge of the general Galileo covariant description of Newtonian gravity).

V. THE HAMILTON EQUATIONS AND THEIR SOLUTION.

In this Section we shall study the Hamilton equations in the preferred 3-orthogonal gauge and we will verify that their solution produces a linearized solution of Einstein's equations. Since, as said in the Introduction, the admissible WSW hyper-surfaces must admit a generalized Fourier transform, we shall use it on our linearized WSW hyper-surfaces (where it coincides with the ordinary Fourier transform on R^3 at the lowest order) to study the Hamilton equations.

As said in the Introduction, in the rest frame instant form of dynamics on WSW hyper-surfaces the Dirac Hamiltonian is $H_D = \hat{E}_{ADM} + \vec{\lambda}(\tau) \cdot \hat{\vec{P}}_{ADM}$. However we can add the gauge fixings $\hat{J}^{\tau r}_{ADM} \approx 0$ [i.e. the vanishing of the weak ADM boost generators of Eq.(A12)] to the three first class constraints $\hat{P}^r_{ADM} \approx 0$ with the consequence $\vec{\lambda}(\tau) = 0$. This means that we have to eliminate the *internal 3-center of mass* of the universe (it is put in the centroid $x^\mu_{(\infty)}(\tau)$ near spatial infinity used as origin of the 3-coordinates on WSW hyper-surfaces), i.e. 3 pairs of global degrees of freedom among the DO's $r_{\bar{a}}(\tau, \vec{\sigma})$, $\pi_{\bar{a}}(\tau, \vec{\sigma})$, reducing them to canonical variables relative to the internal 3-center of mass.

Like in the case of Klein-Gordon and electro-magnetic fields and like for every isolated system treated in the rest frame instant form [5], we should find the canonical transformation from the canonical variables $r_{\bar{a}}(\tau, \vec{\sigma})$, $\pi_{\bar{a}}(\tau, \vec{\sigma})$ to a canonical basis containing the canonical internal 3-center of mass Q^r_{ADM}, weakly equal to the Møller internal 3-center of energy $R^r_{ADM} = -\hat{J}^{\tau r}_{ADM}/\hat{E}_{ADM}$, $\hat{P}^r_{ADM} (\approx 0)$ as the conjugate momentum, and the *internal relative variables* $R_{\bar{a}}$, $\Pi_{\bar{a}}$. Usually [36,37], like in the Klein-Gordon case, we should start with a naive 3-center of mass X^r_{ADM}, conjugate to $\hat{P}^r_{ADM}$, we should find the canonical variables $R^{'}_{\bar{a}}$, $\Pi^{'}_{\bar{a}}$ relative to it and then we should use the Gartenhaus-Schwartz transformation to find [36] the canonical variables with respect to Q^r_{ADM}. Then we should add the gauge fixings $Q^r_{ADM} \approx R^r_{ADM} = -\hat{J}^{\tau r}_{ADM}/\hat{E}_{ADM} \approx 0$, which put the internal 3-centers in the centroid $x^\mu_{(\infty)}$ origin of the 3-coordinates on the WSW hyper-surfaces and implies $H_D = \hat{E}_{ADM}$, to the rest-frame condition $\hat{P}^r_{ADM} \approx 0$ and we should go to Dirac brackets with the result $R_{\bar{a}} \equiv R^{'}_{\bar{a}}$, $\Pi_{\bar{a}} \equiv \Pi^{'}_{\bar{a}}$. Presumably also the final relative variables have to satisfy Eqs.(C2).

However, since also the determination of the naive 3-center of mass X^r_{ADM} of gravity is not trivial, we will fix this final gauge freedom simply by putting equal to zero the Dirac multipliers, $\vec{\lambda}(\tau) = 0$ as a pre-gauge fixing condition compatible with $\hat{J}^{\tau r}_{ADM} \approx 0$.

We can now study the Hamilton equations for the independent canonical degrees of freedom $r_{\bar{a}}(\tau, \vec{\sigma})$, $\pi_{\bar{a}}(\tau, \vec{\sigma})$ of the gravitational field, only restricted by Eqs.(2.15), in our completely fixed 3-orthogonal gauge generated by the Hamiltonian $\hat{E}_{ADM}$ of Eq.(3.1). These Hamilton equations replace the equations $\Box \bar{h}_{\mu\nu} = 0$, $\partial_\nu \bar{h}^{\mu\nu} = 0$ of the standard linearized theory in the harmonic gauge coordinates.

A. The First Half of Hamilton Equations.

The first half of the Hamilton equations associated with the Hamiltonian (3.1) yield the following expression for the velocities $\partial_\tau r_{\bar{a}}(\tau, \vec{\sigma})$

$$\begin{aligned}
\partial_\tau r_{\bar a}(\tau,\vec\sigma) &\overset{\circ}{=} \{r_{\bar a}(\tau,\vec\sigma), \hat E_{ADM}\} = \frac{24\pi G}{c^3}\pi_{\bar a}(\tau,\vec\sigma) + \\
&+ \frac{24\pi G}{c^3}\sum_{u,r,\bar b,(a)} \delta_{(a)u}\Big[\gamma_{\bar a u}\gamma_{\bar b r}\int d^3\sigma_1 \mathcal{T}^{(o)u}_{(a)r}(\vec\sigma,\vec\sigma_1)\pi_{\bar b}(\tau,\vec\sigma_1) + \\
&+ \gamma_{\bar a r}\gamma_{\bar b u}\int d^3\sigma_1 \pi_{\bar b}(\tau,\vec\sigma_1)\mathcal{T}^{(o)u}_{(a)r}(\vec\sigma_1,\vec\sigma)\Big] + \\
&+ \frac{12\pi G}{c^3}\sum_{u,r,s,\bar b}\gamma_{\bar a r}\gamma_{\bar b s}\Big[\sum_{(a)}\int d^3\sigma_1 \mathcal{T}^{(o)u}_{(a)r}(\vec\sigma,\vec\sigma_1)\int d^3\sigma_2 \mathcal{T}^{(o)u}_{(a)s}(\vec\sigma_1,\vec\sigma_2) + \\
&+ \sum_{v(a)(b)}(\delta_{(a)v}\delta_{(b)u} - \delta_{(a)u}\delta_{(b)v})\int d^3\sigma_1 \mathcal{T}^{(o)u}_{(a)r}(\vec\sigma,\vec\sigma_1)\int d^3\sigma_2 \mathcal{T}^{(o)v}_{(b)s}(\vec\sigma_1,\vec\sigma_2)\Big]\pi_{\bar b}(\tau,\vec\sigma_2) + \\
&+ \frac{1}{L}O(r^2_{\bar a}). \qquad (5.1)
\end{aligned}$$

By using the expression of $\mathcal{T}^{(o)u}_{(a)r}$ given in the last two lines of Eqs.(2.13) it can be checked that the two terms linear in the $\mathcal{T}$'s vanish due to a factor $\delta_{(a)u}(1-\delta_{(a)u})$. After some calculations it turns out that the two terms bilinear in the $\mathcal{T}$'s give the same result, so that the final expression for the velocities is

$$\begin{aligned}
\partial_\tau r_{\bar a}(\tau,\vec\sigma) \overset{\circ}{=} \frac{24\pi G}{c^3}\pi_{\bar a}(\tau,\vec\sigma) + \frac{6\pi G}{c^3}\sum_{\bar b rs}\gamma_{\bar a r}\gamma_{\bar b s}\sum_{uv,u\neq v}[1-2(\delta_{ur}+\delta_{vr})][1-2(\delta_{us}+\delta_{vs})] \\
\int_{-\infty}^{\sigma^u} d\sigma^u_1 \int_{-\infty}^{\sigma^u_1} d\sigma^u_2 \int_{\sigma^v}^{\infty} d\sigma^v_1 \int_{\sigma^v_1}^{\infty} d\sigma^v_2 \frac{\partial^4 \pi_{\bar b}(\tau, \sigma^u_2\sigma^v_2\sigma^{k\neq u,v})}{(\partial\sigma^r_2)^2(\partial\sigma^s_2)^2} + \frac{1}{L}O(r^2_{\bar a}). \qquad (5.2)
\end{aligned}$$

In obtaining this result we have made integrations by parts justified by the asymptotic vanishing of $\pi_{\bar a}(\tau,\vec\sigma)$ and we have used $\partial_x\theta(x,y) = -\partial_y\theta(x,y)$, $\partial_x\delta(x,y) = -\partial_y\delta(x,y)$ assumed valid on linearized conformally flat WSW hyper-surfaces.

To invert these equations, to get the momenta in terms of the velocities, we shall assume the validity of the Fourier transform on the linearized WSW CMC-hyper-surfaces. They are conformal to R^3, i.e. they are Euclidean plus corrections of order $O(r_{\bar a})$, irrelevant when acting on the functions $r_{\bar a}(\tau,\vec\sigma)$, $\pi_{\bar a}(\tau,\vec\sigma)$; for the scalar product we have ${}^3\hat g_{rs}k^rh^s = \vec k\cdot\vec h + O(r_{\bar a})$. As a consequence functions of order $O(r_{\bar a})$ are considered as functions over R^3 at the lowest order of approximation.

By using the Fourier transform defined in Appendix C, Eqs.(5.2) become Eqs.(C4)

$$\partial_\tau \tilde r_{\bar a}(\tau,\vec k) = \sum_{\bar b} A_{\bar a\bar b}(\vec k)\tilde\pi_{\bar b}(\tau,\vec k), \qquad (5.3)$$

with the matrix $A_{\bar a\bar b}(\vec k)$ and its inverse given in Eqs.(C5) and (C6), respectively. Let us remark that, notwithstanding $A_{\bar a\bar b}(\vec k)$ diverges for $k^r \to 0$, Eqs.(5.3) are well defined if $\tilde\pi_{\bar a}(\tau,\vec k) \rightarrow_{\vec k\rightarrow 0} (k^1k^2k^3)^{2+\epsilon}\, \tilde f_{\bar a}(\vec k)$, $\epsilon > 0$. *This condition is stronger of the requirement* (C2), i.e. of the Fourier transform of Eqs.(2.15).

As a consequence we get $[\epsilon_{\bar a\bar b} = -\epsilon_{\bar b\bar a},\ \epsilon_{\bar 1\bar 2} = 1]$

$$
\begin{aligned}
\tilde{\pi}_{\bar{a}}(\tau,\vec{k}) &= \sum_{\bar{b}} A^{-1}_{\bar{a}\bar{b}}(\vec{k})\partial_\tau \tilde{r}_{\bar{a}}(\tau,\vec{k}),\\
\pi_{\bar{a}}(\tau,\vec{\sigma}) &= \sum_{\bar{b}} \int d^3\sigma_1 G_{\bar{a}\bar{b}}(\vec{\sigma}-\vec{\sigma}_1)\partial_\tau r_{\bar{b}}(\tau,\vec{\sigma}_1),\\
G_{\bar{a}\bar{b}}(\vec{\sigma}-\vec{\sigma}_1) &= \int \frac{d^3k}{(2\pi)^3} A^{-1}_{\bar{a}\bar{b}}(\vec{k}) e^{i\vec{k}\cdot(\vec{\sigma}-\vec{\sigma}_1)} =\\
&= -\frac{c^3}{8\pi G}\int \frac{d^3w_1 d^3w_2 d^3w_3}{(4\pi)^3|\vec{\sigma}-\vec{w}_1|\,|\vec{w}_1-\vec{w}_2|\,|\vec{w}_2-\vec{w}_3|}\Big[\delta_{\bar{a}\bar{b}}\frac{\partial^6\delta^3(\vec{w}_3-\vec{\sigma}_1)}{(\partial w_3^1)^2(\partial w_3^2)^2(\partial w_3^3)^2}+\\
&+\frac{1}{2}\sum_{r,s,t,\bar{c},\bar{d}} \epsilon_{\bar{a}\bar{c}}\gamma_{\bar{c}r}\epsilon_{\bar{b}\bar{d}}\gamma_{\bar{d}s}(2\delta_{tr}-1)(2\delta_{ts}-1)\frac{\partial^6\delta^3(\vec{w}_3-\vec{\sigma}_1)}{(\partial w_3^r)^2(\partial w_3^s)^2(\partial w_3^t)^2}\Big]. \qquad (5.4)
\end{aligned}
$$

The final result of the inversion is

$$
\begin{aligned}
\pi_{\bar{a}}(\tau,\vec{\sigma}) &= -\frac{c^3}{8\pi G}\int \frac{d^3\sigma_1 d^3\sigma_2 d^3\sigma_3}{(4\pi)^3|\vec{\sigma}-\vec{\sigma}_1|\,|\vec{\sigma}_1-\vec{\sigma}_2|\,|\vec{\sigma}_2-\vec{\sigma}_3|}\\
&\Big[\frac{\partial^6\partial_\tau r_{\bar{a}}(\tau,\vec{\sigma}_3)}{(\partial\sigma_3^1)^2(\partial\sigma_3^2)^2(\partial\sigma_3^3)^2}+\\
&+\frac{1}{2}\sum_{r,s,t,\bar{b},\bar{c},\bar{d}} \epsilon_{\bar{a}\bar{c}}\gamma_{\bar{c}r}\epsilon_{\bar{b}\bar{d}}\gamma_{\bar{d}s}(2\delta_{tr}-1)(2\delta_{ts}-1)\frac{\partial^6\partial_\tau r_{\bar{b}}(\tau,\vec{\sigma}_3)}{(\partial\sigma_3^r)^2(\partial\sigma_3^s)^2(\partial\sigma_3^t)^2}\Big]. \qquad (5.5)
\end{aligned}
$$

Eqs.(5.5) satisfies the condition (2.15) automatically.

B. The Second Half of Hamilton Equations.

Let us now study the second half of Hamilton equations associated with the Hamiltonian (3.1). After some calculations we get

$$
\begin{aligned}
\partial_\tau\pi_{\bar{a}}(\tau,\vec{\sigma}) &\overset{\circ}{=} \{\pi_{\bar{a}}(\tau,\vec{\sigma}),\hat{E}_{ADM}\} =\\
&= \frac{c^3}{24\pi G}\sum_r \frac{\partial^2 r_{\bar{a}}(\tau,\vec{\sigma})}{\partial(\sigma^r)^2} - \frac{c^3}{12\pi G}\sum_{r,\bar{b}} \gamma_{\bar{a}r}\gamma_{\bar{b}r}\frac{\partial^2 r_{\bar{b}}(\tau,\vec{\sigma})}{(\partial\sigma^r)^2} -\\
&- \frac{c^3}{48\pi G}\sum_{\bar{b}rs}\gamma_{\bar{a}r}\gamma_{\bar{b}s}\int \frac{d^3\sigma_1}{4\pi|\vec{\sigma}-\vec{\sigma}_1|}\frac{\partial^4 r_{\bar{b}}(\tau,\vec{\sigma}_1)}{(\partial\sigma_1^r)^2(\partial\sigma_1^s)^2} + \frac{c^3}{GL^2}O(r_{\bar{a}}^2). \qquad (5.6)
\end{aligned}
$$

The Fourier transform of Eqs.(5.6) is given in Eqs. (C7) and (C8), with the remarkable result $\partial_\tau\,\tilde{\pi}_{\bar{a}}(\tau,\vec{k}) \overset{\circ}{=} -|\vec{k}|^2\sum_{\bar{b}} A^{-1}_{\bar{a}\bar{b}}(\vec{k})\,\tilde{r}_{\bar{a}}(\tau,\vec{k})$.

This *implies* $\int_{-\infty}^{\infty} d\sigma^r\, r_{\bar{a}}(\tau,\vec{\sigma}) = 0$, i.e. Eq.(2.16), *as the simplest way to get consistency* between Eqs. (5.6) and (2.15).

Let us remark that, as shown in Eqs. (D9) of Appendix D, our Hamilton equations imply the *satisfaction of the remaining Einstein equations* ${}^4\hat{R}_{rs}(\tau,\vec{\sigma}) \overset{\circ}{=} 0$.

Eqs. (5.4) and (5.6) imply

$$
\begin{aligned}
\partial_\tau^2 r_{\bar a}(\tau,\vec\sigma) \stackrel{\circ}{=}\ & \frac{24\pi G}{c^3}\partial_\tau \pi_{\bar a}(\tau,\vec\sigma) + \\
&+ \frac{6\pi G}{c^3}\sum_{\bar b rs}\gamma_{\bar a r}\gamma_{\bar b s}\sum_{uv,u\neq v}[1-2(\delta_{ur}+\delta_{vr})][1-2(\delta_{us}+\delta_{vs})] \\
&\int_{-\infty}^{\sigma^u} d\sigma_1^u \int_{-\infty}^{\sigma^v} d\sigma_1^v \int_{\sigma_1^u}^{\infty} d\sigma_2^u \int_{\sigma_1^v}^{\infty} d\sigma_2^v (\partial_r^2\partial_s^2\partial_\tau \pi_{\bar b})(\tau,\sigma_2^u,\sigma_2^v,\sigma^{k\neq u,v}) + \\
&+ \frac{1}{L}O(r^2_{\bar a}), \\
& \\
&\Downarrow \\
& \\
\Box r_{\bar a}(\tau,\vec\sigma) =\ & [\partial_\tau^2 - \sum_r \partial_r^2] r_{\bar a}(\tau,\vec\sigma) \stackrel{\circ}{=} \\
\stackrel{\circ}{=}\ & \frac{1}{2}\sum_{\bar b r u}\gamma_{\bar a r}(4\gamma_{\bar b r}-\gamma_{\bar b u})\int \frac{d^3\sigma_1}{4\pi|\vec\sigma-\vec\sigma_1|}\frac{\partial^4 r_{\bar b}(\tau,\vec\sigma_1)}{(\partial\sigma_1^r)^2(\partial\sigma_1^u)^2} + \\
&+ \frac{1}{8}\sum_{\bar b rs}\sum_{\bar c tw}\sum_{uv}(-2\delta_{\bar b\bar c}+4\gamma_{\bar b t}\gamma_{\bar c t}-\gamma_{\bar b t}\gamma_{\bar c w})\gamma_{\bar a r}\gamma_{\bar b s}(1-\delta_{uv})[1-2(\delta_{ur}+\delta_{vr})] \\
&[1-2(\delta_{us}+\delta_{vs})]\int_{-\infty}^{\sigma^u} d\sigma_1^u \int_{-\infty}^{\sigma^v} d\sigma_1^v \int_{\sigma_1^u}^{\infty} d\sigma_2^u \int_{\sigma_1^v}^{\infty} d\sigma_2^v \\
&\int d^3\sigma_3\, \frac{[(\partial_{3r})^2(\partial_{3s})^2(\partial_{3w})^2(\partial_{3t})^2 r_{\bar c}](\tau,\vec\sigma_3)}{4\pi\sqrt{(\sigma_3^u-\sigma_2^u)^2+(\sigma_3^v-\sigma_2^v)^2+(\sigma_3^{k\neq u,v}-\sigma^{k\neq u,v})^2}} + \frac{1}{L^2}O(r^2_{\bar a}). \qquad (5.7)
\end{aligned}
$$

As shown in Appendix C, Eq.(C10), all the terms in the second member of Eq.(5.7) cancel, so that we find that the DO's $r_{\bar a}(\tau,\vec\sigma)$ satisfy the wave equation. Actually, as shown in Eq.(C10), the Fourier transform of the equation of motion is

$$
\begin{aligned}
&\ddot{\tilde r}_{\bar a}(\tau,\vec k) + |\vec k|^2 \tilde r_{\bar a}(\tau,\vec k) = 0, \\
& \\
&\Downarrow \\
& \\
&\Box\, r_{\bar a}(\tau,\vec\sigma) \stackrel{\circ}{=} 0. \qquad (5.8)
\end{aligned}
$$

Therefore, in our radiation gauge we get *the wave equation* $\Box\, r_{\bar a}(\tau,\vec\sigma)\stackrel{\circ}{=}0$ *for the DO's.* Notwithstanding the presence of gravito-magnetism, which, as said, should imply the anisotropy of light propagation, we get an isotropic propagation of gravitational waves in the radiation gauge.

The complicated form (3.1), (C13), of the weak ADM energy shows that *in our gauge we do not have the conventional description of a massless spin two particle over Minkowski space-time like in the TT harmonic gauge.* This fact is connected with the lack of a coordinate-independent notion of gravitational energy density, which has no counterpart in the theory of massless spin two particles in Minkowski space-time. Instead we will see in

Eqs.(5.11) and (5.12) that the weak ADM 3-momentum and angular momentum do admit a standard particle interpretation.

The solutions of the Hamilton equations for the DO's $r_{\bar a}(\tau,\vec\sigma)$ and $\pi_{\bar a}(\tau,\vec\sigma)$ are given in Eq.(C11), (C12)

$$
\begin{aligned}
r_{\bar a}(\tau,\vec\sigma) &= \int \frac{d^3k}{(2\pi)^3}\Big(C_{\bar a}(\vec k)e^{-i\,|\vec k|\tau} + C^*_{\bar a}(-\vec k)e^{i\,|\vec k|\tau}\Big)\, e^{+i\,\vec k\cdot\vec\sigma},\\
\pi_{\bar a}(\tau,\vec\sigma) &= \int \frac{d^3k}{(2\pi)^3}\,\tilde\pi_{\bar a}(\tau,\vec k)\, e^{i\,\vec k\cdot\vec\sigma} = \int \frac{d^3k}{(2\pi)^3}\,\sum_{\bar b} A^{-1}_{\bar a\bar b}\partial_\tau \tilde r_{\bar b}(\tau,\vec k)\, e^{i\,\vec k\cdot\vec\sigma},\\
&= \int \frac{d^3k}{(2\pi)^3}\,\sum_{\bar b} A^{-1}_{\bar a\bar b}(\vec k)\,\Big[-i|\vec k|C_{\bar b}(\vec k)e^{-i|\vec k|\tau} + i|\vec k|C^*_{\bar b}(-\vec k)e^{i|\vec k|\tau}\Big]\, e^{i\,\vec k\cdot\vec\sigma},
\end{aligned}
\tag{5.9}
$$

with the functions $C_{\bar a}(\vec k)$ *vanishing* as $(k^1\,k^2\,k^3)^\epsilon$ for $k^r \rightarrow 0$.

C. Special Solutions for the Background-Independent Gravitational Waves.

Special solutions are the following *plane waves* whose 3-momentum $\vec h$ cannot lie in any coordinate plane $[\vec h \neq \vec 0; (1,0,0); (0,1,0); (0,0,1); (1,1,0); (1,0,1); (0,1,1)]$

$$
\begin{aligned}
& r_{\bar a}(\tau,\vec\sigma) = C_{\bar a}\, e^{i\,(\vec h\cdot\vec\sigma - |\vec h|\,\tau)}, \quad C_{\bar a} = const.,\\
&\qquad \tilde r_{\bar a}(\tau,\vec k) = (2\pi)^3\, C_{\bar a}\, e^{-i\,|\vec h|\,\tau}\,\delta^3(\vec k - \vec h),\\
& \pi_{\bar a}(\tau,\vec\sigma) = -i\,|\vec h|\,\sum_{\bar b} A^{-1}_{\bar a\bar b}(\vec h)\, C_{\bar b}\, e^{i(\vec h\cdot\vec\sigma - |\vec h|\,\tau)},\\
&\qquad \tilde\pi_{\bar a}(\tau,\vec k) = -i\,(2\pi)^3\,|\vec h|\,\sum_{\bar b} A^{-1}_{\bar a\bar b}(\vec h)\, C_{\bar b}\, e^{-i\,|\vec h|\,\tau}\,\delta^3(\vec k - \vec h), \quad \tilde\pi_{\bar a}(\tau,\vec 0) = 0,
\end{aligned}
\tag{5.10}
$$

which verify Eqs. (2.15) and (2.16). Here $\pi_{\bar a}$ has been evaluated with Eq.(5.5) and Eqs.(C4) and (C7) are valid also at $\vec k = 0$.

Hoever these solutions do not satisfy the rest frame conditions $\hat P^r_{ADM} \approx 0$, which restrict the solutions to globally outgoing or ingoing wave packets.

Before looking for these solutions, we shall give the form of the weak ADM charges (A12) in the radiation gauge with our solution of the constraints (see Eqs.(3.1) and (C13) for $\hat E_{ADM}$). The weak ADM 3-momentum and spin of Eqs.(A12) assume the same simple form in terms of Fourier transformed quantities as for free massless fields due to the fact that we are in an instant form of dynamics. This is not true for the ADM boosts, which have a complicated form like the ADM energy.

For the 3-momentum, due to exact cancellations, we get

$$
\begin{aligned}
\hat{P}^r_{ADM} &= - \int d^3\sigma \sum_{\bar c} \Big[\partial_r\, r_{\bar c}(\tau, \vec\sigma) - \\
&- \gamma_{\bar c r} \Big(- \sum_{\bar a u} \gamma_{\bar a u} \int d^3\sigma_1\, \frac{\partial^2_{1u}\, r_{\bar a}(\tau, \vec\sigma_1)}{4\pi\, |\vec\sigma - \vec\sigma_1|} - 2 \sum_{\bar a} \gamma_{\bar a r}\, \partial_r\, r_{\bar a}(\tau, \vec\sigma) \Big) \Big]\, \pi_{\bar c}(\tau, \vec\sigma) + \\
&+ \int d^3\sigma d^3\sigma_1 \sum_{\bar c s} \Big(\gamma_{\bar c s} \\
&\Big[\sum_u \frac{1}{\sqrt{3}} \Big(\frac{1}{2} \sum_{\bar a m} \gamma_{\bar a m}\, \partial_r \int d^3\sigma_2\, \frac{\partial^2_{2m}\, r_{\bar a}(\tau, \vec\sigma_2)}{4\pi\, |\vec\sigma - \vec\sigma_1|} + \sum_{\bar a} \gamma_{\bar a u}\, \partial_r\, r_{\bar a}(\tau, \vec\sigma) \Big) \\
&\delta^u_{(a)}\, \mathcal{T}^{(o)u}_{(a)s}(\vec\sigma, \vec\sigma_1) + \\
&+ \sum_{uv} \Big(\delta^r_u\, \frac{1}{\sqrt{3}} \Big(\frac{1}{2} \sum_{\bar a m} \gamma_{\bar a m}\, \partial_v \int d^3\sigma_2\, \frac{\partial^2_{2m}\, r_{\bar a}(\tau, \vec\sigma_2)}{4\pi\, |\vec\sigma - \vec\sigma_1|} + \sum_{\bar a} \gamma_{\bar a u}\, \partial_v\, r_{\bar a}(\tau, \vec\sigma) \Big) + \\
&+ \delta^r_v\, \frac{1}{\sqrt{3}} \Big(\frac{1}{2} \sum_{\bar a m} \gamma_{\bar a m}\, \partial_u \int d^3\sigma_2\, \frac{\partial^2_{2m}\, r_{\bar a}(\tau, \vec\sigma_2)}{4\pi\, |\vec\sigma - \vec\sigma_1|} + \sum_{\bar a} \gamma_{\bar a v}\, \partial_u\, r_{\bar a}(\tau, \vec\sigma) \Big) \Big) \\
&\delta^u_{(a)}\, \mathcal{T}^{(o)v}_{(a)s}(\vec\sigma, \vec\sigma_1) \Big] \Big)\, \pi_{\bar c}(\tau, \vec\sigma_1) = \\
&= - \int d^3\sigma \sum_{\bar c} \pi_{\bar c}(\tau, \vec\sigma)\, \partial_r\, r_{\bar c}(\tau, \vec\sigma) = \\
&= i \int \frac{d^3k}{(2\pi)^3} \sum_{\bar c} \tilde\pi_{\bar c}(\tau, \vec k)\, k^r\, \tilde r_{\bar c}(\tau, -\vec k),
\end{aligned}
\qquad (5.11)
$$

where we used $\mathcal{T}^{(o)u}_{(u)s} = 0$, see Eq.(2.13).

For the angular momentum, due to similar cancellations, we get

$$
\begin{aligned}
\hat{J}^{rs}_{ADM} &= \int d^3\sigma \sum_{\bar c} \pi_{\bar c}(\tau, \vec\sigma)\, (\sigma^r\, \partial_s - \sigma^s\, \partial_r)\, r_{\bar c}(\tau, \vec\sigma) = \\
&= - \int \frac{d^3k}{(2\pi)^3} \sum_{\bar c} \tilde\pi_{\bar c}(\tau, \vec k)\, \Big(k^r\, \frac{\partial}{\partial k^s} - k^s\, \frac{\partial}{\partial k^r} \Big)\, \tilde r_{\bar c}(\tau, -\vec k).
\end{aligned}
\qquad (5.12)
$$

Eqs. (5.11) and (5.12) reflect the fact that we are in an instant form of the dynamics. Instead for the boosts we get a complicated expression like for the energy (see Eqs.(2.13) for the kernels)

$$
\begin{aligned}
\hat{J}^{\tau r}_{ADM} &= \epsilon \int d^3\sigma\, \sigma^r\, \Big(\frac{c^3}{16\pi\, G} \Big[\frac{1}{3} \sum_r \Big(\frac{1}{2} \Big(\sum_{\bar a m} \gamma_{\bar a m}\, \partial_r \int d^3\sigma_2\, \frac{\partial^2_{2m}\, r_{\bar a}(\tau, \vec\sigma_2)}{4\pi\, |\vec\sigma - \vec\sigma_1|} \Big)^2 - \\
&- \sum_{\bar a} \Big(\partial_r\, r_{\bar a}(\tau, \vec\sigma) \Big)^2 + \sum_{\bar a m} \gamma_{\bar a m}\, \partial_r \int d^3\sigma_2\, \frac{\partial^2_{2m}\, r_{\bar a}(\tau, \vec\sigma_2)}{4\pi\, |\vec\sigma - \vec\sigma_1|} \sum_{\bar b} \gamma_{\bar b r}\, \partial_r\, r_{\bar b}(\tau, \vec\sigma) + \\
&+ 2 \Big(\sum_{\bar b} \gamma_{\bar b r}\, \partial_r\, r_{\bar b}(\tau, \vec\sigma) \Big)^2 \Big) \Big] - \frac{6\pi\, G}{c^3} \Big[2 \sum_{\bar a} \pi^2_{\bar a}(\tau, \vec\sigma) + \\
&+ 4 \sum_{\bar b u} \gamma_{\bar b u}\, \pi_{\bar b}(\tau, \vec\sigma) \int d^3\sigma_1 \sum_m \delta^u_{(a)}\, \mathcal{T}^{(o)u}_{(a)m}(\vec\sigma, \vec\sigma_1) \sum_{\bar a} \gamma_{\bar a m}\, \pi_{\bar a}(\tau, \vec\sigma_1) +
\end{aligned}
$$

$$
\begin{aligned}
&+ \int d^3\sigma_1 d^3\sigma_2 \Big(\sum_u \sum_m \mathcal{T}^{(o)u}_{(a)m}(\vec{\sigma}, \vec{\sigma}_1) \sum_{\bar{a}} \gamma_{\bar{a}m}\, \pi_{\bar{a}}(\tau, \vec{\sigma}_1) \\
&\quad \sum_s \mathcal{T}^{(o)u}_{(a)s}(\vec{\sigma}, \vec{\sigma}_2) \sum_{\bar{b}} \gamma_{\bar{b}s}\, \pi_{\bar{b}}(\tau, \vec{\sigma}_2) + \\
&+ \sum_{uv} (\delta^u_{(b)}\, \delta^v_{(a)} - \delta^u_{(a)}\, \delta^v_{(b)}) \sum_m \mathcal{T}^{(o)u}_{(a)m}(\vec{\sigma}, \vec{\sigma}_1) \sum_{\bar{a}} \gamma_{\bar{a}m}\, \pi_{\bar{a}}(\tau, \vec{\sigma}_1) \\
&\quad \sum_s \mathcal{T}^{(o)v}_{(b)s}(\vec{\sigma}, \vec{\sigma}_2) \sum_{\bar{b}} \gamma_{\bar{b}s}\, \pi_{\bar{b}}(\tau, \vec{\sigma}_2) \Big) \Big] \Big) - \\
&- \frac{\epsilon\, c^3}{8\pi\, G} \int d^3\sigma \sum_{uv} \delta^r_u\, (\delta_{uv} - 1)\, \frac{1}{3} \Big(- \frac{1}{4} \sum_{\bar{a}m} \gamma_{\bar{a}m}\, \partial_u \int d^3\sigma_2\, \frac{\partial^2_{2m}\, r_{\bar{a}}(\tau, \vec{\sigma}_2)}{4\pi\, |\vec{\sigma} - \vec{\sigma}_1|} + \\
&+ \sum_{\bar{a}} (\gamma_{\bar{a}v} - \gamma_{\bar{a}u})\, \partial_u\, r_{\bar{a}}(\tau, \vec{\sigma}) \Big) \\
&\quad \Big(- \sum_{\bar{a}n} \gamma_{\bar{a}n} \int d^3\sigma_2\, \frac{\partial^2_{2n}\, r_{\bar{a}}(\tau, \vec{\sigma}_2)}{4\pi\, |\vec{\sigma} - \vec{\sigma}_1|} + 2 \sum_{\bar{b}} \gamma_{\bar{b}u}\, r_{\bar{b}}(\tau, \vec{\sigma}) \Big) = \\
&= \epsilon \int d^3\sigma \sigma^r \Big(\frac{c^3}{16\pi G} \sum_s [2(\partial_s q(\tau, \vec{\sigma}))^2 - \frac{1}{3} \sum_{\bar{b}} (\partial_s r_{\bar{b}}(\tau, \vec{\sigma}))^2 + \\
&- \frac{2}{\sqrt{3}}\, \partial_s q(\tau, \vec{\sigma}) \sum_{\bar{b}} \gamma_{\bar{b}s} \partial_s r_{\bar{b}}(\tau, \vec{\sigma}) + \frac{2}{3} (\sum_{\bar{b}} \gamma_{\bar{b}s} \partial_s r_{\bar{b}})^2 (\tau, \vec{\sigma})] - \frac{12\pi G}{c^3} [\sum_{\bar{a}} \pi^2_{\bar{a}}(\tau, \vec{\sigma}) + \\
&+ 4 \int d^3\sigma_1 \int d^3\sigma_2 \sum_{u,m,s,(a)} \mathcal{T}^{(o)u}_{(a)m}(\vec{\sigma}, \vec{\sigma}_1) \sum_{\bar{b}} \gamma_{\bar{b}m} \pi_{\bar{b}}(\tau, \vec{\sigma}_1) \mathcal{T}^{(o)u}_{(a)s}(\vec{\sigma}, \vec{\sigma}_2) \sum_{\bar{c}} \gamma_{\bar{c}s} \pi_{\bar{c}}(\tau, \vec{\sigma}_2)] \Big) + \\
&- \epsilon \frac{c^3}{8\pi G} \int d^3\sigma \sum_v (\delta_{rv} - 1)(2q + \frac{2}{\sqrt{3}} \sum_{\bar{a}} \gamma_{\bar{a}r} r_{\bar{a}})(\frac{1}{2}\, \partial_r q + \frac{1}{\sqrt{3}} \sum_{\bar{b}} (\gamma_{\bar{b}v} - \gamma_{\bar{b}r}) \partial_r r_{\bar{b}})(\tau, \vec{\sigma}) + \\
&+ O(r^3_{\bar{a}}).
\end{aligned}
\tag{5.13}
$$

The weak ADM 3-momentum of the solutions (5.9), (C11), has the form

$$
\hat{P}^r_{ADM} = 2 \sum_{\bar{b}\bar{c}} \int \frac{d^3k}{(2\pi)^3}\, A^{-1}_{\bar{c}\bar{b}}(\vec{k})\, |\vec{k}|\, k_r\, C_{\bar{b}}(\vec{k})\, C^*_{\bar{c}}(\vec{k}) \approx 0, \tag{5.14}
$$

so that its vanishing is a condition on the Fourier coefficients $C_{\bar{a}}(\vec{k})$, which, for instance, *cannot be satisfied by the plane waves* (5.10).

A *class of solutions* of the Hamilton equations with the correct vanishing behaviour at spatial infinity, which, besides Eqs.(C2), also satisfies the rest frame condition $\hat{P}^r_{ADM} = 0$ is obtained by taking coefficients $C_{\bar{a}}(\vec{k}) = C_{\bar{a}}\, \frac{(k_1 k_2 k_3)^2}{|\vec{k}|}\, e^{-\vec{k}^2}$ ($C_{\bar{a}} = const.$) in Eq.(5.9)

$$
\begin{aligned}
r_{\bar{a}}(\tau, \vec{\sigma}) &= C_{\bar{a}} \int \frac{d^3k}{(2\pi)^3}\, \frac{(k_1 k_2 k_3)^2}{|\vec{k}|}\, e^{-\vec{k}^2} \Big[e^{-i(|\vec{k}|\, \tau - \vec{k} \cdot \vec{\sigma})} + e^{i(|\vec{k}|\, \tau - \vec{k} \cdot \vec{\sigma})} \Big] = \\
&= - \frac{4\, C_{\bar{a}}}{(2\pi)^2}\, \frac{\partial^2}{\partial^2 \sigma_1}\, \frac{\partial^2}{\partial^2 \sigma_2}\, \frac{\partial^2}{\partial^2 \sigma_3}\, \frac{(1 + |\vec{\sigma}|^2 - \tau^2)}{[1 + (|\vec{\sigma}| + \tau)^2]\, [1 + (|\vec{\sigma}| - \tau)^2]},
\end{aligned}
$$

$$
\pi_{\bar{a}}(\tau, \vec{\sigma}) = -i \int \frac{d^3k}{(2\pi)^3}\, \frac{(k_1 k_2 k_3)^2}{|\vec{k}|}\, e^{-\vec{k}^2}
$$

$$|\vec{k}| \sum_{\bar{b}} A^{-1}_{\bar{a}\bar{b}}(\vec{k})\, C_{\bar{b}} \Big[e^{-i(|\vec{k}|\,\tau - \vec{k}\cdot\vec{\sigma})} - e^{i(|\vec{k}|\,\tau - \vec{k}\cdot\vec{\sigma})} \Big],$$

$$\begin{aligned}
n_r(\tau, \vec{\sigma}) &= i\,\sqrt{3} \frac{4\pi G}{c^3} \sum_{\bar{a}vuc} \gamma_{\bar{a}v} (1 - \delta_{uc})[1 - 2(\delta_{uv} + \delta_{cv})][1 - 2(\delta_{ur} + \delta_{cr})] \\
&\quad \int \frac{d^3k}{(2\pi^3)}\, (k_1 k_2 k_3)^2 e^{-\vec{k}^2} \frac{k_r k_v^2}{k_u^2 k_c^2} \sum_{\bar{b}} A^{-1}_{\bar{a}\bar{b}}(\vec{k})\, C_{\bar{b}}\, sin\,(|\vec{k}|\,\tau) e^{i\vec{k}\cdot\vec{\sigma}}, \\
\hat{E}_{ADM} &= \sum_{\bar{a}\bar{b}} \alpha_{\bar{a}\bar{b}}\, C_{\bar{a}} C_{\bar{b}}, \\
\hat{P}^r_{ADM} &= 0, \\
\hat{J}^r_{ADM} &= \frac{1}{2}\, \epsilon^{ruv}\, \hat{J}^{uv}_{ADM} = \sum_{\bar{a}\bar{b}} \beta^r_{\bar{a}\bar{b}}\, C_{\bar{a}} C_{\bar{b}}.
\end{aligned} \tag{5.15}$$

The two constants $C_{\bar{1}}$, $C_{\bar{2}}$ have to be expressed in terms of the two boundary constants $M = \hat{E}_{ADM}$, $S = |\hat{\vec{J}}_{ADM}|$ defining the mass and spin of the post-Minkowskian Einstein space-time. The solution (5.15) goes like $|\vec{\sigma}|^{-8}$ for $|\vec{\sigma}| \rightarrow \infty$, i.e. much faster than the behaviour (1.2). It should be possible to find solutions saturating Eqs.(1.2), namely such that by using Eqs.(3.7) we should get ${}^3g_{rs} \rightarrow_{r\rightarrow\infty} (1 + \frac{\hat{E}_{ADM}}{2\,r})\delta_{rs}$, $n_r = -\epsilon\, {}^4g_{\tau r} \rightarrow_{r\rightarrow\infty} O(r^{-\epsilon})$. By comparison, in presence of compact matter the expected solution should have a post-Newtonian behaviour ${}^4g_{\tau\tau} \rightarrow_{r\rightarrow\infty} 1 - \frac{2M}{r}$, ${}^4g_{\tau r} \rightarrow_{r\rightarrow\infty} 4\epsilon_{ruv} S^u \frac{\sigma^v}{r^3}$, ${}^4g_{rs} \rightarrow_{r\rightarrow\infty} (1 + \frac{2M}{r})\delta_{rs}$ (see Eq.(19.5) of Ref. [22]).

Therefore, in absence of matter, the rest-frame condition destroys the transversality property of the TT harmonic gauge plane waves.

VI. THE TIME-LIKE GEODESICS, THE GEODESIC DEVIATION EQUATION AND THE POLARIZATION OF GRAVITATIONAL WAVES IN THE RADIATION GAUGE.

A. The Geodesic Equation.

Let us consider a time-like geodesic with affine parameter s, $\sigma^A(s) = \Big(\tau(s); \sigma^u(s)\Big)$. After the linearization in the radiation gauge the *geodesic equation* becomes (see Eqs.(C16) of Appendix C for Fourier transforms)

$$\begin{aligned}
\frac{d^2\tau(s)}{ds^2} &= -{}^4\hat{\Gamma}^\tau_{rs}(\sigma(s))\frac{d\sigma^r(s)}{ds}\frac{d\sigma^s(s)}{ds} + O(r^2_{\bar{a}}),\\
\frac{d^2\sigma^u(s)}{ds^2} &= -\partial_\tau n_u(\sigma(s))(\frac{d\tau(a)}{ds})^2 - 2\,{}^4\hat{\Gamma}^u_{\tau r}(\sigma(s))\frac{d\tau(s)}{ds}\frac{d\sigma^r(s)}{ds} - {}^3\hat{\Gamma}^u_{rs}(\sigma(s))\frac{d\sigma^r(s)}{ds}\frac{d\sigma^s(s)}{ds} +\\
&+ O(r^2_{\bar{a}}),
\end{aligned} \tag{6.1}$$

where Eq.(D5) has to be used for the Christoffel symbols.

If we parametrize the geodesic as $\sigma^A(s) = a^A + b^A s + f^A(s|r_{\bar{a}}, \pi_{\bar{a}}]$, where $a^A + b^A\, s$ is the flat geodesic, we get at the lowest order $\frac{d^2 f^\tau(s)}{ds^2} \approx -b^r b^s\, {}^4\hat{\Gamma}^\tau_{rs}(\sigma(s)) + O(r^2_{\bar{a}}) \approx -b^r b^s\, {}^4\hat{\Gamma}^\tau_{rs}(a + bs) + O(r^2_{\bar{a}})$. Therefore the solution to the first of Eqs.(6.1) is

$$\begin{aligned}
\tau(s) &= a^\tau + b^\tau s - \int_0^s ds_1 \int_0^{s_1} ds_2 {}^4\hat{\Gamma}^\tau_{ru}(a + bs_2) b^r b^u\\
p^\tau(s) &= \frac{d\tau(s)}{ds} = b^\tau - \int_0^s ds_2 {}^4\hat{\Gamma}^\tau_{ru}(a + bs_2) b^r b^u \qquad \text{with} \qquad {}^4\hat{\Gamma}^\tau_{ru}(a + bs_2) \sim O(r_{\bar{a}}).
\end{aligned} \tag{6.2}$$

Since the tangent $p^A(s) = \frac{d\sigma^A(s)}{ds}$ to the geodesic must satisfy either ${}^4g^{AB}\, p_A\, p_B = \epsilon$ (time-like geodesic with s as proper time) or ${}^4g^{AB}\, p_A\, p_B = 0$ (null geodesic) at the initial time, the constant b^τ is determined in terms of the b^r's and of the initial data. Therefore we will have $b^\tau = b^\tau_{(o)} + b^\tau_{(1)}$ with $b^\tau_{(1)} = O(r_{\bar{a}})$. Therefore Eq.(6.2) may be rewritten in the form

$$\begin{aligned}
\tau(s) &= a^\tau + [b^\tau_{(o)} + b^\tau_{(1)}]\, s - \int_0^s ds_1 \int_0^{s_1} ds_2 {}^4\hat{\Gamma}^\tau_{ru}(a + b_{(o)} s_2) b^r b^u\\
p^\tau(s) &= \frac{d\tau(s)}{ds} = b^\tau_{(o)} + b^\tau_{(1)} - \int_0^s ds_2 {}^4\hat{\Gamma}^\tau_{ru}(a + b_{(o)} s_2)\, b^r b^u.
\end{aligned} \tag{6.3}$$

The spatial part of Eqs.(6.1) becomes

$$\begin{aligned}
\frac{d^2 f^u(s)}{ds^2} &= -\partial_\tau n_u(a + b_{(o)} s)(b^\tau_{(o)})^2 - 2\,{}^4\hat{\Gamma}^u_{\tau r}(a + b_{(o)} s) b^\tau_{(o)} b^r - {}^3\hat{\Gamma}^u_{rt}(a + b_{(o)} s) b^r b^t + O(r^2_{\bar{a}})\\
\sigma^u(s) &= a^u + b^u - \int_0^s ds_1 \int_0^{s_1} ds_2 \Big[(b^\tau_{(o)})^2 - 2\,{}^4\hat{\Gamma}^u_{\tau r}(a + b_{(o)} s_2) b^\tau_{(o)}\, b^r - {}^3\hat{\Gamma}^u_{rt}(a + b_{(o)} s_2) b^r b^t\Big]\\
p^u(s) &= \frac{d\sigma^u(s)}{ds} = b^u - \int_0^s ds_2 \Big[(b^\tau_{(o)})^2 - 2\,{}^4\hat{\Gamma}^u_{\tau r}(a + b_{(o)} s_2) b^\tau_{(o)}\, b^r - {}^3\hat{\Gamma}^u_{rt}(a + b_{(o)} s_2) b^r b^t\Big].
\end{aligned} \tag{6.4}$$

In the time-like case at $s=0$, where $\tau(0)=a^\tau$, we have the condition

$$
\begin{aligned}
{}^4g_{AB}p^Ap^B|_{s=0} &= \epsilon\Big[(p^\tau(s))^2 - 2n_r(s,r_{\bar a}(s))p^r(s)p^\tau(s) - [\delta_{rt} - \epsilon\,{}^4h_{rt}(s,r_{\bar a}(s))p^r(s)p^t(s)]\Big]|_{s=0} = \\
&= \epsilon,
\end{aligned}
$$

$$
1 = (b^\tau)^2 - 2b^\tau b^r n_r(a^\tau, r_{\bar a}(a^\tau)) - [\delta_{rt} - \epsilon\,{}^4h_{rt}(a^\tau, r_{\bar a}(a^\tau))b^r b^t], \quad at\, s=0, \qquad (6.5)
$$

whose solution is

$$
\begin{aligned}
b^\tau &= b^\tau_{(o)} + b^\tau_{(1)} = b^r n_r(a, r_{\bar a}(a)) \pm \sqrt{(b^r n_r(a^\tau, r_{\bar a}(a^\tau)))^2 + 1 + [\delta_{rt} - \epsilon\,{}^4h_{rt}(a^\tau, r_{\bar a}(a^\tau))]\, b^r b^t}, \\
b^\tau_{(o)} &= \pm\sqrt{1+\delta_{rs}b^r b^s}, \\
b^\tau_{(1)} &= n_r(a^\tau, r_{\bar a}(a^\tau))\, b^r \mp \frac{\epsilon}{2}\, \frac{{}^4h_{rt}(a^\tau, r_{\bar a}(a^\tau))b^r b^t}{\sqrt{1+\delta_{rs}b^r b^s}} + O(r^2_{\bar a}). \qquad (6.6)
\end{aligned}
$$

If we choose $b^r=0$, we can take the solution $f^\tau(s)=0$ so that we get $\tau(s) = a^\tau + b^\tau s \rightarrow_{b^\tau=1} s + a^\tau$.

With the choice $b^\tau = 1$, $b^r = 0$, $a^\tau = 0$, we get $\tau(s) = s$ and $\frac{d^2 f^u(s)}{ds^2} = -\partial_\tau n_u(\tau(s), \vec\sigma(s) = \vec a + \vec f(s)) = 0$ at the lowest order, with the solution $f^u(s) = -\int^s ds' n_u(s', a^r) + O(r^2_{\bar a})$. Therefore, with $b^\tau = 1$, $b^r = 0$, $a^\tau = 0$ we get the following expression for the geodesics

$$
\begin{aligned}
\sigma^A(s) &= \Big(\tau(s) = s;\ \sigma^u(s) = a^u - \int_0^s dw\, n_u(w, a^v)\Big) + O(r^2_{\bar a}), \\
\frac{d\sigma^A(s)}{ds} &= \Big(1;\ -n_u(s, a^v)\Big) + O(r^2_{\bar a}). \qquad (6.7)
\end{aligned}
$$

This is the *trajectory of a massive test particle in our coordinates.*

B. The Geodesic Deviation Equation.

If we consider two nearby freely falling particles following two nearby geodesics (6.7) such that $\sigma^A_1(0) = (0;\vec 0)$ and $\sigma^A_2(0) = (0; a^u)$ with an infinitesimal a^u, we get for $\triangle x^A(s) = \sigma^A_2(s) - \sigma^A_1(s)$: $\triangle x^\tau(s) = 0$, $\triangle x^r(s) \approx [\delta^{ru} - \int_0^s dw\, \frac{\partial n^r(w,a^v)}{\partial a^u}]\, a^u$. If $a^u = \delta^{ur}\, a$ is their coordinate distance, then their *proper distance* is $\triangle l = \sqrt{-\epsilon\, {}^4g_{AB}\, \triangle x^A\, \triangle x^B} = \triangle x^r\, \sqrt{{}^3g_{rr}} = (1 + \frac{1}{2}k_r)\, \triangle x^r + O(r^2_{\bar a}) \approx [1 + \frac{1}{2}k_r - \int_0^s dw\, \frac{\partial n^r(w,a^v)}{\partial a^r}]\, a + O(r^2_{\bar a})$ by using Eqs.(3.7) [in a TT harmonic gauge one gets $\triangle l \approx [1 + h^{TT}_{rr}]\, a$ if r is one of the two polarization directions].

As a consequence the connecting vector $\triangle x^A(s)$ satisfies

$$
\triangle x^\tau(s) = 0, \qquad \frac{d^2 \triangle x^r(s)}{ds^2} = -\frac{d}{ds}\Big(\frac{\partial n^r(s, a^v)}{\partial a^u}\Big)\, a^u. \qquad (6.8)
$$

This is the *geodesic deviation equation* along the geodesic $\sigma^A_1(s)$ of the form (6.7): it shows explicitly the action of the *tidal forces* in our 4-coordinate system. More in general, by using the equation

$$\frac{d\sigma^C(s)}{ds}\,{}^4\nabla_C \triangle x^A(s) = \frac{d\triangle x^A(s)}{ds} + \frac{d\sigma^C(s)}{ds}\,{}^4\hat{\Gamma}^A_{CE}(s,a^u)\triangle x^E(s) + O(r^2_{\bar{a}}) =$$
$$= \frac{d\triangle x^A(s)}{ds} + {}^4\hat{\Gamma}^A_{\tau E}(s,a^u)\triangle x^E(s) + O(r^2_{\bar{a}}), \qquad (6.9)$$

the geodesic deviation equation [22,24] takes the form

$$\frac{d\sigma^B(s)}{ds}\,{}^4\nabla_B\Big(\frac{d\sigma^C(s)}{ds}\,{}^4\nabla_C\,\triangle x^A(s)\Big) = -{}^4\hat{R}^A{}_{BCD}(\sigma(s))\,\triangle x^C(s)\frac{d\sigma^B(s)}{ds}\frac{d\sigma^D(s)}{ds},$$

$$\Downarrow$$

$$\frac{d^2\triangle x^A(s)}{ds^2} + 2\,{}^4\hat{\Gamma}^A_{\tau E}(s,a^u)\frac{d\triangle x^E(s)}{ds} + \frac{d\,{}^4\hat{\Gamma}^A_{\tau E}(s,a^u)}{ds}\triangle x^E(s) = -{}^4\hat{R}^A{}_{\tau C\tau}(s,a^u)\triangle x^C(s),$$

$$\Downarrow$$

$$\frac{d^2\triangle x^\tau(s)}{ds^2} = 0 + O(r^2_{\bar{a}}), \quad \Rightarrow \quad \triangle x^\tau(s) = cs + d,$$

$$\frac{d^2\triangle x^r(s)}{ds^2} + 2\Big[{}^4\hat{\Gamma}^r_{\tau\tau}(s,a^u)c + {}^4\hat{\Gamma}^r_{\tau s}(s,a^u)\frac{d\triangle x^s(s)}{ds} + \frac{d\,{}^4\hat{\Gamma}^r_{\tau\tau}(s,a^u)}{ds}(cs+d) +$$
$$+ \frac{d\,{}^4\hat{\Gamma}^r_{\tau s}(s,a^u)}{ds}\triangle x^s(s) = -{}^4\hat{R}^r{}_{\tau s\tau}(s,a^u)\triangle x^s(s) + O(r^2_{\bar{a}}). \qquad (6.10)$$

If we take $\triangle x^\tau(s) = 0$ as it happens for the geodesic (6.7), i.e. $c = d = 0$, the geodesic deviation equation becomes

$$\frac{d^2\triangle x^r(s)}{ds^2} + 2\,{}^4\hat{\Gamma}^r_{\tau s}(s,a^u)\frac{d\triangle x^s(s)}{ds} + \frac{d\,{}^4\hat{\Gamma}^r_{\tau s}(s,a^u)}{ds}\triangle x^s(s) =$$
$$= -{}^4\hat{R}^r{}_{\tau s\tau}(s,a^u)\triangle x^s(s) + O(r^2_{\bar{a}}). \qquad (6.11)$$

If we write $\triangle x^r(s) = \triangle_{(o)}x^r(s) + \triangle_{(1)}x^r(s)$, with $\triangle_{(o)}x^r(s) = (o; a^u)$ the flat deviation and $\triangle_{(1)}x^r(s) = \Big(0; -\int_0^s dw\,[n^u(w,a^v) - n^u(w,\vec{0})]\Big) \approx \Big(0; -\int_0^s dw\,\frac{\partial n^u(w,a^v)}{\partial a^v}|_{a^v=0}\,a^v\Big) = O(r_{\bar{a}})$, Eq.(6.11) becomes

$$\frac{d^2}{ds^2}\Big(\triangle_{(o)}x^r(s) + \triangle_{(1)}x^r(s)\Big) = \frac{d^2\,\triangle_{(1)}x^r(s)}{ds^2} =$$
$$= -2\,{}^4\hat{\Gamma}^r_{\tau s}(s,a^u)\frac{d\triangle_{(o)}x^s(s)}{ds} - \frac{d\,{}^4\hat{\Gamma}^r_{\tau s}(s,a^u)}{ds}\triangle_{(o)}x^s(s) -$$
$$-{}^4\hat{R}^r{}_{\tau s\tau}(s,a^u)\triangle_{(o)}x^s(s) + O(r^2_{\bar{a}}). \qquad (6.12)$$

The use of Eqs.(D5) and (D7) allows to check that Eq.(6.12) coincides with Eq.(6.8).

C. The Tidal Forces Generated by the DO's on Test Particles.

Eq.(6.8) can be used to see the effect of the tidal forces generated by the two DO's $r_{\bar a}(\tau, \vec\sigma)$ of our radiation gauge, replacing the two polarizations of the TT harmonic gauge, on a sphere of test particles surrounding a test particle sitting in the origin ($\tau = 0; \vec\sigma = \vec 0$) at $s = 0$ and described by the geodesic $\sigma^A_o(s) = (s; \sigma^u_o(s) = -\int_0^s dw\, n^u(w, \vec 0)\,)$. If we define $a^u = a\,\eta^u$ with ${}^3g_{uv}\,\eta^u\,\eta^v = 1 + O(r_{\bar a})$ and $a << 1$ (the radius of the sphere), the test particles on the sphere at $s = 0$ will follow the geodesics $\sigma^A(s) = (s; \sigma^u(s) = a\,\eta^u - \int_0^s dw\, n^u(s, a\,\eta^v)\,)$. The connecting vectors $\triangle x^A(s) = \triangle_{(o)}x^A(s) + \triangle_{(1)}x^A(s) = \sigma^A(s) - \sigma^A_o(s)$ satisfy Eq.(6.8), namely

$$\triangle x^\tau(s) = 0, \qquad \frac{d^2 \triangle_{(1)} x^r(s)}{ds^2} = -\frac{d}{ds}\Big(\frac{\partial n^r(s, a^v)}{\partial a^v}|_{a^v=0}\Big)\, a\,\eta^v + O(r^2_{\bar a}). \tag{6.13}$$

We have to solve these equations with the shift functions (3.5) evaluated on the solutions (5.15), which are not transverse plane waves due to the rest frame conditions (this forces us to consider a sphere of test particles).

The quasi-Shanmugadhasan canonical basis (1.2) adapted to our gauge is not unique, because there is still the freedom associated with the numerical constants $\gamma_{\bar a r}$ (their variation amounts to a redefinition of the DO's). As a consequence of the conditions stated after Eqs.(1.5), the $\gamma_{\bar a r}$'s may be written in term of an angle ψ with $0 \leq \psi \leq 2\pi$

$$\begin{aligned}
\gamma_{\bar 1 r} &= \Big(\frac{cos\,\psi}{\sqrt6} - \frac{sin\,\psi}{\sqrt2}, \frac{cos\,\psi}{\sqrt6} + \frac{sin\,\psi}{\sqrt2}, -\sqrt{\frac{2}{3}}\,cos\,\psi\Big),\\
\gamma_{\bar 2 r} &= \Big(-\frac{cos\,\psi}{\sqrt2} - \frac{sin\,\psi}{\sqrt6}, \frac{cos\,\psi}{\sqrt2} - \frac{sin\,\psi}{\sqrt6}, \sqrt{\frac{2}{3}}\,cos\,\psi\Big).
\end{aligned} \tag{6.14}$$

By defining the DO's, i.e.our polarizations, with the convention $\psi = 0$, we get $\gamma_{\bar 1 r} = \frac{1}{\sqrt6}\,(1, 1, -2)$, $\gamma_{\bar 2 r} = \frac{1}{\sqrt2}\,(-1, 1, 0)$. Then Eqs.(3.7) imply

$$-\epsilon\,{}^3\hat g^{(\bar 1)}_{rs}(\tau, \vec\sigma) = \delta_{rs} + {}^4h^{(\bar 1)}_{rs}(\tau, \vec\sigma) \overset{def}{=} \delta_{rs}\,[1 + k^{(\bar 1)}_r(\tau, \vec\sigma)] + O(r^2_{\bar a}),$$

$$\begin{aligned}
k^{\bar 1}_1(\tau, \vec\sigma) = k^{\bar 1}_2(\tau, \vec\sigma) &= -\epsilon\,\frac{\sqrt2}{3}\,[r_{\bar 1}(\tau, \vec\sigma) - \frac{1}{2}\int d^3\sigma_1\,\frac{(\partial^2_{1,1} + \partial^2_{1,2} + \partial^2_{1,3})\,r_{\bar 1}(\tau, \vec\sigma_1)}{4\pi|\vec\sigma - \vec\sigma_1|}],\\
k^{\bar 1}_3(\tau, \vec\sigma) &= -\epsilon\,\frac{\sqrt2}{3}\,[-2 r_{\bar 1}(\tau, \vec\sigma) - \frac{1}{2}\int d^3\sigma_1\,\frac{(\partial^2_{1,1} + \partial^2_{1,2} + \partial^2_{1,3})\,r_{\bar 1}(\tau, \vec\sigma_1)}{4\pi|\vec\sigma - \vec\sigma_1|}],
\end{aligned}$$

$$-\epsilon\,{}^3\hat g^{(\bar 2)}_{rs}(\tau, \vec\sigma) = \delta_{rs} + {}^4h^{(\bar 2)}_{rs}(\tau, \vec\sigma) \overset{def}{=} \delta_{rs}\,[1 + k^{(\bar 2)}_r(\tau, \vec\sigma)] + O(r^2_{\bar a}),$$

$$k^{\bar 2}_1(\tau, \vec\sigma) = -\epsilon\,\sqrt{\frac{2}{3}}\,[-r_{\bar 2}(\tau, \vec\sigma) + \frac{1}{2}\int d^3\sigma_1\,\frac{(\partial^2_{1,1} - \partial^2_{1,2})\,r_{\bar 1}(\tau, \vec\sigma_1)}{4\pi|\vec\sigma - \vec\sigma_1|}],$$

$$
\begin{aligned}
k_2^{\bar{2}}(\tau,\vec{\sigma}) &= -\epsilon\,\sqrt{\frac{2}{3}}\,[r_{\bar{2}}(\tau,\vec{\sigma}) + \frac{1}{2}\int d^3\sigma_1\,\frac{(\partial^2_{1,1}-\partial^2_{1,2})\,r_{\bar{1}}(\tau,\vec{\sigma}_1)}{4\pi|\vec{\sigma}-\vec{\sigma}_1|}],\\
k_3^{\bar{2}}(\tau,\vec{\sigma}) &= -\epsilon\,\sqrt{\frac{2}{3}}\,[0 + \frac{1}{2}\int d^3\sigma_1\,\frac{(\partial^2_{1,1}-\partial^2_{1,2})\,r_{\bar{1}}(\tau,\vec{\sigma}_1)}{4\pi|\vec{\sigma}-\vec{\sigma}_1|}].
\end{aligned}
\tag{6.15}
$$

If we consider our (non transverse) solution given by the wave packet (5.15) and we solve Eqs.(6.13) numerically for a sphere of particle at rest around the origin of the 3-coordinates on a WSW hyper-surface, we obtain the two 3-dimensional deformation patterns replacing the usual 2-dimensional ones for the polarization in the TT harmonic gauge:

i) in figure 1 there is the deformation pattern for the case $C_{\bar{1}} \neq 0$, $C_{\bar{2}} = 0$, namely for $r_{\bar{1}}(\tau,\vec{\sigma}) \neq 0$, $r_{\bar{2}}(\tau,\vec{\sigma}) = 0$;

ii) in figure 2 there is the deformation pattern for the case $C_{\bar{1}} = 0$, $C_{\bar{2}} \neq 0$, namely for $r_{\bar{1}}(\tau,\vec{\sigma}) = 0$, $r_{\bar{2}}(\tau,\vec{\sigma}) \neq 0$.

In the two figures are reported the snapshots at three different times ($t = -1, -0.5, 0$) of the sphere of particles originally at rest (bottom) and the time evolution (from $t = -3$ to $t = 3$) of the six particles at the intersection of the three axes and the sphere of particle (top), whose initial 3-coordinates are $(1,0,0)$ and $(-1,0,0)$ on the x-axis, $(0,1,0)$ and $(0,-1,0)$ on the y-axis, $(0,0,1)$ and $(0,0,-1)$ on the z-axis, respectively. Since a particle on a x axis will remain on the same axes during the evolution, only the x coordinates are reported for the two particles on x axes. The same representation rule has been applied for the particles lying on the other axes.

FIGURES



FIG. 1. Deformation of a sphere of particle at rest induced by the passage of the gravitational wave packet of Eq. (5.15) for $C_{\bar{1}} \neq 0$, $C_{\bar{2}} = 0$.




Particles at t = -1

Particles at t = -0.5

Particles at t = 0.

FIG. 2. Deformation of a sphere of particle at rest induced by the passage of the gravitational wave packet of Eq. (5.15) for $C_{\bar{1}} = 0$, $C_{\bar{2}} \neq 0$.

D. A Relativistic Harmonic Oscillator as a Resonant Detector.

Let us remark that in a TT harmonic gauge the basic idealization of a test resonant detector on the Earth for gravitational waves is a non-relativistic damped harmonic oscillator [22,24]. Given two equal masses m located along the r axis at the positions $r_1(t)$ and $r_2(t)$, connected by a massless spring with spring constant k, damping constant ν and unstretched length l_o, one considers the proper extension $l(t) = \int_{r_1(t)}^{r_2(t)} dt\, \sqrt{1 + h^{TT}_{rr}(t)} \approx [1 + \frac{1}{2}\, h^{TT}_{rr}(t)]\, l_o$ in the metric of the gravitational wave and defines $\xi = l - l_o \approx r_2 - r_1 - l_o + \frac{1}{2}\, h^{TT}_{rr}(r_2 - r_1)$. This leads to modify the equation of motion $\frac{d^2\xi}{dt^2} + 2\gamma\, \frac{d\xi}{dt} + \omega_o^2\, \xi = 0$ ($\omega_o^2 = 2k/m$, $\gamma = \nu/m$), valid in absence of gravitational radiation, to

$$\frac{d^2\xi}{dt^2} + 2\gamma\, \frac{d\xi}{dt} + \omega_o^2\, \xi = \frac{1}{2}\, l_o\, \frac{d^2 h^{TT}_{rr}}{dt^2}. \tag{6.16}$$

This same equation can be obtained as a consequence of the tidal force at the lowest order from the geodesic deviation equation [24], by considering the center of mass of the detector as moving along a geodesic and by identifying ξ with the spatial part of the connecting vector $\triangle x^A(s)$ with $\triangle x^\tau(s) = 0$ (ξ is twice the connecting vector from the center of mass to one of the masses). Note that due to the additional force terms the nearby world-line is no more a geodesic.

Let us try to generalize this resonant detector to *special relativity* and to see the effect on it of our post-Minkowskian gravitational wave. As shown in Ref. [36], there is a description of the relativistic 2-body problem on arbitrary space-like hyper-surfaces Σ_τ, leaves of a 3+1 splitting of Minkowski space-time. If $z^\mu(\tau, \vec{\sigma})$ is the embedding of Σ_τ into Minkowski space-time, the two positive energy scalar particles are described by 3-coordinates $\eta^r_i(\tau)$, $i = 1, 2$, on Σ_τ, such that $x^\mu_i(\tau) = z^\mu(\tau, \vec{\eta}_i(\tau))$ and by the conjugate momenta $\kappa_{ir}(\tau)$. When the hyper-surfaces are the Wigner hyper-planes, orthogonal to the total 4-momentum of the 2-body system, we know how to replace the canonical coordinates $\vec{\eta}_i$, $\vec{\kappa}_i$ with: i) the canonical internal relativistic center of mass $\vec{q}_+$ and its conjugate momentum $\vec{\kappa}_+ \approx 0$ (it is vanishing since we are in the rest-frame instant form of dynamics); ii) relative coordinates $\vec{\rho}$ and $\vec{\pi}$. Even if we do not know explicitly how to express the old basis $\vec{\eta}_i$,

$\vec{\kappa}_i$ in terms of the new basis due to the non-linearity of the canonical transformation, we are able to evaluate the Hamiltonian for the relative motion in the rest frame. In the free case it is $H = \sqrt{m_1^2 + 2\vec{\pi}^2} + \sqrt{m_2^2 + 2\vec{\pi}^2}$. We can add action-at-a-distance interactions either inside the square roots (DrozVincent - Komar - Todorov (DVKT) models [38]), $H = \sqrt{m_1^2 + V(\vec{\rho}^2) + 2\vec{\pi}^2} + \sqrt{m_2^2 + V(\vec{\rho}^2) + 2\vec{\pi}^2}$, or outside them (like the Coulomb potential for charged particles [39]), $H = \sqrt{m_1^2 + 2\vec{\pi}^2} + \sqrt{m_2^2 + 2\vec{\pi}^2} + U(\vec{\rho}^2)$. Due to the square roots we cannot find explicitly the Lagrangian and the Euler-Lagrange equations for the relative motion in the rest frame, except in the special equal mass case. For $m_1 = m_2 = m$ we get [36] $L = -m\sqrt{4 - \dot{\vec{\rho}}^2}$ and an interaction of the DVKT type is introduced by the replacement $m \mapsto M = \sqrt{m^2 + V(\vec{\rho}^2)}$. The Euler-Lagrange equations are

$$\frac{d}{d\tau}\,\frac{M\,\dot{\vec{\rho}}}{\sqrt{4 - \dot{\vec{\rho}}^2}} \overset{\circ}{=} \frac{\sqrt{4 - \dot{\vec{\rho}}^2}}{M}\,V'(\vec{\rho}^2)\,\vec{\rho}, \qquad V'(\vec{\rho}^2) = \frac{dV(\vec{\rho}^2)}{d\,\vec{\rho}^2},$$

or

$$\ddot{\vec{\rho}} \overset{\circ}{=} \frac{4 - \dot{\vec{\rho}}^2}{m^2 + V(\vec{\rho}^2)}\,V'(\vec{\rho}^2)\,[\vec{\rho} - \frac{1}{2}\,\vec{\rho}\cdot\dot{\vec{\rho}}\,\dot{\vec{\rho}}]. \tag{6.17}$$

For $V = -\frac{m}{2}k\,\vec{\rho}^2$ (relativistic harmonic oscillator), the non-relativistic limit of these equations becomes $\ddot{\vec{\rho}} + \omega_o^2\,\vec{\rho} = 0$. A damping term $-\nu\,\dot{\vec{\rho}}$ may be added by hand.

Like in the non-relativistic case, we can adapt this relativistic model, in the case of a test 2-body system, to curved space-time: i) the relativistic center of mass moves along a geodesic Γ, with 4-coordinates $\sigma^A(s)$; ii) the relative variable $\vec{\rho}$ is the connecting vector in Σ_τ to a nearby (non-geodesic) world-line Γ_1. However, since Σ_τ is a Riemannian 3-manifold, the flat vector $\vec{\rho}$ has to be reinterpreted as the field of tangent vectors to the 3-geodesic joining Γ to Γ_1 on each Σ_τ [40]: if $\theta^r(s,\zeta)$ is a 3-geodesic with affine parameter ζ [21] such that Γ is $\theta^r(s,0)$ and Γ_1 is $\theta^r(s,1)$, then we replace $V(\vec{\rho}^2(\tau))$ with $V(\mathcal{R}^2(s))$, where $\mathcal{R}^2(s) = \int_0^1 d\zeta\,\frac{\partial\theta^r(s,\zeta)}{\partial\zeta}\,{}^3g_{rs}(\tau,\theta(s,\zeta))\,\frac{\partial\theta^s(s,\zeta)}{\partial\zeta}$ (it is independent from ζ because the tangent vector $\frac{\partial\theta^r(s,\zeta)}{\partial\zeta}$ is parallel transported along the 3-geodesic). Analogously we make the replacements $\vec{\rho}\cdot\dot{\vec{\rho}} \mapsto \mathcal{R}_1(s) = \int_0^1 d\zeta\,\frac{\partial\theta^r(s,\zeta)}{\partial\zeta}\,{}^3g_{rs}(\tau,\theta(s,\zeta))\,\frac{\partial^2\theta^s(s,\zeta)}{\partial s\,\partial\zeta}$ and $\dot{\vec{\rho}}^2 \mapsto \mathcal{R}_2(s) = \int_0^1 d\zeta\,\frac{\partial^2\theta^r(s,\zeta)}{\partial s\,\partial\zeta}\,{}^3g_{rs}(\tau,\theta(s,\zeta))\,\frac{\partial^2\theta^s(s,\zeta)}{\partial s\,\partial\zeta}$.

By identifying $\rho^r(\tau) \mapsto \triangle\sigma^r(s)$, $\dot{\rho}^r(\tau) \mapsto \frac{d\triangle\sigma^r(s)}{ds}$, $\ddot{\vec{\rho}}^r \mapsto \frac{d^2\triangle\sigma^r(s)}{ds^2}$, and by adding the force terms (damping included) of Eqs.(6.17) in the geodesic deviation equation (6.9) we get

$$\frac{d^2\triangle x^r(s)}{ds^2} + \Big[2\,{}^4\hat{\Gamma}^r_{\tau s}(s) + \frac{4 - \mathcal{R}_2(s)}{m^2 + V(\mathcal{R}^2(s))}\,V'(\mathcal{R}^2(s))\,(\nu + \frac{1}{2}\,\mathcal{R}_1(s))\Big]\,\frac{d\triangle x^s(s)}{ds} +$$
$$+ \Big[\frac{d\,{}^4\hat{\Gamma}^r_{\tau s}(s)}{ds} + {}^4\hat{R}^r{}_{\tau s\tau}(s) - \frac{4 - \mathcal{R}_2(s)}{m^2 + V(\mathcal{R}^2(s))}\,V'(\mathcal{R}^2(s))\Big]\,\triangle x^s(s) = 0. \tag{6.18}$$

---

[21] $\frac{\partial^2\theta^r(s,\zeta)}{\partial\zeta^2} + {}^3\hat{\Gamma}^r_{uv}(\theta(s,\zeta))\,\frac{\partial\theta^u(s,\zeta)}{\partial\zeta}\,\frac{\partial\theta^v(s,\zeta)}{\partial\zeta} = 0$ with the 3-Christoffel symbol of Eqs.(D1).

This is the post-Minkowskian counterpart in our coordinates of the non-relativistic equation (6.16) of the TT harmonic gauge.

For very small $|\vec{\rho}|$ we can make the approximations $\mathcal{R}^2 \approx \frac{\partial \theta^r(s,\zeta)}{\partial \zeta}|_{\zeta=0}\, {}^3g_{rs}(\sigma^A(s))\, \frac{\partial \theta^s(s,\zeta)}{\partial \zeta}|_{\zeta=0}$, $\mathcal{R}_1 \approx \frac{\partial \theta^r(s,\zeta)}{\partial \zeta}|_{\zeta=0}\, {}^3g_{rs}(\sigma^A(s))\, \frac{\partial^2 \theta^s(s,\zeta)}{\partial s\, \partial \zeta}|_{\zeta=0}$, $\mathcal{R}_2 \approx \frac{\partial^2 \theta^r(s,\zeta)}{\partial s\, \partial \zeta}|_{\zeta=0}\, {}^3g_{rs}(\sigma^A(s))\, \frac{\partial^2 \theta^s(s,\zeta)}{\partial s\, \partial \zeta}|_{\zeta=0}$: this is a pole-dipole approximation of the test 2-body problem.

VII. THE EMBEDDING OF THE WSW HYPER-SURFACES, THE ASSOCIATED CONGRUENCE OF TIME-LIKE OBSERVERS AND THE NULL GEODESICS.

A. The Post-Minkowskian WSW Hyper-surfaces.

The WSW triads ${}^3e^{(WSW)\,r}_{(a)}(\tau,\vec{\sigma}) \rightarrow_{|\vec{\sigma}|\rightarrow\infty} \delta^r_{(a)}$, parallel transported from spatial infinity on the linearized WSW CMC hyper-surface, and the associated cotriads ${}^3e^{(WSW)}_{(a)r}(\tau,\vec{\sigma})$ [22] are the solution of Eqs.(A18), whose linearized form is

$$
\begin{aligned}
&\partial_r\Big(\sqrt{\hat{\gamma}}\,{}^3e^{(WSW)r}_{(a)}\Big)(\tau,\vec{\sigma}) = \partial_r\,{}^3e^{(WSW)r}_{(a)}(\tau,\vec{\sigma}) + 3\delta^r_{(a)}\partial_r q(\tau,\vec{\sigma}) + O(r^2_{\bar{a}}) = 0,\\
&\partial_1\,{}^3e^{(WSW)}_{(2)3}(\tau,\vec{\sigma}) + \partial_3\,{}^3e^{(WSW)}_{(1)2}(\tau,\vec{\sigma}) + \partial_2\,{}^3e^{(WSW)}_{(3)1}(\tau,\vec{\sigma}) =\\
&= \Big[{}^3\hat{\Gamma}^2_{13} + {}^3\hat{\Gamma}^1_{32} + {}^3\hat{\Gamma}^3_{21}\Big](\tau,\vec{\sigma}) + O(r^2_{\bar{a}}) = 0 + O(r^2_{\bar{a}}),
\end{aligned}
\tag{7.1}
$$

where Eqs.(D1) have been used.

These equations imply

$$
\begin{aligned}
{}^3e^{(WSW)\,r}_{(a)}(\tau,\vec{\sigma}) &= f^r_{(a)\perp}(\tau,\vec{\sigma}) + \frac{\sqrt{3}}{2}\,\delta^r_{(a)}\sum_{\bar{a}u}\gamma_{\bar{a}u}\int d^3\sigma_1\,\frac{\partial^2_{1u}\,r_{\bar{a}}(\tau,\vec{\sigma}_1)}{4\pi\,|\vec{\sigma}-\vec{\sigma}_1|},\\
&\partial_r\,f^r_{(a)\perp}(\tau,\vec{\sigma}) = 0, \qquad f^r_{(a)\perp}(\tau,\vec{\sigma}) \rightarrow_{|\vec{\sigma}|\rightarrow\infty} \delta^r_{(a)},\\
&\partial_1\,f^3_{(2)\perp} + \partial_3\,f^2_{(1)\perp} + \partial_2\,f^1_{(3)\perp} = 0 + O(r^2_{\bar{a}}),\\
&\Rightarrow\quad f^r_{(a)\perp} = \delta^r_{(a)} + (\delta^{rs} + \frac{\partial^r\,\partial^s}{\triangle})\,g_{(a)s},
\end{aligned}
\tag{7.2}
$$

with $g_{(a)s}$ arbitrary. The simplest solution is to take $g_{(a)s} = 0$, so that the WSW triads are

$$
{}^3e^{(WSW)r}_{(a)} = \delta^r_{(a)}\,\Big[1 + \frac{\sqrt{3}}{2}\sum_{\bar{a}u}\gamma_{\bar{a}u}\int d^3\sigma_1\,\frac{\partial^2_{1u}\,r_{\bar{a}}(\tau,\vec{\sigma}_1)}{4\pi\,|\vec{\sigma}-\vec{\sigma}_1|}\Big].
\tag{7.3}
$$

Eqs.(A19), allow to find the associated WSW adapted tetrads (preferred ADM Eulerian observers or *asymptotic fixed stars* giving a *local compass of inertia* to be compared with local (Fermi-Walker transported or not) gyroscopes)

$$
\begin{aligned}
{}^4_{(\Sigma)}\check{\tilde{E}}^{(WSW)A}_{(o)}(\tau,\vec{\sigma}) &= -\epsilon\Big(1;\,-n_r(\tau,\vec{\sigma})\Big),\\
{}^4_{(\Sigma)}\check{\tilde{E}}^{(WSW)A}_{(a)}(\tau,\vec{\sigma}) &= \Big(0;\,{}^3e^{(WSW)\,r}_{(a)}(\tau,\vec{\sigma})\Big).
\end{aligned}
\tag{7.4}
$$

Once the WSW triads are known, Eqs.(A20) and (A21) give the embedding of the linearized WSW CMC hyper-surfaces into the linearized space-time

---

[22] ${}^3e^{(WSW)\,r}_{(a)} = {}^3\hat{g}^{rs}\,{}^3e^{(WSW)}_{(a)s} = -\epsilon\delta^{rs}\,{}^3e^{(WSW)}_{(a)s} + 2\epsilon\delta^r_{(a)}[q + \frac{2}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}r}r_{\bar{a}}] + O(r^2_{\bar{a}})$.

$$z^\mu_{(WSW)}(\tau,\vec{\sigma}) = x^\mu_{(\infty)}(0) + b^\mu_A(\tau,\vec{\sigma}) F^A(\tau,\vec{\sigma}),$$

$$\begin{aligned} &F^\tau(\tau,\vec{\sigma}) = -\epsilon\tau,\\ &F^s(\tau,\vec{\sigma}) = {}^3e^{(WSW)\,s}_{(a)}(\tau,\vec{\sigma})\delta_{(a)r}\sigma^r + \epsilon\tau n_s(\tau,\vec{\sigma}), \end{aligned} \tag{7.5}$$

with the transition coefficients $b^\mu_A = \frac{\partial z^\mu_{(WSW)}}{\partial\sigma^A}$ solution of the linear partial differential equations

$$F^B(\tau,\vec{\sigma}) \frac{\partial b^\mu_B(\tau,\vec{\sigma})}{\partial\sigma^A} = \Big(\delta^B_A - \frac{\partial F^B(\tau,\vec{\sigma})}{\partial\sigma^A}\Big) b^\mu_B(\tau,\vec{\sigma}). \tag{7.6}$$

In adapted coordinates Eqs.(7.5) of the embedding of linearized WSW CMC hyper-surfaces Σ_τ become

$$z^A_{(WSW)}(\tau,\vec{\sigma}) = x^A_{(\infty)}(0) + F^A(\tau,\vec{\sigma}), \tag{7.7}$$

with $z^\tau_{(WSW)}(\tau,\vec{\sigma}) = x^\tau_{(\infty)}(0) - \epsilon\,\tau$. The time-like evolution vector is $z^\mu_{(WSW)\tau}(\tau,\vec{\sigma}) = \Big(-\epsilon; [\epsilon\, n_r + \epsilon\,\tau\,\partial_\tau\, n_r + \partial_\tau\, {}^3e^{(WSW)r}_{(a)}\,\sigma^a](\tau,\vec{\sigma})\Big)$.

B. Congruences of Timelike Observers.

In Eqs.(3.9) we have the expression of the $\Sigma^{(WSW)}_\tau$-adapted tetrads and cotetrads. In particular there is the expression for the contra- and co-variant normals, $l^A(\tau,\vec{\sigma})$ and $l_A(\tau,\vec{\sigma})$, to $\Sigma^{(WSW)}_\tau$. The associated 4-velocity field defines a (non-rotating, surface forming) congruence of time-like observers orthogonal to $\Sigma^{(WSW)}_\tau$, using the parameter τ, labeling the leaves of the foliation, as evolution parameter [23]. Let us remark that τ *is not in general the proper time of any observer of the congruence.*

As for any congruence, we have the decomposition ($P_{AB} = {}^4g_{AB} - l_A\, l_B$)

$$\begin{aligned} {}^4\nabla_A\, l_B &= l_A\, a_B + \frac{1}{3}\,\Theta\, P_{AB} + \sigma_{AB} + \omega_{AB},\\ a^A &= l^B\, {}^4\nabla_B\, l^A, \qquad 4-acceleration, \end{aligned}$$

---

[23] This is the *hyper-surface point of view* according to Ref. [41]. Instead, the *threading point of view* is a description involving only a rotating congruence of observers (like the one which can be built with the 4-velocity field associated to $z^A_{(WSW)\tau}(\tau,\vec{\sigma})$) : since this congruence, being rotating, is not surface-forming (non-zero vorticity), in each point we can only divide the tangent space in the direction parallel to the 4-velocity and the orthogonal complement (the local rest frame). On the other hand, the *slicing point of view*, originally adopted in ADM canonical gravity, uses two congruences: the non-rotating one with the normals to Σ_τ as 4-velocity fields and a second (rotating, non-surface-forming) congruence of observers, whose 4-velocity field is the field of time-like unit vectors determined by the τ derivative of the embeddings identifying the leaves Σ_τ (the so-called evolution vector field). Furthermore, it uses the affine parameter describing the world-lines of this second family of observers as Hamiltonian evolution parameter.

$$\begin{aligned}
&\Theta = {}^4\nabla_A\, l^A, \quad \mathit{scalar\,(volume)\,rate\,of\,expansion},\\
&\sigma_{AB} = \frac{1}{2}\,(a_A\, l_B + a_B\, l_A) + \frac{1}{2}\,({}^4\nabla_A\, l_B + {}^4\nabla_B\, l_A) - \frac{1}{3}\,\Theta\, P_{AB},\\
&\mathit{rate\,of\,shear\,tensor\,(with\,magnitude}\, \sigma^2 = \frac{1}{2}\sigma_{AB}\sigma^{AB}),\\
&\omega_{AB} = -\omega_{BA} = \epsilon_{ABCD}\, \omega^C\, l^D = \frac{1}{2}\,(a_A\, l_B - a_B\, l_A) + \frac{1}{2}\,({}^4\nabla_A\, l_B - {}^4\nabla_B\, l_A) = 0,\\
&\mathit{twist\,or\,vorticity\,tensor}, \qquad \omega^A = \frac{1}{2}\,\epsilon^{ABCD}\,\omega_{BC}\, l_D = 0, \quad \mathit{vorticity\,vector}. \qquad (7.8)
\end{aligned}$$

Θ is the *expansion* (it measures the average expansion of the infinitesimally nearby world-lines surrounding a given world-line in the congruence), σ_{AB} the *shear* (it measures how an initial sphere in the tangent space to the given world-line, which is Lie transported along l^A, will distort toward an ellipsoid with principal axes given by the eigenvectors of $\sigma^A{}_B$ with rate given by the eigenvalues of $\sigma^A{}_B$) and ω_{AB} the *twist or vorticity* (it measures the rotation of the infinitesimally nearby world-lines surrounding the given one); σ_{AB} and ω_{AB} are purely spatial ($\sigma_{AB}l^B = \omega_{AB}l^B = 0$). Due to the Frobenius theorem, the congruence is (locally) hyper-surface orthogonal if and only if $\omega_{AB} = 0$. The equation $\frac{1}{l}\, l^A\, \partial_A\, l = \frac{1}{3}\,\Theta$ defines a representative length l along the world-line of l^A, describing the volume expansion (contraction) behaviour of the congruence completely.

The linearized acceleration of the observers vanishes (i.e. at the lowest level we get inertial observers) so that there is no gravito-electric force on test particles in the sense of Ref. [41] (geodesic in local rest frame)

$$a^A = l^B\, \partial_B\, l^A + {}^4\Gamma^A_{BC}\, l^B l^C = (0; -\partial_\tau\, n_r + {}^4\Gamma^r_{\tau\tau}) + O(r^2_{\bar a}) = 0 + O(r^2_{\bar a}). \qquad (7.9)$$

By using Eqs.(D5) we get ${}^4\nabla_A\, l_B = -{}^4\Gamma^C_{AB}\, l_C = \epsilon\, {}^4\Gamma^\tau_{AB} + O(r^2_{\bar a})$. The expansion of the congruence is

$$\Theta = {}^4\nabla_A\, l^A = \partial_A\, l^A + {}^4\Gamma^A_{AB}\, l^B = \epsilon\vec{\partial}\cdot\vec{n} - \epsilon\,\vec{\partial}\cdot\vec{n} + O(r^2_{\bar a}) = 0 + O(r^2_{\bar a}). \qquad (7.10)$$

The shear is not zero so that there is gravito-magnetism on test particles in the sense of Ref. [41] (geodesic in local rest frame)

$$\sigma_{AB} = \frac{1}{2}\,(a_A\, l_B + a_B\, l_A) + \frac{1}{2}\,({}^4\nabla_A\, l_B + {}^4\nabla_B\, l_A) = \epsilon\, {}^4\Gamma^\tau_{AB} + O(r^2_{\bar a}), \qquad (7.11)$$

while the vorticity vanishes (the congruence is surface forming)

$$\omega_{AB} = \frac{1}{2}\,(a_A\, l_B - a_B\, l_A) + \frac{1}{2}\,({}^4\nabla_A\, l_B - {}^4\nabla_B\, l_A) = 0 + O(r^2_{\bar a}). \qquad (7.12)$$

If $x^A_{\vec{\sigma}_o}(\tau)$ is the time-like world-line, in adapted coordinates, of the observer crossing the leave Σ_{τ_o} at $\vec{\sigma}_o$, we have

$$
\begin{aligned}
x^A_{\vec{\sigma}_o}(\tau) &= \Big(\tau; \vec{\rho}_{\vec{\sigma}_o}(\tau)\Big) = \\
&= \Big(\tau_o - \epsilon\, [z^\tau_{(WSW)}(\tau, \vec{\rho}_{\vec{\sigma}_o}(\tau)\,) - z^\tau_{(WSW)}(\tau_o, \vec{\sigma}_o)]; \\
&\quad \sigma^r_o + z^r_{(WSW)}(\tau, \vec{\rho}_{\vec{\sigma}_o}(\tau)\,) - z^r_{(WSW)}(\tau_o, \vec{\sigma}_o)\Big), \\
x^A_{\vec{\sigma}_o}(\tau_o) &= \Big(\tau_o; \vec{\sigma}_o\Big), \qquad \vec{\rho}_{\vec{\sigma}_o}(\tau_o) = \vec{\sigma}_o.
\end{aligned}
\tag{7.13}
$$

The effective trajectory $\vec{\rho}_{\vec{\sigma}_o}(\tau)$ is determined by solving the equations $\vec{\rho}_{\vec{\sigma}_o}(\tau) = \vec{\sigma}_o + \vec{z}_{(WSW)}(\tau, \vec{\rho}_{\vec{\sigma}_o}(\tau)\,) - \vec{z}_{(WSW)}(\tau_o, \vec{\sigma}_o)$ with $\vec{z}_{(WSW)}$ given by Eq.(7.7). The 4-velocity $\dot{x}^A_{\vec{\sigma}_o}(\tau) = \frac{dx^A_{\vec{\sigma}_o}(\tau)}{d\tau}$ satisfies

$$
l^A_{\vec{\sigma}_o}(\tau) = l^A(\tau, \vec{\rho}_{\vec{\sigma}_o}(\tau)) = \frac{\dot{x}^A_{\vec{\sigma}_o}(\tau)}{\sqrt{{}^4g_{BC}(x_{\vec{\sigma}_o}(\tau))\, \dot{x}^B_{\vec{\sigma}_o}(\tau)\, \dot{x}^C_{\vec{\sigma}_o}(\tau)}},
\tag{7.14}
$$

with the observer acceleration given by $a^A_{\vec{\sigma}_o}(\tau) = \frac{dl^A_{\vec{\sigma}_o}(\tau)}{d\tau} = 0 + O(r^2_{\bar{a}})$, $a^A_{\vec{\sigma}_o}(\tau)\, l_{\vec{\sigma}_o\, A}(\tau) = 0$.

Since $\dot{x}^\tau_{\vec{\sigma}_o}(\tau) = \tau$, $\dot{x}^r_{\vec{\sigma}_o}(\tau) = O(r_{\bar{a}})$, we get ${}^4g_{AB}\, \dot{x}^A_{\vec{\sigma}_o}\, \dot{x}^B_{\vec{\sigma}_o} = \epsilon + O(r^2_{\bar{a}})$, namely that to this order τ is the proper time of any observer of the congruence.

Yet, the ADM canonical formalism gives us an additional information. Actually, on each space-like hyper-surface $\Sigma^{(WSW)}_\tau$ of the foliation, there is a privileged contravariant space-like direction identified by the lapse and shift gauge variables

$$
\begin{aligned}
\mathcal{N}^A(\tau, \vec{\sigma}) &= \frac{1}{|\vec{N}(\tau, \vec{\sigma})|} \Big(0; n^r(\tau, \vec{\sigma})\Big) = \Big(0; n_r(\tau, \vec{\sigma})\Big) + O(r^2_{\bar{a}}), \\
\mathcal{N}_A(\tau, \vec{\sigma}) &= |\vec{N}(\tau, \vec{\sigma})| \Big(1; \frac{N_r(\tau, \vec{\sigma})}{|\vec{N}(\tau, \vec{\sigma})|^2}\Big) = \Big(|\vec{n}(\tau, \vec{\sigma})|; n_r(\tau, \vec{\sigma})\Big) + O(r^2_{\bar{a}}), \\
&\mathcal{N}^A(\tau, \vec{\sigma})\, l_A(\tau, \vec{\sigma}) = 0, \qquad \mathcal{N}^\mu(\tau, \vec{\sigma})\, \mathcal{N}_\mu(\tau, \vec{\sigma}) = -\epsilon, \\
&|\vec{N}(\tau, \vec{\sigma})| = \sqrt{({}^3g_{rs}\, N^r\, N^s)(\tau, \vec{\sigma})} = |\vec{n}(\tau, \vec{\sigma})| + O(r^2_{\bar{a}}).
\end{aligned}
\tag{7.15}
$$

If 4-coordinates exist, corresponding to an on-shell complete Hamiltonian gauge fixing, such that the vector field identified by $\mathcal{N}^A(\tau, \vec{\sigma})$ on each $\Sigma^{(WSW)}_\tau$ is surface-forming (zero vorticity [24]), then each Σ_τ can be foliated with 2-surfaces, and the 3+1 splitting of space-time becomes a (2+1)+1 splitting corresponding to the 2+2 splittings studied by Stachel and d'Inverno [42].

We have therefore a natural candidate for one of the three spatial triads of each observer: $E^A_{\vec{\sigma}_o\,(\mathcal{N})}(\tau) = \mathcal{N}^A_{\vec{\sigma}_o}(\tau) = \mathcal{N}^A(\tau, \vec{\rho}_{\vec{\sigma}_o}(\tau))$. By means of $l^A_{\vec{\sigma}_o}(\tau) = l^A(\tau, \vec{\rho}_{\vec{\sigma}_o}(\tau))$ and $\mathcal{N}^A_{\vec{\sigma}_o}(\tau)$, we can construct two null vectors at each space-time point

---

[24] This requires that $\mathcal{N}_A\, d\sigma^A$ is a closed 1-form, namely that we have $\partial_\tau\, n_r = \partial_r\, |\vec{n}| + O(r^2_{\bar{a}})$ and $\partial_r\, n_s = \partial_s\, n_r + O(r^2_{\bar{a}})$. In turn, this requires $n_r = \partial_r\, f + O(r^2_{\bar{a}})$ with $\partial_\tau\, f = |\vec{n}| + const$. Our gauge has a non-surface forming $\mathcal{N}^A$.

$$\mathcal{K}^A_{\vec{\sigma}_o}(\tau) = \sqrt{\frac{|\vec{n}|}{2}}\Big(l^A_{\vec{\sigma}_o}(\tau) + \mathcal{N}^A_{\vec{\sigma}_o}(\tau)\Big) = \sqrt{\frac{1}{2}}\Big(l^A_{\vec{\sigma}_o}(\tau) + \mathcal{N}^A_{\vec{\sigma}_o}(\tau)\Big) + O(r^2_{\bar{a}}),$$
$$\mathcal{L}^A_{\vec{\sigma}_o}(\tau) = \frac{1}{\sqrt{2\,|\vec{n}|}}\Big(l^A_{\vec{\sigma}_o}(\tau) - \mathcal{N}^A_{\vec{\sigma}_o}(\tau)\Big) = \frac{1}{\sqrt{2}}\Big(l^A_{\vec{\sigma}_o}(\tau) - \mathcal{N}^A_{\vec{\sigma}_o}(\tau)\Big) + O(r^2_{\bar{a}}). \qquad (7.16)$$

and then we can arrive at a *null tetrad* of the type used in the Newman-Penrose formalism [43]. The last two axes of the spatial triad can be chosen as two space-like circular complex polarization vectors $E^\mu_{\vec{\sigma}_o\,(\pm)}(\tau)$, like in electromagnetism. They are built starting from the transverse helicity polarization vectors $E^\mu_{\vec{\sigma}_o\,(1,2)}(\tau)$, which are the first and second columns of the standard Wigner helicity boost generating $\mathcal{K}^\mu_{\vec{\sigma}_o}(\tau)$ from the reference vector $\overset{\circ}{\mathcal{K}}{}^\mu_{\vec{\sigma}_o}(\tau) = |\vec{N}|\,\Big(1; 001\Big)$.

Let us call $E^{(ADM)\mu}_{\vec{\sigma}_o\,(\alpha)}(\tau)$ the ADM tetrad formed by $l^\mu_{\vec{\sigma}_o}(\tau)$, $\mathcal{N}^\mu_{\vec{\sigma}_o}(\tau)$, $E^\mu_{\vec{\sigma}_o\,(1,2)}(\tau)$. This tetrad will not be in general Fermi-Walker transported along the world-line $x^\mu_{\vec{\sigma}_o}(\tau)$ of the observer.

Another possible (but only on-shell) choice of the spatial triad together with the unit normal to $\Sigma^{(WSW)}_\tau$ is the *local WSW (on-shell) compass of inertia* defined in Eqs.(7.4). This local compass corresponds to the *standard of non rotation with respect to the fixed stars* and its τ-evolution, dictated by Einstein's equations, does not correspond to the FW transport, which is defined independently from them using only local geometrical and group-theoretical concepts. The WSW local compass of inertia corresponds to pointing to the fixed stars with a telescope and is needed in a satellite like Gravity Probe B to detect the frame dragging (or gravito-magnetic Lense-Thirring effect) of the inertial frames by means of the rotation with respect to it of a FW transported gyroscope.

Given the 4-velocity $l^A_{\vec{\sigma}_o}(\tau) = E^A_{\vec{\sigma}_o}(\tau)$ of the observer, the observer spatial triads $E^A_{\vec{\sigma}_o\,(a)}(\tau)$, $a = 1, 2, 3$, have to be chosen in a conventional way, namely by means of a conventional assignment of an origin for the local measurements of rotations. Usually, the choice corresponds to Fermi-Walker (FW) transported (*gyroscope-type transport, non-rotating observer*) tetrads $E^{(FW)\,A}_{\vec{\sigma}_o\,(\alpha)}(\tau)$, such that (we show also the implication of the linearization)

$$\frac{D}{D\tau}\,E^{(FW)\,A}_{\vec{\sigma}_o\,(a)}(\tau) = \Omega^{(FW)A}_{\vec{\sigma}_o}{}_B(\tau)\,E^{(FW)\,B}_{\vec{\sigma}_o\,(a)}(\tau) = l^A_{\vec{\sigma}_o}(\tau)\,a_{\vec{\sigma}_o\,B}(\tau)\,E^{(FW)\,B}_{\vec{\sigma}_o\,(a)}(\tau) = 0 + O(r^2_{\bar{a}}),$$
$$\Omega^{(FW)AB}_{\vec{\sigma}_o}(\tau) = a^A_{\vec{\sigma}_o}(\tau)\,l^B_{\vec{\sigma}_o}(\tau) - a^B_{\vec{\sigma}_o}(\tau)\,l^A_{\vec{\sigma}_o}(\tau). \qquad (7.17)$$

The triad $E^{(FW)\,A}_{\vec{\sigma}_o\,(a)}(\tau)$ is the correct relativistic generalization of global Galilean non-rotating frames. Naturally any other choice of the triads (Lie transport, co-rotating-FW transport,...) is possible. A generic triad $E^A_{\vec{\sigma}_o\,(a)}(\tau)$ will satisfy $\frac{D}{D\tau}\,E^A_{\vec{\sigma}_o\,(a)}(\tau) = \Omega_{\vec{\sigma}_o}{}^A{}_B(\tau)\,E^B_{\vec{\sigma}_o\,(a)}(\tau)$ with $\Omega^{AB}_{\vec{\sigma}_o} = \Omega^{(FW)\,AB}_{\vec{\sigma}_o} + \Omega^{(SR)\,AB}_{\vec{\sigma}_o}$ with the spatial rotation part $\Omega^{(SR)\,AB}_{\vec{\sigma}_o} = \epsilon^{ABCD}\,l_{\vec{\sigma}_o\,C}\,J_{\vec{\sigma}_o\,D}$, $J^A_{\vec{\sigma}_o}\,l_{\vec{\sigma}_o\,A} = 0$, producing a rotation of the gyroscope in the local space-like 2-plane orthogonal to $l^A_{\vec{\sigma}_o}$ and $J^A_{\vec{\sigma}_o}$.

In the linearized theory on the WSW hyper-surfaces of our gauge FW transport implies no τ-dependence for the triads $E^{(FW)\,A}_{\vec{\sigma}_o\,(a)}(\tau)$. Neither the ADM tetrads nor the WSW tetrads (7.4) are FW transported.

See Ref. [41] for the description of a geodesics $y^\mu(\tau)$, the world-line of a scalar test particle, from the point of view of those observers $\gamma_{\vec{\sigma}_o, y(\tau)}$ in the congruence which intersect it, namely such that at τ it holds $x^{\mu}_{\vec{\sigma}_o, y(\tau)}(\tau) = y^{\mu}(\tau)$. The family of these observers is called a *relative observer world 2-sheet* in Ref. [41].

C. The Null Geodesics, the Deformed Light-Cone and the Eikonal.

The solution of Eqs.(6.1) for *null geodesics* has still the form

$$\begin{aligned}
\tau(s) &= a^{\tau} + [b^{\tau}_{(o)} + b^{\tau}_{(1)}]\, s - \int_0^s ds_1 \int_0^{s_1} ds_2 {}^4\hat{\Gamma}^{\tau}_{ru}(a + b_{(o)} s_2) b^r b^u \\
\sigma^u(s) &= a^u + b^u - \int_0^s ds_1 \int_0^{s_1} ds_2 \Big[(b^{\tau}_{(o)})^2 - 2\, {}^4\hat{\Gamma}^u_{\tau r}(a + b_{(o)} s_2) b^{\tau}_{(o)}\, b^r - {}^3\hat{\Gamma}^u_{rt}(a + b_{(o)} s_2) b^r b^t \Big]
\end{aligned}$$

$$with$$

$$\begin{aligned}
p^{\tau}(s) &= \frac{d\tau(s)}{ds} = b^{\tau}_{(o)} + b^{\tau}_{(1)} - \int_0^s ds_2 {}^4\hat{\Gamma}^{\tau}_{ru}(a + b_{(o)} s_2)\, b^r b^u, \\
p^u(s) &= \frac{d\sigma^u(s)}{ds} = b^u - \int_0^s ds_2 \Big[(b^{\tau}_{(o)})^2 - 2\, {}^4\hat{\Gamma}^u_{\tau r}(a + b_{(o)} s_2) b^{\tau}_{(o)}\, b^r - {}^3\hat{\Gamma}^u_{rt}(a + b_{(o)} s_2) b^r b^t \Big],
\end{aligned} \tag{7.18}$$

but now at $s = 0$ we have

$$\begin{aligned}
{}^4g_{AB} p^A p^B|_{s=0} &= \epsilon \Big[(p^{\tau}(s))^2 - 2 n_r(s, r_{\bar{a}}(s)) p^r(s) p^{\tau}(s) - [\delta_{rt} - \epsilon^4 h_{rt}(s, r_{\bar{a}}(s)) p^r(s) p^t(s)] \Big]|_{s=0} = \\
&= 0,
\end{aligned}$$

$$0 = (b^{\tau})^2 - 2 b^{\tau} b^r n_r(a^{\tau}, r_{\bar{a}}(a^{\tau})) - [\delta_{rt} - \epsilon^4 h_{rt}(a^{\tau}, r_{\bar{a}}(a^{\tau}))]\, b^r b^t, \quad at\, s = 0, \tag{7.19}$$

whose solution is

$$b^{\tau} = b^{\tau}_{(o)} + b^{\tau}_{(1)} == b^r n_r(a^{\tau}, r_{\bar{a}}(a^{\tau})) \pm \sqrt{(b^r n_r(a^{\tau}, r_{\bar{a}}(a^{\tau})))^2 + [\delta_{rt} - \epsilon^4 h_{rt}(a^{\tau}, r_{\bar{a}}(a^{\tau}))]\, b^r b^t},$$

$$\begin{aligned}
b^{\tau}_{(o)} &= \pm \sqrt{\delta_{rs} b^r b^s}, \\
b^{\tau}_{(1)} &= n_r(a^{\tau}, r_{\bar{a}}(a^{\tau}))\, b^r \mp \frac{\epsilon}{2} \frac{{}^4h_{rt}(a^{\tau}, r_{\bar{a}}(a^{\tau})) b^r b^t}{\sqrt{\delta_{rs} b^r b^s}} + O(r^2_{\bar{a}}).
\end{aligned} \tag{7.20}$$

Therefore, if we consider the family of null geodesics emanating from a fixed point of space-time and parametrized by $\{b^r\}$, we obtain the null surface describing the *deformed light-cone* through that point in our post-Minkowskian Einstein space-time.

Let us now consider the *eikonal equation* [7] ${}^4g^{AB}(\tau, \vec{\sigma})\, \partial_A\, U(\tau, \vec{\sigma})\, \partial_B\, U(\tau, \vec{\sigma}) = 0$, whose solution, the so-called *optical function* U, is used to find the null hyper-surfaces $U(\tau, \vec{\sigma}) = 0$ tangent to the deformed light-cones, generalizing the planes $x^{\pm} = \frac{1}{\sqrt{2}}\, (x^o \pm x^3) = 0$ tangent to the light-cone in special relativity, where they are used for the front (or null) form of dynamics in light-cone coordinates.

In Minkowski space-time with Cartesian coordinates the eikonal equation is $(\partial_\tau\, U_M(\tau, \vec{\sigma}))^2 - (\vec{\partial}\, U_M(\tau, \vec{\sigma}))^2 = 0$, namely $\partial_\tau\, U_M = \alpha\, |\vec{\partial}\, U_M|$ with $\alpha = \pm$. The solutions are $U_M(\tau, \vec{\sigma}) = f(\tau \pm \sum_r A_r\, \sigma^r)$ with $\sum_r A_r^2 = 1$.

After the linearization in our gauge the eikonal equation becomes ($\hat{\gamma}^{rs} = \gamma_r\, \delta^{rs}$)

$$(\partial_\tau\, U(\tau, \vec{\sigma}))^2 - 2\, \partial_\tau\, U(\tau, \vec{\sigma})\, \sum_r n_r(\tau, \vec{\sigma})\, \partial_r\, U(\tau, \vec{\sigma}) - \sum_r \gamma_r(\tau, \vec{\sigma})\, (\partial_r\, U(\tau, \vec{\sigma}))^2 = 0. \quad (7.21)$$

Let us put $U = U_M + V$ with $V = O(r_{\bar{a}})$. By disregarding terms of order V^2 the eikonal equation becomes the following quasi-linear partial differential equation for V

$$\begin{aligned} &2\, \Big[\alpha |\vec{\partial}\, U_M| - \sum_r n_r\, \partial_r\, U_M\Big]\, \partial_\tau\, V - 2\, \sum_r\, \Big[\alpha\, |\vec{\partial}\, U_M|\, n_r + \gamma_r\, \partial_r\, U_m\Big]\, \partial_r V = \\ &= \sum_r (\gamma_r - 1)\, (\partial_r\, U_M)^2 + 2\, \alpha\, |\vec{\partial}\, U_M|\, \sum_r n_r\, \partial_r\, U_M. \end{aligned} \quad (7.22)$$

If we write this equation in the form $a\, \partial_\tau\, V + \sum_r a_r\, \partial_r\, V = F$, then a solution $U = U_M + V$ of the linearized eikonal equation can be obtained with the method of characteristics [25] if it is possible to find an explicit solution of the system

$$\frac{dV}{F} = \frac{d\tau}{a} = \frac{d\sigma^r}{a_r}. \quad (7.23)$$

VIII. CONCLUSIONS.

We have defined a background-independent Hamiltonian linearization of vacuum canonical tetrad gravity in a completely fixed 3-orthogonal gauge in the framework of its rest-frame instant form, where the evolution is governed by the weak ADM energy. In this *non-harmonic* gauge every quantity is expressed in terms of the DO's $r_{\bar{a}}(\tau, \vec{\sigma})$, $\pi_{\bar{a}}(\tau, \vec{\sigma})$, $\bar{a} = 1, 2$, parametrizing the independent degrees of freedom of the gravitational field. The method is based on the linearization of the Lichnerowicz equation for the conformal factor of the 3-metric and on the restriction of the weak ADM energy to its part quadratic in the DO's.

As a consequence we succeed for the first time in solving all the constraints of tetrad gravity (super-hamiltonian constraint included) and to find a solution of the linearized Einstein equations in the uniquely defined 4-coordinate system induced by the chosen gauge, which corresponds to a *post-Minkowskian Einstein space-time* of the Christodoulou-Klainermann type. The DO's $r_{\bar{a}}(\tau, \vec{\sigma})$ turn out to satisfy the massless wave equation even if we are not in a harmonic gauge and we get non-zero shift functions (namely our 4-coordinates are non-synchronous). Besides re-opening the Hamiltonian approach to gravity, we can show explicitly the role played by the deterministically predictable DO's of the gravitational field in deforming the structures of the flat Minkowski space-time without having used it as a background for the propagation of a massless spin 2 field as it happens in the standard treatment of gravitational waves in the harmonic gauges. In particular the two configuration DO's $r_{\bar{a}}(\tau, \vec{\sigma})$, $\bar{a} = 1, 2$, of our gauge, where the 3-metric is diagonal, replace the two polarizations of the TT harmonic gauge (with its non-diagonal 3-metric) and, through the geodesic deviation equation, they induce two well defined patterns of deformation on a sphere of test particles. Even if we were able to eliminate the background with our Hamiltonian linearization, our results are still coordinate (i.e. gauge)-dependent like in all existing treatments of gravitational waves: indeed our DO's $r_{\bar{a}}$, $\pi_{\bar{a}}$ are coordinate-dependent non-tensorial quantities. To get a coordinate-independent description of them, we have to verify the main conjecture of Ref. [15]. It states that by means of a Hamiltonian re-formulation of the Newman-Penrose formalism [43] it should be possible to find a Shanmugadhsan canonical basis in which the DO's are also (coordinate-independent) Bergmann observables and also the gauge variables are coordinate-independent.

We have also made some comments on the coordinate dependence of the gravito-magnetic effects: for instance in our non-harmonic gauge the gravito-electric-magnetic analogy does not hold. Again the verification of the main conjecture of Ref. [15] would allow a coordinate-independent description of gravito-magnetism.

These results about background-independent gravitational waves in post-Minkowskian space-times are welcome because *they open the possibility, after the introduction of matter, to study the emission of gravitational waves from relativistic sources without any kind of post-Newtonian approximation.* For instance this is the case for the relativistic motion (but still in the weak field regime) of the binaries before the beginning of the final inspiral phase: it is known that in this phase the post-Newtonian approximation does not work and that , till now, only numerical gravity may help. In a future paper [44] we will add a relativistic perfect fluid, described by suitable Lagrangian [45] or Eulerian [46] variables, to tetrad gravity, we will define a Hamiltonian linearization of the system in the completely fixed 3-orthogonal gauge, we will find the Hamilton equations for the DO's both of the gravitational field and of the fluid, we will find the relativistic version of the Newton and gravito-magnetic action-at-a-distance potentials acting inside the fluid present in the weak ADM energy and finally, by using a multipolar expansion, we will find the relativistic counterpart of the post-Newtonian quadrupole emission formula.

Let us remark that till now we have a treatment of the generation of gravitational waves from a compact localized source of size R and mean internal velocity v only [22] for *nearly Newtonian slow motion* sources for which $v << c$, $\frac{\lambda}{2\pi} >> R$: outgoing gravitational waves are observed in the *radiation zone* (far field, $r >> \frac{\lambda}{2\pi}$), while deep in the *near zone* ($R < r << \frac{\lambda}{2\pi}$), for example $r \leq 1000\, \frac{\lambda}{2\pi}$, vacuum Newtonian gravitation theory is valid. On the contrary with our approach in suitable 4-coordinates we are going to obtain a *weak field approximation but with fast relativistic motion in the source* subject to the restriction that the total invariant mass and the mass currents are compatible with the weak field requirement. This is enough to get relativistic results conceptually equivalent to the re-summation of the post-Newtonian expansion.

Moreover we will have to explore whether our Hamiltonian approach is suitable for doing Hamiltonian numerical gravity on the full non-linearized theory.

APPENDIX A: RESULTS ON TETRAD GRAVITY.

In this Appendix we reproduce those results of Ref. [3], which are needed in this paper.

The Lichnerowicz equation in the 3-orthogonal gauge $\pi_\phi(\tau,\vec\sigma)\approx 0$ given in Eq.(193) of Ref. [3] is (see Eq.(A4) for the definition of the $\mathcal{T}^u_{(a)r}$'s)

$$\begin{aligned}
&(-\tilde{\triangle}[r_{\bar a}]+\frac{1}{8}{}^3\tilde{R}[r_{\bar a}])(\tau,\vec\sigma)\phi(\tau,\vec\sigma)=\frac{12\pi^2G^2}{c^6}\Big[2(\phi^{-7}\sum_{\bar a}\pi^2_{\bar a})(\tau,\vec\sigma)+\\
&\quad+4\Big(\phi^{-5}\sum_u e^{\frac{1}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a u}r_{\bar a}}\sum_{\bar b}\gamma_{\bar b u}\pi_{\bar b}\Big)(\tau,\vec\sigma)\times\\
&\qquad\int d^3\sigma_1\sum_r\delta^u_{(a)}\mathcal{T}^u_{(a)r}(\vec\sigma,\vec\sigma_1,\tau)\Big(\phi^{-2}e^{-\frac{1}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a r}r_{\bar a}}\sum_{\bar b}\gamma_{\bar b r}\pi_{\bar b}\Big)(\tau,\vec\sigma_1)+\\
&\quad+\phi^{-3}(\tau,\vec\sigma)\int d^3\sigma_1 d^3\sigma_2\Big(\sum_u e^{\frac{2}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a u}r_{\bar a}(\tau,\vec\sigma)}\times\\
&\qquad\sum_r\mathcal{T}^u_{(a)r}(\vec\sigma,\vec\sigma_1,\tau)\Big(\phi^{-2}e^{-\frac{1}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a r}r_{\bar a}}\sum_{\bar b}\gamma_{\bar b r}\pi_{\bar b}\Big)(\tau,\vec\sigma_1)\times\\
&\qquad\sum_s\mathcal{T}^u_{(a)s}(\vec\sigma,\vec\sigma_2,\tau)\Big(\phi^{-2}e^{-\frac{1}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a s}r_{\bar a}}\sum_{\bar c}\gamma_{\bar c s}\pi_{\bar c}\Big)(\tau,\vec\sigma_2)+\\
&\quad+\sum_{uv}e^{\frac{1}{\sqrt{3}}\sum_{\bar a}(\gamma_{\bar a u}+\gamma_{\bar a v})r_{\bar a}(\tau,\vec\sigma)}(\delta^u_{(b)}\delta^v_{(a)}-\delta^u_{(a)}\delta^v_{(b)})\times\\
&\qquad\sum_r\mathcal{T}^u_{(a)r}(\vec\sigma,\vec\sigma_1,\tau)\Big(\phi^{-2}e^{-\frac{1}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a r}r_{\bar a}}\sum_{\bar b}\gamma_{\bar b r}\pi_{\bar b}\Big)(\tau,\vec\sigma_1)\\
&\qquad\sum_s\mathcal{T}^v_{(b)s}(\vec\sigma,\vec\sigma_2,\tau)\Big(\phi^{-2}e^{-\frac{1}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a s}r_{\bar a}}\sum_{\bar c}\gamma_{\bar c s}\pi_{\bar c}\Big)(\tau,\vec\sigma_2)\Big)\Big].
\end{aligned}\tag{A1}$$

This equation for the conformal factor $\phi(\tau,\vec\sigma)$ of the 3-metric is implied by the superhamiltonian constraint, Eq.(191) of Ref. [3], which is

$$\begin{aligned}
\tilde{\mathcal{H}}_R(\tau,\vec\sigma)\,|_{\pi_\phi=0}&=\epsilon\phi(\tau,\vec\sigma)\Big[\frac{c^3}{16\pi G}(-8\tilde{\triangle}[r_{\bar a}]+{}^3\tilde{R}[r_{\bar a}])\phi-\\
&-\frac{6\pi G}{c^3}\Big[\Big(2\phi^{-7}\sum_{\bar a}\pi^2_{\bar a}\Big)(\tau,\vec\sigma)+\\
&+4\Big(\phi^{-5}\sum_u e^{\frac{1}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a u}r_{\bar a}}\sum_{\bar b}\gamma_{\bar b u}\pi_{\bar b}\Big)(\tau,\vec\sigma)\times\\
&\quad\int d^3\sigma_1\sum_r\delta^u_{(a)}\mathcal{T}^u_{(a)r}(\vec\sigma,\vec\sigma_1,\tau)\Big(\phi^{-2}e^{-\frac{1}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a r}r_{\bar a}}\sum_{\bar b}\gamma_{\bar b r}\pi_{\bar b}\Big)(\tau,\vec\sigma_1)+\\
&+\phi^{-3}(\tau,\vec\sigma)\int d^3\sigma_1 d^3\sigma_2\Big(\sum_u e^{\frac{2}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a u}r_{\bar a}(\tau,\vec\sigma)}\times\\
&\quad\sum_r\mathcal{T}^u_{(a)r}(\vec\sigma,\vec\sigma_1,\tau)\Big(\phi^{-2}e^{-\frac{1}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a r}r_{\bar a}}\sum_{\bar b}\gamma_{\bar b r}\pi_{\bar b}\Big)(\tau,\vec\sigma_1)\times\\
&\quad\sum_s\mathcal{T}^u_{(a)s}(\vec\sigma,\vec\sigma_2,\tau)\Big(\phi^{-2}e^{-\frac{1}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a s}r_{\bar a}}\sum_{\bar c}\gamma_{\bar c s}\pi_{\bar c}\Big)(\tau,\vec\sigma_2)+\\
&+\sum_{uv}e^{\frac{1}{\sqrt{3}}\sum_{\bar a}(\gamma_{\bar a u}+\gamma_{\bar a v})r_{\bar a}(\tau,\vec\sigma)}(\delta^u_{(b)}\delta^v_{(a)}-\delta^u_{(a)}\delta^v_{(b)})\times
\end{aligned}$$

$$\begin{aligned}
&\sum_r \mathcal{T}^u_{(a)r}(\vec{\sigma},\vec{\sigma}_1,\tau)\Big(\phi^{-2}e^{-\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}r}r_{\bar{a}}}\sum_{\bar{b}}\gamma_{\bar{b}r}\pi_{\bar{b}}\Big)(\tau,\vec{\sigma}_1)\\
&\sum_s \mathcal{T}^v_{(b)s}(\vec{\sigma},\vec{\sigma}_2,\tau)\Big(\phi^{-2}e^{-\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}s}r_{\bar{a}}}\sum_{\bar{c}}\gamma_{\bar{c}s}\pi_{\bar{c}}\Big)(\tau,\vec{\sigma}_2)\Big)\Big]\Big] \approx 0. \qquad \text{(A2)}
\end{aligned}$$

The scalar curvature and the Laplace-Beltrami operator appearing in these equations, Eq.(190) of Ref. [3], are

$$\begin{aligned}
{}^3\hat{R}[\phi,r_{\bar{a}}] &= -\sum_{uv}\{(2\partial_v ln\,\phi+\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}u}\partial_v r_{\bar{a}})(4\partial_v ln\,\phi-\frac{1}{\sqrt{3}}\sum_{\bar{b}}\gamma_{\bar{b}u}\partial_v r_{\bar{b}})+\\
&+\phi^{-4}e^{\frac{2}{\sqrt{3}}\sum_{\bar{c}}\gamma_{\bar{c}v}r_{\bar{c}}}[2\partial_v^2 ln\,\phi+\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}u}\partial_v^2 r_{\bar{a}}+\\
&+\frac{2}{\sqrt{3}}(2\partial_v ln\,\phi+\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}u}\partial_v r_{\bar{a}})\sum_{\bar{b}}(\gamma_{\bar{b}u}-\gamma_{\bar{b}v})\partial_v r_{\bar{b}}-\\
&-(2\partial_v ln\,\phi+\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}v}\partial_v r_{\bar{a}})(2\partial_v ln\,\phi+\frac{1}{\sqrt{3}}\sum_{\bar{b}}\gamma_{\bar{b}u}\partial_v r_{\bar{b}}]\}+\\
&+\phi^{-4}\sum_u e^{\frac{2}{\sqrt{3}}\sum_{\bar{c}}\gamma_{\bar{c}u}r_{\bar{c}}}[-2\partial_u^2 ln\,\phi+\frac{2}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}u}\partial_u^2 r_{\bar{a}}+\\
&+(2\partial_u ln\,\phi+\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}u}\partial_u r_{\bar{a}})(2\partial_u ln\,\phi-\frac{2}{\sqrt{3}}\sum_{\bar{b}}\gamma_{\bar{b}u}\partial_u r_{\bar{b}})],\\
\tilde{\triangle}[r_{\bar{a}}] &= \partial_r[{}^3\tilde{g}^{rs}\,\partial_s] = {}^3\tilde{g}^{rs}\,{}^3\tilde{\nabla}_r\,{}^3\tilde{\nabla}_s = \sum_r e^{-\frac{2}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}r}r_{\bar{a}}}[\partial_r^2-\frac{2}{\sqrt{3}}\sum_{\bar{b}}\gamma_{\bar{b}r}\partial_r r_{\bar{b}}\partial_r]. \qquad \text{(A3)}
\end{aligned}$$

If $\tilde{\pi}^{\vec{\alpha}}_{(a)}(\tau,\vec{\sigma})\approx 0$ and $\tilde{\pi}^{\vec{\xi}}_r(\tau,\vec{\sigma})\approx 0$ are the Abelianized momenta corresponding to the rotation and super-momentum constraints, the connection of the old cotriad momenta to the final ones, see Eq.(1.4), after the quasi-Shanmugadhasan canonical transformation (1.3), i.e. Eq.(185) of Ref. [3], is

$$\begin{aligned}
{}^3\hat{\tilde{\pi}}^r_{(a)}(\tau,\vec{\sigma}) &= {}^3\tilde{\pi}^r_{(a)}(\tau,\vec{\sigma})|_{\alpha_{(a)}=0,\xi^r=\sigma^r,\tilde{\pi}^{\vec{\alpha}}_{(a)}=\tilde{\pi}^{\vec{\xi}}_r=0}=\\
&=\sum_s\int d^3\sigma_1\mathcal{K}^r_{(a)s}(\vec{\sigma},\vec{\sigma}_1;\tau)\,\tilde{\Pi}^s(\tau,\vec{\sigma}_1)=\\
&=\sum_s\int d^3\sigma_1\mathcal{K}^r_{(a)s}(\vec{\sigma},\vec{\sigma}_1;\tau)(\phi^{-2}e^{-\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}s}r_{\bar{a}}})(\tau,\vec{\sigma}_1)\\
&\quad\Big[\frac{\pi_\phi}{6\,\phi}+\sqrt{3}\sum_{\bar{b}}\gamma_{\bar{b}s}\pi_{\bar{b}}\Big](\tau,\vec{\sigma}_1),\\
\rightarrow_{\pi_\phi\rightarrow 0}\quad &\sqrt{3}\sum_{s,\bar{b}}\gamma_{\bar{b}s}\int d^3\sigma_1\mathcal{K}^r_{(a)s}(\vec{\sigma},\vec{\sigma}_1;\tau|\phi,r_{\bar{a}}](\phi^{-2}e^{-\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}s}r_{\bar{a}}}\pi_{\bar{b}})(\tau,\vec{\sigma}_1),
\end{aligned}$$

with the kernel

$$\mathcal{K}^r_{(a)s}(\vec{\sigma},\vec{\sigma}_1,\tau)=\tilde{\mathcal{K}}^r_{(a)s}(\vec{\sigma},\vec{\sigma}_1,\tau|\phi,r_{\bar{a}}]\overset{def}{=}\delta^r_{(a)}\delta^r_s\delta^3(\vec{\sigma},\vec{\sigma}_1)+\mathcal{T}^r_{(a)s}(\vec{\sigma},\vec{\sigma}_1,\tau),$$

$$\mathcal{T}^r_{(a)s}(\vec{\sigma},\vec{\sigma}_1,\tau)=\tilde{\mathcal{T}}^r_{(a)s}(\vec{\sigma},\vec{\sigma}_1,\tau|\phi,r_{\bar{a}}]=$$

$$= -Q_s(\tau ,\vec{\sigma}_1)\frac{\partial G^{rs}_{(a)}(\vec{\sigma},\vec{\sigma}_1;\tau |\phi ,r_{\bar a}]}{\partial \sigma_1^s} - \sum_v \frac{\partial Q_s(\tau ,\vec{\sigma}_1)}{\partial \sigma_1^v} G^{rv}_{(a)}(\vec{\sigma},\vec{\sigma}_1;\tau |\phi ,r_{\bar a}], \tag{A4}$$

The canonicity of the transformation (1.3) implies that the kernels $\mathcal{T}^r_{(o)s}$'s are determined by the kernels $G^{rs}_{(a)}$'s.

Eqs.(156) of Ref. [3], giving the partial differential equations for the kernels $G^{ru}_{(a)}$'s, are

$$\begin{aligned}
&\epsilon_{(a)(r_1)(b)}\delta^s_{r_1}\frac{\delta^3(\vec{\sigma}_1,\vec{\sigma})}{Q_{r_1}(\tau ,\vec{\sigma})} - \sum_u \epsilon_{(u)(r_1)(b)}\frac{Q_u(\tau ,\vec{\sigma})}{Q_{r_1}(\tau ,\vec{\sigma})}\frac{\partial G^{su}_{(a)}(\vec{\sigma}_1,\vec{\sigma};\tau)}{\partial \sigma^{r_1}} =\\
&= \epsilon_{(a)(r_2)(b)}\delta^s_{r_2}\frac{\delta^3(\vec{\sigma}_1,\vec{\sigma})}{Q_{r_2}(\tau ,\vec{\sigma})} - \sum_u \epsilon_{(u)(r_2)(b)}\frac{Q_u(\tau ,\vec{\sigma})}{Q_{r_2}(\tau ,\vec{\sigma})}\frac{\partial G^{su}_{(a)}(\vec{\sigma}_1,\vec{\sigma};\tau)}{\partial \sigma^{r_2}},\\
&\Downarrow\\
&\frac{1}{Q^2_{r_1}(\tau ,\vec{\sigma})}\frac{\partial G^{sr_2}_{(a)}(\vec{\sigma}_1,\vec{\sigma};\tau)}{\partial \sigma^{r_1}} + \frac{1}{Q^2_{r_2}(\tau ,\vec{\sigma})}\frac{\partial G^{sr_1}_{(a)}(\vec{\sigma}_1,\vec{\sigma};\tau)}{\partial \sigma^{r_2}} =\\
&= \Big[\frac{\delta_{(a)r_1}\delta^s_{r_2}}{Q_{r_1}(\tau ,\vec{\sigma})Q^2_{r_2}(\tau ,\vec{\sigma})} + \frac{\delta_{(a)r_2}\delta^s_{r_1}}{Q^2_{r_1}(\tau ,\vec{\sigma})Q_{r_2}(\tau ,\vec{\sigma})}\Big]\delta^3(\vec{\sigma}_1,\vec{\sigma}),\\
&\Downarrow\\
&1)\ s=a \quad \text{homogeneous equations :}\\
&\frac{1}{Q^2_1(\tau ,\vec{\sigma})}\frac{\partial G^{a2}_{(a)}(\vec{\sigma}_1,\vec{\sigma};\tau)}{\partial \sigma^1} + \frac{1}{Q^2_2(\tau ,\vec{\sigma})}\frac{\partial G^{a1}_{(a)}(\vec{\sigma}_1,\vec{\sigma};\tau)}{\partial \sigma^2} =\\
&= \frac{1}{Q^2_2(\tau ,\vec{\sigma})}\frac{\partial G^{a3}_{(a)}(\vec{\sigma}_1,\vec{\sigma};\tau)}{\partial \sigma^2} + \frac{1}{Q^2_3(\tau ,\vec{\sigma})}\frac{\partial G^{a2}_{(a)}(\vec{\sigma}_1,\vec{\sigma};\tau)}{\partial \sigma^3} =\\
&= \frac{1}{Q^2_3(\tau ,\vec{\sigma})}\frac{\partial G^{a1}_{(a)}(\vec{\sigma}_1,\vec{\sigma};\tau)}{\partial \sigma^3} + \frac{1}{Q^2_1(\tau ,\vec{\sigma})}\frac{\partial G^{a3}_{(a)}(\vec{\sigma}_1,\vec{\sigma};\tau)}{\partial \sigma^1} = 0, \qquad a=1,2,3;\\
&2)\ s\not= a\ [s\not= r,\, r\not= a] \quad \text{in-homogeneous equations :}\\
&\frac{1}{Q^2_s(\tau ,\vec{\sigma})}\frac{\partial G^{sr}_{(a)}(\vec{\sigma}_1,\vec{\sigma};\tau)}{\partial \sigma^s} + \frac{1}{Q^2_r(\tau ,\vec{\sigma})}\frac{\partial G^{ss}_{(a)}(\vec{\sigma}_1,\vec{\sigma};\tau)}{\partial \sigma^r} =\\
&= \frac{1}{Q^2_r(\tau ,\vec{\sigma})}\frac{\partial G^{sa}_{(a)}(\vec{\sigma}_1,\vec{\sigma};\tau)}{\partial \sigma^r} + \frac{1}{Q^2_a(\tau ,\vec{\sigma})}\frac{\partial G^{sr}_{(a)}(\vec{\sigma}_1,\vec{\sigma};\tau)}{\partial \sigma^a} = 0,\\
&\frac{1}{Q^2_a(\tau ,\vec{\sigma})}\frac{\partial G^{ss}_{(a)}(\vec{\sigma}_1,\vec{\sigma};\tau)}{\partial \sigma^a} + \frac{1}{Q^2_s(\tau ,\vec{\sigma})}\frac{\partial G^{sa}_{(a)}(\vec{\sigma}_1,\vec{\sigma};\tau)}{\partial \sigma^s} = \frac{\delta^3(\vec{\sigma}_1;\vec{\sigma})}{Q_a(\tau ,\vec{\sigma})Q^2_s(\tau ,\vec{\sigma})}.
\end{aligned} \tag{A5}$$

Eqs.(159) of Ref. [3] for the kernels $\mathcal{K}^r_{(a)u}$'s, which ensure the simultaneous satisfaction of the rotation and super-momentum constraints, are $[Q_u = \phi^2\, e^{\frac{1}{\sqrt{3}}\sum_{\bar a} \gamma_{\bar a u}\, r_{\bar a}}]$

$$
\begin{aligned}
&\sum_r Q_r(\tau,\vec{\sigma})\Big[\delta_{(a)r}\mathcal{K}^r_{(b)u} - \delta_{(b)r}\mathcal{K}^r_{(a)u}\Big](\vec{\sigma},\vec{\sigma}_1;\tau) =\\
&= Q_u(\tau,\vec{\sigma}_1)\Big[Q_a(\tau,\vec{\sigma})\frac{\partial G^{au}_{(b)}(\vec{\sigma},\vec{\sigma}_1;\tau)}{\partial \sigma_1^u} - Q_b(\tau,\vec{\sigma})\frac{\partial G^{bu}_{(a)}(\vec{\sigma},\vec{\sigma}_1;\tau)}{\partial \sigma_1^u}\Big] +\\
&+\sum_v \frac{\partial Q_u(\tau,\vec{\sigma}_1)}{\partial \sigma_1^v}\Big[Q_a(\tau,\vec{\sigma})G^{av}_{(b)}(\vec{\sigma},\vec{\sigma}_1;\tau) - Q_b(\tau,\vec{\sigma})G^{bv}_{(a)}(\vec{\sigma},\vec{\sigma}_1;\tau)\Big] =\\
&= 0, \qquad a \neq b,
\end{aligned}
$$

$$
\begin{aligned}
&\hat{D}^{(\hat{\omega})}_{(a)(b)r}(\tau,\vec{\sigma})\mathcal{K}^r_{(b)u}(\vec{\sigma},\vec{\sigma}_1;\tau) =\\
&= \Big(\delta_{(a)(b)}\partial_r + \epsilon_{(a)(b)(c)}\, {}^3\hat{\omega}_{r(c)}(\tau,\vec{\sigma})\Big)\mathcal{K}^r_{(b)u}(\vec{\sigma},\vec{\sigma}_1;\tau) =\\
&= \Big(\delta_{(a)(b)}\partial_r + \sum_u(\delta_{(a)r}\delta_{(b)u} - \delta_{(a)u}\delta_{(b)r})\frac{\partial_u Q_r(\tau,\vec{\sigma})}{Q_u(\tau,\vec{\sigma})}\Big)\Big[\delta^r_{(b)}\delta_{(b)u}\delta^3(\vec{\sigma},\vec{\sigma}_1) -\\
&- Q_u(\tau,\vec{\sigma}_1)\frac{\partial G^{ru}_{(b)}(\vec{\sigma},\vec{\sigma}_1;\tau)}{\partial \sigma_1^u} - \sum_v \frac{\partial Q_u(\tau,\vec{\sigma}_1)}{\partial \sigma_1^v}\, G^{rv}_{(b)}(\vec{\sigma},\vec{\sigma}_1;\tau)\Big] = 0,\\
&{}^3\hat{\omega}_{r(a)}(\tau,\vec{\sigma}) = \sum_u \epsilon_{(a)(m)(n)}\delta_{(m)r}\delta_{(n)u}\frac{\partial_u Q_r(\tau,\vec{\sigma})}{Q_u(\tau,\vec{\sigma})} \qquad \text{(A6)}
\end{aligned}
$$

The cotriads, the triads, the 3-metric, the Christoffel symbols and the spin connection on the WSW hyper-surfaces of our gauge, given in Eq.(184) of Ref. [3], have the following expressions in the canonical basis (1.3)

$$
\begin{aligned}
{}^3\hat{e}_{(a)r} &= \delta_{(a)r}\phi^2 e^{\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}r}r_{\bar{a}}},\\
{}^3\hat{e}^r_{(a)} &= \delta^r_{(a)}\phi^{-2} e^{-\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}r}r_{\bar{a}}},\\
{}^3\hat{g}_{rs} &= \delta_{rs}\phi^4 e^{\frac{2}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}r}r_{\bar{a}}},\\
{}^3\hat{g}^{rs} &= \delta^{rs}\phi^{-4} e^{-\frac{2}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}r}r_{\bar{a}}},\\
{}^3\hat{e} &= \sqrt{\hat{\gamma}} = \phi^6,\\
{}^3\hat{\Gamma}^r_{uv} &= -\delta_{uv}\sum_s \delta^r_s e^{\frac{2}{\sqrt{3}}\sum_{\bar{a}}(\gamma_{\bar{a}u}-\gamma_{\bar{a}s})r_{\bar{a}}}\Big[2\partial_s ln\,\phi + \frac{1}{\sqrt{3}}\sum_{\bar{b}}\gamma_{\bar{b}u}\partial_s r_{\bar{b}}\Big] +\\
&+\delta^r_u\Big[2\partial_v ln\,\phi + \frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}u}\partial_v r_{\bar{a}}\Big] + \delta^r_v\Big[2\partial_u ln\,\phi + \frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}v}\partial_u r_{\bar{a}}\Big],\\
&\sum_u {}^3\hat{\Gamma}^u_{uv} = 6\partial_v ln\,\phi,\\
{}^3\hat{\omega}_{r(a)} &= \epsilon_{(a)(b)(c)}\,\delta_{(b)r}\delta_{(c)u}\, e^{\frac{1}{\sqrt{3}}\sum_{\bar{a}}(\gamma_{\bar{a}r}-\gamma_{\bar{a}u})r_{\bar{a}}}\Big[2\partial_u ln\,\phi + \frac{1}{\sqrt{3}}\sum_{\bar{b}}\gamma_{\bar{b}r}\partial_u r_{\bar{b}}\Big]\\
&\rightarrow_{r_{\bar{a}}\rightarrow 0}\ \epsilon_{(a)(b)(c)}\,\delta_{(b)r}\delta_{(c)u}\, 2\partial_u ln\,\phi. \qquad \text{(A7)}
\end{aligned}
$$

Eqs.(102) and (105) of Ref. [3] define the Green function of the covariant divergence

$$\hat{D}^{(\omega)}_{(a)(b)r}(\tau,\vec{\sigma})\,\zeta^{(\omega)r}_{(b)(c)}(\vec{\sigma},{\vec{\sigma}}^{'};\tau) = -\delta_{(a)(c)}\delta^3(\vec{\sigma},{\vec{\sigma}}^{'}). \tag{A8}$$

$$\begin{aligned}
&\zeta^{(\omega)r}_{(a)(b)}(\vec{\sigma},{\vec{\sigma}}^{'};\tau) = d^r_{\gamma_{pp'}}(\vec{\sigma},{\vec{\sigma}}^{'})\Big(P_{\gamma_{pp'}}\, e^{\int_{{\vec{\sigma}}^{'}}^{\vec{\sigma}} d\sigma_1^s\, \hat{R}^{(c)}\, {}^3\omega_{s(c)}(\tau,\vec{\sigma}_1)}\Big)_{(a)(b)}\\
\rightarrow_{\phi\rightarrow 1, r_{\bar{a}}\rightarrow 0}\quad &\zeta^{(o)r}_{(a)(b)}(\vec{\sigma},\vec{\sigma}_1) = -\frac{\sigma^r-\sigma^r_1}{4\pi|\vec{\sigma}-\vec{\sigma}_1|^3}\delta_{(a)(b)} = -\delta_{(a)(b)}\frac{\partial}{\partial\sigma^r_1}\frac{1}{4\pi|\vec{\sigma}-\vec{\sigma}_1|} =\\
&= \delta_{(a)(b)}\, c^r(\vec{\sigma}-\vec{\sigma}_1),
\end{aligned} \tag{A9}$$

since the linearization reduces $d^r_{\gamma}(\vec{\sigma},\vec{\sigma}_1)$ to the Green function of the ordinary divergence $c^r(\vec{\sigma}-\vec{\sigma}_1)$.

The weak and strong ADM Poincare' generators of Eqs.(25) of Ref. [3] are

$$\begin{aligned}
\hat{P}^{\tau}_{ADM} &= \int d^3\sigma\epsilon[\frac{c^3}{16\pi G}\sqrt{\gamma}\ {}^3g^{\check{r}\check{s}}({}^3\Gamma^{\check{u}}_{\check{r}\check{v}}\,{}^3\Gamma^{\check{v}}_{\check{s}\check{u}} - {}^3\Gamma^{\check{u}}_{\check{r}\check{s}}\,{}^3\Gamma^{\check{v}}_{\check{v}\check{u}}) -\\
&- \frac{8\pi G}{c^3\sqrt{\gamma}}{}^3G_{\check{r}\check{s}\check{u}\check{v}}\,{}^3\tilde{\Pi}^{\check{r}\check{s}}\,{}^3\tilde{\Pi}^{\check{u}\check{v}}](\tau,\vec{\sigma}),\\
\hat{P}^{\check{r}}_{ADM} &= -2\int d^3\sigma\,{}^3\Gamma^{\check{r}}_{\check{s}\check{u}}(\tau,\vec{\sigma})\,{}^3\tilde{\Pi}^{\check{s}\check{u}}(\tau,\vec{\sigma}),\\
\hat{J}^{\tau\check{r}}_{ADM} &= -\hat{J}^{\check{r}\tau}_{ADM} = \int d^3\sigma\epsilon\{\sigma^{\check{r}}\\
&\qquad [\frac{c^3}{16\pi G}\sqrt{\gamma}\ {}^3g^{\check{n}\check{s}}({}^3\Gamma^{\check{u}}_{\check{n}\check{v}}\,{}^3\Gamma^{\check{v}}_{\check{s}\check{u}} - {}^3\Gamma^{\check{u}}_{\check{n}\check{s}}\,{}^3\Gamma^{\check{v}}_{\check{v}\check{u}}) - \frac{8\pi G}{c^3\sqrt{\gamma}}{}^3G_{\check{n}\check{s}\check{u}\check{v}}\,{}^3\tilde{\Pi}^{\check{n}\check{s}}\,{}^3\tilde{\Pi}^{\check{u}\check{v}}] +\\
&+ \frac{c^3}{16\pi G}\delta^{\check{r}}_{\check{u}}({}^3g_{\check{v}\check{s}} - \delta_{\check{v}\check{s}})\partial_{\check{n}}[\sqrt{\gamma}({}^3g^{\check{n}\check{s}}\,{}^3g^{\check{u}\check{v}} - {}^3g^{\check{n}\check{u}}\,{}^3g^{\check{s}\check{v}})]\}(\tau,\vec{\sigma}),\\
\hat{J}^{\check{r}\check{s}}_{ADM} &= \int d^3\sigma[(\sigma^{\check{r}}\,{}^3\Gamma^{\check{s}}_{\check{u}\check{v}} - \sigma^{\check{s}}\,{}^3\Gamma^{\check{r}}_{\check{u}\check{v}})\,{}^3\tilde{\Pi}^{\check{u}\check{v}}](\tau,\vec{\sigma}),\\
&\\
P^{\tau}_{ADM} &= \hat{P}^{\tau}_{ADM} + \int d^3\sigma\tilde{\mathcal{H}}(\tau,\vec{\sigma}) \approx \hat{P}^{\tau}_{ADM},\\
P^{\check{r}}_{ADM} &= \hat{P}^{\check{r}}_{ADM} + \int d^3\sigma\,{}^3\tilde{\mathcal{H}}^{\check{r}}(\tau,\vec{\sigma}) \approx \hat{P}^{\check{r}}_{ADM},\\
J^{\tau\check{r}}_{ADM} &= \hat{J}^{\tau\check{r}}_{ADM} + \frac{1}{2}\int d^3\sigma\sigma^{\check{r}}\,\tilde{\mathcal{H}}(\tau,\vec{\sigma}) \approx \hat{J}^{\tau\check{r}}_{ADM},\\
J^{\check{r}\check{s}}_{ADM} &= \hat{J}^{\check{r}\check{s}}_{ADM} + \int d^3\sigma[\sigma^{\check{s}}\,{}^3\tilde{\mathcal{H}}^{\check{r}}(\tau,\vec{\sigma}) - \sigma^{\check{r}}\,{}^3\tilde{\mathcal{H}}^{\check{s}}(\tau,\vec{\sigma})] \approx \hat{J}^{\check{r}\check{s}}_{ADM}.
\end{aligned} \tag{A10}$$

Eq.(197) of Ref. [3] identifies the Hamiltonian of the rest-frame instant form of tetrad gravity with the weak ADM energy

$$\hat{H}^{(WSW)'}_{(D)ADM} = -\epsilon\hat{P}^{\tau}_{ADM,R} = \hat{E}_{ADM}. \tag{A11}$$

The expression of the weak ADM Poincare' generators after the quasi-Shanmugadhasan transformation (1.3) and the restriction to the $\pi_{\phi}(\tau,\vec{\sigma}) = 0$ 3-orthogonal gauge, given in Eqs.(227) of Appendix B of Ref. [3], is

$$\hat{P}^{\tau}_{ADM,R} = \epsilon\int d^3\sigma\Big(\frac{c^3}{16\pi G}\Big[\phi^2\sum_r e^{-\frac{2}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}r}r_{\bar{a}}}\times$$

$$
\begin{aligned}
&\Big(8(\partial_r ln\, \phi)^2 - \frac{1}{3}\sum_{\bar b}(\partial_r r_{\bar b})^2 - \\
&- \frac{4}{\sqrt{3}}\partial_r ln\, \phi \sum_{\bar b}\gamma_{\bar b r}\partial_r r_{\bar b} + \frac{2}{3}(\sum_{\bar b}\gamma_{\bar b r}\partial_r r_{\bar b})^2)\Big)\Big](\tau,\vec\sigma) - \\
&- \frac{6\pi G}{c^3}\phi^{-2}(\tau,\vec\sigma)\Big[2(\phi^{-4}\sum_{\bar a}\pi^2_{\bar a})(\tau,\vec\sigma) + \\
&+ 4(\phi^{-2}\sum_u e^{\frac{1}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a u} r_{\bar a}}\sum_{\bar b}\gamma_{\bar b u}\pi_{\bar b})(\tau,\vec\sigma) \times \\
&\int d^3\sigma_1 \sum_r \delta^u_{(a)} \tilde{\mathcal{T}}^u_{(a)r}(\vec\sigma,\vec\sigma_1,\tau|\phi, r_{\bar a}]\Big(\phi^{-2}e^{-\frac{1}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a r} r_{\bar a}}\sum_{\bar b}\gamma_{\bar b r}\pi_{\bar b}\Big)(\tau,\vec\sigma_1) + \\
&+ \int d^3\sigma_1 d^3\sigma_2 \Big(\sum_u e^{\frac{2}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a u} r_{\bar a}(\tau,\vec\sigma)} \times \\
&\sum_r \tilde{\mathcal{T}}^u_{(a)r}(\vec\sigma,\vec\sigma_1,\tau|\phi, r_{\bar a}]\Big(\phi^{-2}e^{-\frac{1}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a r} r_{\bar a}}\sum_{\bar b}\gamma_{\bar b r}\pi_{\bar b}\Big)(\tau,\vec\sigma_1) \times \\
&\sum_s \tilde{\mathcal{T}}^u_{(a)s}(\vec\sigma,\vec\sigma_2,\tau|\phi, r_{\bar a}]\Big(\phi^{-2}e^{-\frac{1}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a s} r_{\bar a}}\sum_{\bar c}\gamma_{\bar c s}\pi_{\bar c}\Big)(\tau,\vec\sigma_2) + \\
&+ \sum_{uv} e^{\frac{1}{\sqrt{3}}\sum_{\bar a}(\gamma_{\bar a u}+\gamma_{\bar a v}) r_{\bar a}(\tau,\vec\sigma)}(\delta^u_{(b)}\delta^v_{(a)} - \delta^u_{(a)}\delta^v_{(b)}) \times \\
&\sum_r \tilde{\mathcal{T}}^u_{(a)r}(\vec\sigma,\vec\sigma_1,\tau|\phi, r_{\bar a}]\Big(\phi^{-2}e^{-\frac{1}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a r} r_{\bar a}}\sum_{\bar b}\gamma_{\bar b r}\pi_{\bar b}\Big)(\tau,\vec\sigma_1) \\
&\sum_s \tilde{\mathcal{T}}^v_{(b)s}(\vec\sigma,\vec\sigma_2,\tau|\phi, r_{\bar a}]\Big(\phi^{-2}e^{-\frac{1}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a s} r_{\bar a}}\sum_{\bar c}\gamma_{\bar c s}\pi_{\bar c}\Big)(\tau,\vec\sigma_2)\Big)\Big]\Big),
\end{aligned}
$$

$$
\begin{aligned}
\hat P^r_{ADM,R} = &- \int d^3\sigma \phi^{-2}(\tau,\vec\sigma)\Big(\phi^{-2}(\tau,\vec\sigma) \\
&\Big[e^{-\frac{2}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a r} r_{\bar a}}\sum_{\bar b}\partial_r r_{\bar b}\pi_{\bar b} + \\
&+ 2\sqrt{3}e^{-\frac{2}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a r} r_{\bar a}}(2\partial_r ln\, \phi + \frac{1}{\sqrt{3}}\sum_{\bar b}\gamma_{\bar b r}\partial_r r_{\bar b}) \\
&\sum_{\bar c}\gamma_{\bar c r}\pi_{\bar c}\Big](\tau,\vec\sigma) + \\
&+ \sqrt{3}\int d^3\sigma_1 \sum_s \Big[-\sum_u \Big(e^{\frac{1}{\sqrt{3}}\sum_{\bar a}(\gamma_{\bar a u} - 2\gamma_{\bar a r}) r_{\bar a}}(2\partial_r ln\, \phi + \frac{1}{\sqrt{3}}\sum_{\bar b}\gamma_{\bar b u}\partial_r r_{\bar b})\Big)(\tau,\vec\sigma) \\
&\delta^u_{(a)}\tilde{\mathcal{T}}^u_{(a)s}(\vec\sigma,\vec\sigma_1,\tau|\phi, r_{\bar a}] \\
&+ \sum_{uv}\Big(e^{-\frac{1}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a u} r_{\bar a}}\Big(\delta^r_u(2\partial_v ln\, \phi + \frac{1}{\sqrt{3}}\sum_{\bar b}\gamma_{\bar b u}\partial_v r_{\bar b}) + \\
&+ \delta^r_v(2\partial_u ln\, \phi + \frac{1}{\sqrt{3}}\sum_{\bar b}\gamma_{\bar b v}\partial_u r_{\bar b})\Big)\Big)(\tau,\vec\sigma) \\
&\delta^u_{(a)}\tilde{\mathcal{T}}^v_{(a)s}(\vec\sigma,\vec\sigma_1,\tau|\phi, r_{\bar a}]\Big] \\
&\Big(\phi^{-2}e^{-\frac{1}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a s} r_{\bar a}}\sum_{\bar c}\gamma_{\bar c s}\pi_{\bar c}\Big)(\tau,\vec\sigma_1)\Big),
\end{aligned}
$$

$$
\begin{aligned}
\hat{J}^{rs}_{ADM,R} &= \int d^3\sigma \phi^{-2}(\tau,\vec{\sigma})\Big(\phi^{-2}(\tau,\vec{\sigma}) \\
&\quad \Big[e^{-\frac{2}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}r}r_{\bar{a}}}\sum_{\bar{b}}(\sigma^r\partial_s-\sigma^s\partial_r)r_{\bar{b}}\pi_{\bar{b}}) + \\
&+ 2\sqrt{3}\sum_u e^{-\frac{2}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}u}r_{\bar{a}}}(\sigma^r\delta^s_u-\sigma^s\delta^r_u)(2\partial_u ln\,\phi+\frac{1}{\sqrt{3}}\sum_{\bar{b}}\gamma_{\bar{b}u}\partial_u r_{\bar{b}})\sum_{\bar{c}}\gamma_{\bar{c}u}\pi_{\bar{c}}\Big](\tau,\vec{\sigma}) + \\
&+ \sqrt{3}\int d^3\sigma_1\sum_w\Big[\Big(-\sigma^r\sum_u\Big(e^{\frac{1}{\sqrt{3}}\sum_{\bar{a}}(\gamma_{\bar{a}u}-2\gamma_{\bar{a}s})r_{\bar{a}}}(2\partial_s ln\,\phi+\frac{1}{\sqrt{3}}\sum_{\bar{b}}\gamma_{\bar{b}u}\partial_s r_{\bar{b}})\Big)(\tau,\vec{\sigma}) - \\
&- \sigma^s\sum_u\Big(e^{\frac{1}{\sqrt{3}}\sum_{\bar{a}}(\gamma_{\bar{a}u}-2\gamma_{\bar{a}r})r_{\bar{a}}}(2\partial_r ln\,\phi+\frac{1}{\sqrt{3}}\sum_{\bar{b}}\gamma_{\bar{b}u}\partial_r r_{\bar{b}})\Big)(\tau,\vec{\sigma})\Big) \\
&\quad \delta^u_{(a)}\tilde{\mathcal{T}}^u_{(a)w}(\vec{\sigma},\vec{\sigma}_1,\tau|\phi,r_{\bar{a}}] + \\
&+ \sum_{uv}\Big(e^{-\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}u}r_{\bar{a}}}\Big((\sigma^r\delta^s_u-\sigma^s\delta^r_u)(2\partial_v ln\,\phi+\frac{1}{\sqrt{3}}\sum_{\bar{b}}\gamma_{\bar{b}u}\partial_v r_{\bar{b}}) + \\
&+ (\sigma^r\delta^s_v-\sigma^s\delta^r_v)(2\partial_u ln\,\phi+\frac{1}{\sqrt{3}}\sum_{\bar{b}}\gamma_{\bar{b}v}\partial_u r_{\bar{b}})\Big)\Big)(\tau,\vec{\sigma}) \\
&\quad \delta^u_{(a)}\tilde{\mathcal{T}}^v_{(a)w}(\vec{\sigma},\vec{\sigma}_1,\tau|\phi,r_{\bar{a}}]\Big] \\
&\quad \Big(\phi^{-2}e^{-\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}s}r_{\bar{a}}}\sum_{\bar{c}}\gamma_{\bar{c}s}\pi_{\bar{c}}\Big)(\tau,\vec{\sigma}_1)\Big),
\end{aligned}
$$

$$
\begin{aligned}
\hat{J}^{\tau r}_{ADM,R} &= \epsilon\int d^3\sigma\sigma^r\Big(\frac{c^3}{16\pi G}\Big[\phi^2\sum_r e^{-\frac{2}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}r}r_{\bar{a}}} \times \\
&\quad \Big(8(\partial_r ln\,\phi)^2-\frac{1}{3}\sum_{\bar{b}}(\partial_r r_{\bar{b}})^2 - \\
&- \frac{4}{\sqrt{3}}\partial_r\phi\sum_{\bar{b}}\gamma_{\bar{b}r}\partial_r r_{\bar{b}}+\frac{2}{3}(\sum_{\bar{b}}\gamma_{\bar{b}r}\partial_r r_{\bar{b}})^2\Big)\Big](\tau,\vec{\sigma}) - \\
&- \frac{6\pi G}{c^3}\phi^{-2}(\tau,\vec{\sigma})\Big\{2(\phi^{-4}\sum_{\bar{a}}\pi^2_{\bar{a}})(\tau,\vec{\sigma}) + \\
&\quad +4\Big[\phi^{-2}\sum_u e^{\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}u}r_{\bar{a}}}\sum_{\bar{b}}\gamma_{\bar{b}u}\pi_{\bar{b}})(\tau,\vec{\sigma}) \times \\
&\qquad \int d^3\sigma_1\sum_m\delta^u_{(a)}\tilde{\mathcal{T}}^u_{(a)m}(\vec{\sigma},\vec{\sigma}_1,\tau|\phi,r_{\bar{a}}]\Big(\phi^{-2}e^{-\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}m}r_{\bar{a}}}\sum_{\bar{b}}\gamma_{\bar{b}m}\pi_{\bar{b}}\Big)(\tau,\vec{\sigma}_1) + \\
&\quad + \int d^3\sigma_1 d^3\sigma_2\Big(\sum_u e^{\frac{2}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}u}r_{\bar{a}}(\tau,\vec{\sigma})} \times \\
&\qquad \sum_m\tilde{\mathcal{T}}^u_{(a)m}(\vec{\sigma},\vec{\sigma}_1,\tau|\phi,r_{\bar{a}}]\Big(\phi^{-2}e^{-\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}m}r_{\bar{a}}}\sum_{\bar{b}}\gamma_{\bar{b}m}\pi_{\bar{b}}\Big)(\tau,\vec{\sigma}_1) \times \\
&\qquad \sum_s\tilde{\mathcal{T}}^u_{(a)s}(\vec{\sigma},\vec{\sigma}_2,\tau|\phi,r_{\bar{a}}]\Big(\phi^{-2}e^{-\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}s}r_{\bar{a}}}\sum_{\bar{c}}\gamma_{\bar{c}s}\pi_{\bar{c}}\Big)(\tau,\vec{\sigma}_2) + \\
&\quad + \sum_{uv}e^{\frac{1}{\sqrt{3}}\sum_{\bar{a}}(\gamma_{\bar{a}u}+\gamma_{\bar{a}v})r_{\bar{a}}(\tau,\vec{\sigma})}(\delta^u_{(b)}\delta^v_{(a)}-\delta^u_{(a)}\delta^v_{(b)}) \times
\end{aligned}
$$

$$\sum_m \tilde{\mathcal{T}}^u_{(a)m}(\vec{\sigma}, \vec{\sigma}_1, \tau|\phi, r_{\bar{a}}]\Big(\phi^{-2} e^{-\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}m} r_{\bar{a}}} \sum_{\bar{b}} \gamma_{\bar{b}m}\pi_{\bar{b}}\Big)(\tau, \vec{\sigma}_1)$$

$$\sum_s \tilde{\mathcal{T}}^v_{(b)s}(\vec{\sigma}, \vec{\sigma}_2, \tau|\phi, r_{\bar{a}}]\Big(\phi^{-2} e^{-\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}s} r_{\bar{a}}} \sum_{\bar{c}} \gamma_{\bar{c}s}\pi_{\bar{c}}\Big)(\tau, \vec{\sigma}_2)\,\Big)\Big]\Big\} -$$

$$- \frac{\epsilon c^3}{8\pi G}\int d^3\sigma \Big[\phi^{-2}\sum_{uv}\delta^r_u(\delta_{uv} - 1)(\phi^4 e^{\frac{2}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}u} r_{\bar{a}}} - 1)$$

$$e^{-\frac{2}{\sqrt{3}}\sum_{\bar{a}}(\gamma_{\bar{a}v} - \gamma_{\bar{a}u}) r_{\bar{a}}}(\partial_u ln\, \phi + \frac{1}{\sqrt{3}}\sum_{\bar{b}}(\gamma_{\bar{b}v} - \gamma_{\bar{b}u})\partial_u r_{\bar{b}})\Big](\tau, \vec{\sigma}). \qquad \text{(A12)}$$

Eqs.(194) of Ref. [3] give the equation for the determination of the lapse function $n(\tau, \vec{\sigma})$ as a consequence of the time constancy of the gauge fixing $\rho(\tau, \vec{\sigma}) = \frac{\pi_\phi(\tau, \vec{\sigma})}{2\,\phi(\tau, \vec{\sigma})} \approx 0$

$$\begin{aligned}
\partial_\tau \rho(\tau, \vec{\sigma}) &\,\circeq \{\rho(\tau, \vec{\sigma}), \hat{H}_{(D)ADM,R}\} = \int d^3\sigma_1 n(\tau, \vec{\sigma}_1)\{\rho(\tau, \vec{\sigma}), \hat{\mathcal{H}}_R(\tau, \vec{\sigma}_1)\} + \\
&+ \tilde{\lambda}_\tau(\tau)\{\rho(\tau, \vec{\sigma}), \hat{P}^\tau_{ADM,R}\} + \tilde{\lambda}_r(\tau)\{\rho(\tau, \vec{\sigma}), \hat{P}^r_{ADM,R}\} \approx \\
&\approx -\frac{1}{2}\phi(\tau, \vec{\sigma})\Big[\int d^3\sigma_1 n(\tau, \vec{\sigma}_1)\frac{\delta \hat{\mathcal{H}}_R(\tau, \vec{\sigma}_1)}{\delta \phi(\tau, \vec{\sigma})} + \\
&+ \tilde{\lambda}_\tau \frac{\delta \hat{P}^\tau_{ADM,R}}{\delta \phi(\tau, \vec{\sigma})} + \tilde{\lambda}_r(\tau)\frac{\delta \hat{P}^r_{ADM,R}}{\delta \phi(\tau, \vec{\sigma})}\Big] \approx 0, \\
\Rightarrow \qquad & n(\tau, \vec{\sigma}) - \hat{n}(\tau, \vec{\sigma}|r_{\bar{a}}, \pi_{\bar{a}}, \tilde{\lambda}_A] \approx 0, \\
\partial_\tau \qquad & \Big[n(\tau, \vec{\sigma}) - \hat{n}(\tau, \vec{\sigma}|r_{\bar{a}}, \pi_{\bar{a}}, \tilde{\lambda}_A]\Big] = \\
&= \lambda_n(\tau, \vec{\sigma}) - \{\hat{n}(\tau, \vec{\sigma}|r_{\bar{a}}, \pi_{\bar{a}}, \tilde{\lambda}_A], \hat{H}^{(WSW)}_{(D)ADM,R}\} \approx 0, \\
\Rightarrow \qquad & \lambda_n(\tau, \vec{\sigma}) \quad \text{determined;}
\end{aligned}$$

the rest-frame instant form expression of this equation is

$$\int d^3\sigma_1 \quad n(\tau, \vec{\sigma}_1)\frac{\delta \hat{\mathcal{H}}_R(\tau, \vec{\sigma}_1)}{\delta \phi(\tau, \vec{\sigma})} = -\epsilon \frac{\delta \hat{P}^\tau_{ADM,R}}{\delta \phi(\tau, \vec{\sigma})}. \qquad \text{(A13)}$$

For $\pi_\phi(\tau, \vec{\sigma}) = 0$ the shift functions $n_r(\tau, \vec{\sigma})$ of our gauge, as functions of the lapse function $n(\tau, \vec{\sigma})$, have the form given in Eq.(187) of Ref. [3], i.e.

$$\begin{aligned}
n_r(\tau, \vec{\sigma}) \approx & -\frac{\epsilon\, 4\sqrt{3}\pi G}{c^3}\Big[\phi^2 e^{\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}r} r_{\bar{a}}}\Big](\tau, \vec{\sigma})\int d^3\sigma_1[\epsilon - n(\tau, \vec{\sigma}_1)]\phi^{-2}(\tau, \vec{\sigma}_1) \\
& \sum_{wu} e^{\frac{1}{\sqrt{3}}\sum_{\bar{a}}(\gamma_{\bar{a}w} + \gamma_{\bar{a}u}) r_{\bar{a}}(\tau, \vec{\sigma}_1)}\,\Big(\delta_{wu}\delta_{(b)(d)} + \delta_{(b)u}\delta_{(d)w} - \delta_{(b)w}\delta_{(d)u}\Big) \\
& \sum_v \int d^3\sigma_2 \mathcal{K}^w_{(b)v}(\vec{\sigma}_1, \vec{\sigma}_2; \tau)\Big[\phi^{-2} e^{-\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}u} r_{\bar{a}}}\Big](\tau, \vec{\sigma}_2) \\
& \sum_{\bar{b}} \gamma_{\bar{b}v}\pi_{\bar{b}}(\tau, \vec{\sigma}_2)\, G^{ur}_{(d)}(\vec{\sigma}_1, \vec{\sigma}; \tau).
\end{aligned} \qquad \text{(A14)}$$

For $\pi_\phi = 0$ the extrinsic curvature of our WSW hyper-surfaces, the ADM metric momentum and the DeWitt supermetric, given by Eqs.(186) of Ref. [3], are

$$
\begin{aligned}
{}^3\hat{K}_{rs}(\tau,\vec{\sigma}) &= \frac{\epsilon\, 4\pi G}{c^3}\Big[e^{\frac{1}{\sqrt{3}}\sum_{\bar{c}}(\gamma_{\bar{c}r}+\gamma_{\bar{c}s})r_{\bar{c}}}\sum_u(\delta_{ru}\delta_{(a)s}+\delta_{su}\delta_{(a)r}-\delta_{rs}\delta_{(a)u})\\
&\quad e^{\frac{1}{\sqrt{3}}\sum_{\bar{c}}\gamma_{\bar{c}u}r_{\bar{c}}}\,{}^3\hat{\tilde{\pi}}^u_{(a)}\Big](\tau,\vec{\sigma}),\\[2ex]
{}^3\hat{K}(\tau,\vec{\sigma}) &= -\frac{\epsilon\, 4\pi G}{c^3}\Big[\phi^{-4}\sum_u\delta_{(a)u}e^{\frac{1}{\sqrt{3}}\sum_{\bar{c}}\gamma_{\bar{c}u}r_{\bar{c}}}\,{}^3\hat{\tilde{\pi}}^u_{(a)}\Big](\tau,\vec{\sigma}) =\\
&= -\frac{\epsilon\, 4\sqrt{3}\pi G}{c^3}\phi^{-4}(\tau,\vec{\sigma})\sum_u\delta_{(a)u}e^{\frac{1}{\sqrt{3}}\sum_{\bar{c}}\gamma_{\bar{c}u}r_{\bar{c}}(\tau,\vec{\sigma})}\\
&\quad \sum_s\sum_{\bar{b}}\gamma_{\bar{b}s}\int d^3\sigma_1\mathcal{K}^r_{(a)s}(\vec{\sigma},\vec{\sigma}_1;\tau|\phi,r_{\bar{a}}](\phi^{-2}e^{-\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}s}r_{\bar{a}}}\pi_{\bar{b}})(\tau,\vec{\sigma}_1),\\[3ex]
{}^3\hat{\tilde{\Pi}}^{rs}(\tau,\vec{\sigma}) &= \frac{1}{4}\Big[{}^3\hat{e}^r_{(a)}\,{}^3\hat{\tilde{\pi}}^s_{(a)}+{}^3\hat{e}^s_{(a)}\,{}^3\hat{\tilde{\pi}}^r_{(a)}\Big](\tau,\vec{\sigma}) =\\
&= \frac{1}{4}\phi^{-2}(\tau,\vec{\sigma})\Big[e^{-\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}r}r_{\bar{a}}}\delta^r_{(a)}\,{}^3\hat{\tilde{\pi}}^s_{(a)}+\\
&+ e^{-\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}s}r_{\bar{a}}}\delta^s_{(a)}\,{}^3\hat{\tilde{\pi}}^r_{(a)}\Big](\tau,\vec{\sigma}) =\\
&= \frac{\sqrt{3}}{4}\phi^{-2}(\tau,\vec{\sigma})\Big[e^{-\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}r}r_{\bar{a}}}\delta^r_{(a)}\\
&\quad \sum_u\int d^3\sigma_1\mathcal{K}^r_{(a)u}(\vec{\sigma},\vec{\sigma}_1,\tau|\phi,r_{\bar{a}},\tilde{\Pi}]+\\
&+ e^{-\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}s}r_{\bar{a}}}\delta^s_{(a)}\sum_u\int d^3\sigma_1\mathcal{K}^s_{(a)u}(\vec{\sigma},\vec{\sigma}_1,\tau|\phi,r_{\bar{a}},\tilde{\Pi}]\Big]\\
&\quad (\phi^{-2}e^{-\frac{1}{\sqrt{3}}\sum_{\bar{a}}\gamma_{\bar{a}u}r_{\bar{a}}})(\tau,\vec{\sigma}_1)\ \sum_{\bar{b}}\gamma_{\bar{b}u}\pi_{\bar{b}}(\tau,\vec{\sigma}_1),\\
{}^3\hat{G}_{rsuv}(\tau,\vec{\sigma}) &= \Big[{}^3\hat{g}_{ru}\,{}^3\hat{g}_{sv}+{}^3\hat{g}_{rv}\,{}^3\hat{g}_{su}-{}^3\hat{g}_{rs}\,{}^3\hat{g}_{uv}\Big](\tau,\vec{\sigma}) =\\
&= \phi^8(\tau,\vec{\sigma})\Big[e^{\frac{2}{\sqrt{3}}\sum_{\bar{a}}(\gamma_{\bar{a}r}+\gamma_{\bar{a}s})r_{\bar{a}}}(\delta_{ru}\delta_{sv}+\delta_{rv}\delta_{su})-\\
&- e^{\frac{2}{\sqrt{3}}\sum_{\bar{a}}(\gamma_{\bar{a}r}+\gamma_{\bar{a}u})r_{\bar{a}}}\delta_{rs}\delta_{uv}\Big](\tau,\vec{\sigma}).
\end{aligned}
\tag{A15}
$$

The tetrads and cotetrads adapted to the WSW hyper-surfaces of our gauge, given in Eq.(1) of Ref. [3], are

$$
\begin{aligned}
&{}^4_{(\Sigma)}\check{E}^\mu_{(\alpha)} = \{{}^4_{(\Sigma)}\check{E}^\mu_{(o)} = l^\mu = \hat{b}^\mu_l = \frac{1}{N}(b^\mu_\tau - N^r b^\mu_r);\ {}^4_{(\Sigma)}\check{E}^\mu_{(a)} = {}^3e^s_{(a)}b^\mu_s\},\\
&{}^4_{(\Sigma)}\check{E}^{(\alpha)}_\mu = \{{}^4_{(\Sigma)}\check{E}^{(o)}_\mu = \epsilon l_\mu = \hat{b}^l_\mu = N b^\tau_\mu;\ {}^4_{(\Sigma)}\check{E}^{(a)}_\mu = {}^3e^{(a)}_s\hat{b}^s_\mu\},
\end{aligned}
$$

$$
{}^4_{(\Sigma)}\check{E}^\mu_{(\alpha)}\quad {}^4g_{\mu\nu}\ {}^4_{(\Sigma)}\check{E}^\nu_{(\beta)} = {}^4\eta_{(\alpha)(\beta)},
$$

$$
{}^4_{(\Sigma)}\check{\tilde{E}}^A_{(\alpha)} = {}^4_{(\Sigma)}\check{E}^\mu_{(\alpha)}\,b^A_\mu,\quad \Rightarrow {}^4_{(\Sigma)}\check{\tilde{E}}^A_{(o)} = \epsilon l^A,
$$

$$\begin{aligned}
&{}^4_{(\Sigma)}\check{\tilde{E}}^{\tau}_{(o)} = \frac{1}{N}, \qquad {}^4_{(\Sigma)}\check{\tilde{E}}^{\tau}_{(a)} = 0,\\
&{}^4_{(\Sigma)}\check{\tilde{E}}^{r}_{(o)} = -\frac{N^r}{N}, \qquad {}^4_{(\Sigma)}\check{\tilde{E}}^{r}_{(a)} = {}^3e^r_{(a)};\\
{}^4_{(\Sigma)}\check{\tilde{E}}^{(\alpha)}_{A} = {}&{}^4_{(\Sigma)}\check{E}^{(\alpha)}_{\mu}\, b^{\mu}_{A}, \quad \Rightarrow {}^4_{(\Sigma)}\check{\tilde{E}}^{(o)}_{A} = l_A,\\
&\\
&{}^4_{(\Sigma)}\check{\tilde{E}}^{(o)}_{\tau} = N, \qquad {}^4_{(\Sigma)}\check{\tilde{E}}^{(a)}_{\tau} = N^r\, {}^3e^{(a)}_r = N^{(a)},\\
&{}^4_{(\Sigma)}\check{\tilde{E}}^{(o)}_{r} = 0, \qquad {}^4_{(\Sigma)}\check{\tilde{E}}^{(a)}_{r} = {}^3e^{(a)}_r,\\
&\\
&{}^4_{(\Sigma)}\check{E}^A_{(\alpha)}\, {}^4g_{AB}\, {}^4_{(\Sigma)}\check{E}^B_{(\beta)} = {}^4\eta_{(\alpha)(\beta)}.
\end{aligned} \tag{A16}$$

The 3-metric on our WSW hyper-surfaces and the line element of our space-time, given by Eqs.(183) of Ref. [3], are

$$\begin{aligned}
&{}^3\hat{g}_{rs} = e^{2q}\begin{pmatrix} e^{\frac{2}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a1}r_{\bar a}} & 0 & 0\\ 0 & e^{\frac{2}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a2}r_{\bar a}} & 0\\ 0 & 0 & e^{\frac{2}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a3}r_{\bar a}}\end{pmatrix} = \phi^4\, {}^3\hat{g}^{diag}_{rs},\\
&\quad \hat{\gamma} = {}^3\hat{g} = {}^3\hat{e}^2 = e^{6q} = \phi^{12}, \qquad det\, |\hat{g}^{diag}_{rs}| = 1,\\
&\\
&\quad d\hat{s}^2 = \epsilon\Big([-\epsilon + n]^2 (d\tau)^2 -\\
&\qquad -\delta_{uv}[\phi^2 e^{\frac{2}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a u}r_{\bar a}} d\sigma^u + \phi^{-2} e^{-\frac{2}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a u}r_{\bar a}}\, n_u\, d\tau]\\
&\qquad [\phi^2 e^{\frac{2}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a v}r_{\bar a}} d\sigma^v + \phi^{-2} e^{-\frac{2}{\sqrt{3}}\sum_{\bar a}\gamma_{\bar a v}r_{\bar a}}\, n_v\, d\tau]\Big),\\
&\\
&\quad q = 2 ln\, \phi = \frac{1}{6}\, ln\, {}^3\hat{g}, \qquad r_{\bar a} = \frac{\sqrt{3}}{2}\sum_r \gamma_{\bar a r}\, ln\, \frac{{}^3\hat{g}_{rr}}{{}^3\hat{g}}.
\end{aligned} \tag{A17}$$

The Frauendiener equations for the WSW triads, given in Eq.(200) of Ref. [3], are

$$\begin{aligned}
&{}^3\nabla_r\, {}^3e^{(WSW)r}_{(1)} = {}^3\nabla_r\, {}^3e^{(WSW)r}_{(2)} = 0,\\
&{}^3\nabla_r\, {}^3e^{(WSW)r}_{(3)} = -\alpha {}^3K,\\
&{}^3e^{(WSW)r}_{(1)}\, {}^3e^{(WSW)s}_{(3)}\, {}^3\nabla_r\, {}^3e^{(WSW)}_{(2)s} + {}^3e^{(WSW)r}_{(3)}\, {}^3e^{(WSW)s}_{(2)}\, {}^3\nabla_r\, {}^3e^{(WSW)}_{(1)s} +\\
+&{}^3e^{(WSW)r}_{(2)}\, {}^3e^{(WSW)s}_{(1)}\, {}^3\nabla_r\, {}^3e^{(WSW)}_{(3)s} = 0.
\end{aligned} \tag{A18}$$

The WSW tetrads of Eq.(201) of Ref. [3] are

$$\begin{aligned}
&{}^4_{(\Sigma)}\check{\tilde{E}}^{(WSW)}{}^{A}_{(o)} = \frac{1}{-\epsilon + n}(1; -n^r),\\
&{}^4_{(\Sigma)}\check{\tilde{E}}^{(WSW)}{}^{A}_{(a)} = (0; {}^3e^{(WSW)r}{}_{(a)}),
\end{aligned}$$

$$
\begin{aligned}
{}^4_{(\Sigma)}\check{E}^{(WSW)\mu}{}_{(o)} &= l^\mu,\\
{}^4_{(\Sigma)}\check{E}^{\mu}_{(a)} &= b^\mu_s \;\; {}^3e^{(WSW)s}{}_{(a)}.
\end{aligned}
\tag{A19}
$$

The embedding of the WSW hyper-surfaces of oue gauge into the space-time, given in Eq.(208) of Ref. [3], are

$$
\begin{aligned}
z^\mu_{(WSW)}(\tau,\vec{\sigma}) &= \delta^\mu_{(\mu)} x^{(\mu)}_{(\infty)}(0) + l^\mu(\tau,\vec{\sigma})\tau + \epsilon^\mu_r(\tau,\vec{\sigma})\sigma^r =\\
&= x^\mu_{(\infty)}(0) + l^\mu(\tau,\vec{\sigma})\tau + b^\mu_s(\tau,\vec{\sigma})\,{}^3e^{(WSW)s}{}_{(a)}(\tau,\vec{\sigma})\delta_{(a)r}\sigma^r =\\
&= x^\mu_{(\infty)}(0) + b^\mu_A(\tau,\vec{\sigma})F^A(\tau,\vec{\sigma}),\\
&\\
F^\tau(\tau,\vec{\sigma}) &= \frac{\tau}{-\epsilon + n(\tau,\vec{\sigma})},\\
F^s(\tau,\vec{\sigma}) &= {}^3e^{(WSW)s}{}_{(a)}(\tau,\vec{\sigma})\delta_{(a)r}\sigma^r - \frac{n^s(\tau,\vec{\sigma})}{-\epsilon + n(\tau,\vec{\sigma})}\tau,
\end{aligned}
\tag{A20}
$$

with the transition coefficients b^μ_A determined by Eqs.(209) of Ref. [3]

$$
\begin{aligned}
b^\mu_A &= \frac{\partial z^\mu_{(WSW)}}{\partial\sigma^A} = b^\mu_B\frac{\partial F^B}{\partial\sigma^A} + \frac{\partial b^\mu_B}{\partial\sigma^A}F^B,\\
A_A{}^B &= \delta^B_A - \frac{\partial F^B}{\partial\sigma^A},\\
F^B\frac{\partial b^\mu_B}{\partial\sigma^A} &= A_A{}^B b^\mu_B,\\
&or \quad b^\mu_b = (A^{-1})_B{}^a F^C\frac{\partial b^\mu_C}{\partial\sigma^A}.
\end{aligned}
\tag{A21}
$$

APPENDIX B: THE KERNELS $F^{(O)R}_{(A)(B)}$.

As shown in Ref. [3] the kernels $F^{(o)r}_{(a)(b)}$'s, zero curvature limit of the kernels $F^{r}_{(a)(b)}$'s of Eq.(1.4), are given by the equations

$$
\begin{aligned}
F^{(o)s}_{(a)(b)}(\vec{\sigma},\vec{\sigma}_1) &= \frac{1}{2}\Big[\epsilon_{(a)(s)(b)}\delta^3(\vec{\sigma},\vec{\sigma}_1) - \sum_{u,r}\epsilon_{(u)(r)(b)}\frac{\partial G^{(o)su}_{(a)}(\vec{\sigma},\vec{\sigma}_1)}{\partial \sigma_1^r}\Big],\\
F^{(o)r}_{(a)(1)}(\vec{\sigma}_1,\vec{\sigma};\tau) &= \frac{\partial G^{(o)r3}_{(a)}(\vec{\sigma}_1,\vec{\sigma};\tau)}{\partial \sigma_1^2} - \delta_{(a)3}\delta^r_2\delta^3(\vec{\sigma},\vec{\sigma}_1) =\\
&= \delta_{(a)2}\delta^r_3\delta^3(\vec{\sigma},\vec{\sigma}_1) - \frac{\partial G^{(o)r2}_{(a)}(\vec{\sigma}_1,\vec{\sigma};\tau)}{\partial \sigma_1^3},\\
F^{(o)r}_{(a)(2)}(\vec{\sigma}_1,\vec{\sigma};\tau) &= \frac{\partial G^{(o)r1}_{(a)}(\vec{\sigma}_1,\vec{\sigma};\tau)}{\partial \sigma_1^3} - \delta_{(a)1}\delta^r_3\delta^3(\vec{\sigma},\vec{\sigma}_1) =\\
&= \delta_{(a)3}\delta^r_1\delta^3(\vec{\sigma},\vec{\sigma}_1) - \frac{\partial G^{(o)r3}_{(a)}(\vec{\sigma}_1,\vec{\sigma};\tau)}{\partial \sigma_1^1},\\
F^{(o)r}_{(a)(3)}(\vec{\sigma}_1,\vec{\sigma};\tau) &= \frac{\partial G^{(o)r2}_{(a)}(\vec{\sigma}_1,\vec{\sigma};\tau)}{\partial \sigma_1^1} - \delta_{(a)2}\delta^r_1\delta^3(\vec{\sigma},\vec{\sigma}_1) =\\
&= \delta_{(a)1}\delta^r_2\delta^3(\vec{\sigma},\vec{\sigma}_1) - \frac{\partial G^{(o)r1}_{(a)}(\vec{\sigma}_1,\vec{\sigma};\tau)}{\partial \sigma_1^2},
\end{aligned}
\tag{B1}
$$

so that from Eqs.(2.12) we get

$$
F^{(o)a}_{(a)(b)}(\vec{\sigma},\vec{\sigma}_1) = 0, \qquad a,b=1,2,3,
$$

$$
\begin{aligned}
F^{(o)3}_{(2)(1)}(\vec{\sigma},\vec{\sigma}_1) &= -F^{(o)2}_{(3)(1)}(\vec{\sigma},\vec{\sigma}_1) = F^{(o)1}_{(3)(2)}(\vec{\sigma},\vec{\sigma}_1) = -F^{(o)3}_{(1)(2)}(\vec{\sigma},\vec{\sigma}_1) =\\
&= F^{(o)2}_{(1)(3)}(\vec{\sigma},\vec{\sigma}_1) = -F^{(o)1}_{(2)(3)}(\vec{\sigma},\vec{\sigma}_1) = \frac{1}{2}\delta^3(\vec{\sigma},\vec{\sigma}_1),
\end{aligned}
$$

$$
\begin{aligned}
F^{(o)2}_{(1)(1)}(\vec{\sigma},\vec{\sigma}_1) &= F^{(o)1}_{(2)(1)}(\vec{\sigma},\vec{\sigma}_1) =\\
&= -\frac{1}{2}\frac{\partial}{\partial \sigma_1^3}\int_{-\infty}^{\sigma_1^1} dw_1^1\delta^3(\vec{\sigma},w_1^1\sigma_1^2\sigma_1^3) =\\
&= -\frac{1}{2}\theta(\sigma_1^1,\sigma^1)\delta(\sigma^2,\sigma_1^2)\frac{\partial\delta(\sigma^3,\sigma_1^3)}{\partial \sigma_1^3},\\
F^{(o)3}_{(2)(3)}(\vec{\sigma},\vec{\sigma}_1) &= F^{(o)2}_{(3)(3)}(\vec{\sigma},\vec{\sigma}_1) =\\
&= \frac{1}{2}\frac{\partial}{\partial \sigma_1^1}\int_{-\infty}^{\sigma_1^3} dw_1^3\delta^3(\vec{\sigma},\sigma_1^1\sigma_1^2w_1^3) =\\
&= \frac{1}{2}\frac{\partial\delta(\sigma^1,\sigma_1^1)}{\partial \sigma_1^1}\delta(\sigma^2,\sigma_1^2)\theta(\sigma_1^3,\sigma^3),\\
F^{(o)3}_{(1)(1)}(\vec{\sigma},\vec{\sigma}_1) &= F^{(o)1}_{(3)(1)}(\vec{\sigma},\vec{\sigma}_1) =
\end{aligned}
$$

$$
\begin{aligned}
&= \frac{1}{2}\frac{\partial}{\partial \sigma_1^2}\int_{-\infty}^{\sigma_1^1} dw_1^1 \delta^3(\vec{\sigma}, w_1^1\sigma_1^2\sigma_1^3) = \\
&= \frac{1}{2}\theta(\sigma_1^1,\sigma^1)\frac{\partial\delta(\sigma^2,\sigma_1^2)}{\partial\sigma_1^2}\delta(\sigma^3,\sigma_1^3), \\
F^{(o)3}_{(2)(2)}(\vec{\sigma},\vec{\sigma}_1) &= F^{(o)2}_{(3)(2)}(\vec{\sigma},\vec{\sigma}_1) = \\
&= -\frac{1}{2}\frac{\partial}{\partial \sigma_1^1}\int_{-\infty}^{\sigma_1^2} dw_1^2 \delta^3(\vec{\sigma}, \sigma_1^1 w_1^2\sigma_1^3) = \\
&= -\frac{1}{2}\frac{\partial\delta(\sigma^1,\sigma_1^1)}{\partial\sigma_1^1}\theta(\sigma_1^2,\sigma^2)\delta(\sigma^3,\sigma_1^3), \\
F^{(o)2}_{(1)(2)}(\vec{\sigma},\vec{\sigma}_1) &= F^{(o)1}_{(2)(2)}(\vec{\sigma},\vec{\sigma}_1) = \\
&= \frac{1}{2}\frac{\partial}{\partial \sigma_1^3}\int_{-\infty}^{\sigma_1^2} dw_1^2 \delta^3(\vec{\sigma}, \sigma_1^1 w_1^2\sigma_1^3) = \\
&= \frac{1}{2}\delta(\sigma^1,\sigma_1^1)\theta(\sigma_1^2,\sigma^2)\frac{\partial\delta(\sigma^3,\sigma_1^3)}{\partial\sigma_1^3}, \\
F^{(o)3}_{(1)(3)}(\vec{\sigma},\vec{\sigma}_1) &= F^{(o)1}_{(3)(3)}(\vec{\sigma},\vec{\sigma}_1) = \\
&= -\frac{1}{2}\frac{\partial}{\partial \sigma_1^2}\int_{-\infty}^{\sigma_1^3} dw_1^3 \delta^3(\vec{\sigma}, \sigma_1^1\sigma_1^2 w_1^3) = \\
&= -\frac{1}{2}\delta(\sigma^1,\sigma_1^1)\frac{\partial\delta(\sigma^2,\sigma_1^2)}{\partial\sigma_1^2}\theta(\sigma_1^3,\sigma^3). \qquad \text{(B2)}
\end{aligned}
$$

APPENDIX C: FOURIER TRANSFORMS.

By assuming the validity of the Fourier transform on the linearized WSW CMC-hypersurfaces, which are assimilated to flat R^3 surfaces at the lowest level, for the real functions $r_{\bar a}(\tau,\vec\sigma)$, $\pi_{\bar a}(\tau,\vec\sigma)$ we have

$$
\begin{aligned}
r_{\bar a}(\tau,\vec\sigma) &= \int \frac{d^3k}{(2\pi)^3}\tilde r_{\bar a}(\tau,\vec k)e^{i\vec k\cdot\vec\sigma}, \qquad \tilde r^*_{\bar a}(\tau,\vec k)=\tilde r_{\bar a}(\tau,-\vec k),\\
\pi_{\bar a}(\tau,\vec\sigma) &= \int \frac{d^3k}{(2\pi)^3}\tilde \pi_{\bar a}(\tau,\vec k)e^{i\vec k\cdot\vec\sigma}, \qquad \tilde \pi^*_{\bar a}(\tau,\vec k)=\tilde \pi_{\bar a}(\tau,-\vec k),\\
&\tilde r_{\bar a}(\tau,\vec k)=\int d^3\sigma r_{\bar a}(\tau,\vec\sigma)e^{-i\vec k\cdot\vec\sigma}, \qquad \tilde \pi_{\bar a}(\tau,\vec k)=\int d^3\sigma \pi_{\bar a}(\tau,\vec\sigma)e^{-i\vec k\cdot\vec\sigma},
\end{aligned}
$$

$$
\begin{aligned}
\{\tilde r_{\bar a}(\tau,\vec k),\tilde\pi_{\bar b}(\tau,\vec k_1)\} &= \{\int d^3\sigma r_{\bar a}(\tau,\vec\sigma)e^{-i\vec k\cdot\vec\sigma},\int d^3\sigma_1\pi_{\bar b}(\tau,\vec\sigma_1)e^{-i\vec k_1\cdot\vec\sigma_1}\}\\
&= \delta_{\bar a\bar b}\int d^3\sigma_1\int d^3\sigma e^{-i\vec k_1\cdot\vec\sigma_1}e^{-i\vec k\cdot\vec\sigma}\delta(\vec\sigma-\vec\sigma_1)\\
&= \delta_{\bar a\bar b}\int d^3\sigma e^{-i(\vec k+\vec k_1)\cdot\vec\sigma}=(2\pi)^3\delta_{\bar a\bar b}\delta(\vec k+\vec k_1). \qquad \text{(C1)}
\end{aligned}
$$

The conditions (2.15) and (2.16) become

$$
\begin{aligned}
&\tilde\pi_{\bar a}(\tau,0k^2k^3)=\tilde\pi_{\bar a}(\tau,k^10k^3)=\tilde\pi_{\bar a}(\tau,k^1k^20)=0,\\
&\tilde r_{\bar a}(\tau,0k^2k^3)=\tilde r_{\bar a}(\tau,k^10k^3)=\tilde r_{\bar a}(\tau,k^1k^20)=0. \qquad \text{(C2)}
\end{aligned}
$$

Some useful relations are

$$
\begin{aligned}
&\theta(x)=lim_{\epsilon\rightarrow 0}\int\frac{dk}{2\pi i}\frac{e^{ikx}}{k-i\epsilon}, \quad \int\frac{d^3k}{(2\pi)^3}\frac{e^{i\vec k\cdot\vec\sigma}}{|\vec k|^2}=\frac{1}{4\pi|\vec\sigma|},\\
&\psi(x)=\int dy\theta(x-y)f(y) \quad \Rightarrow \quad \tilde f(k)=ik\tilde\psi(k),\\
&\Box r_{\bar a}(\tau,\vec\sigma)=\int\frac{d^3k}{(2\pi)^3}[\partial_\tau^2+|\vec k|^2]\tilde r_{\bar a}(\tau,\vec k)e^{i\vec k\cdot\vec\sigma},\\
&\int d^3\sigma\, e^{-i\vec k\cdot\vec\sigma}\int d^3\sigma_1\frac{f(\vec\sigma_1)}{4\pi|\vec\sigma-\vec\sigma_1|}=\frac{\tilde f(\vec k)}{|\vec k|^2},\\
&\int d^3\sigma_1\frac{r_{\bar a}(\tau,\vec\sigma_1)}{4\pi|\vec\sigma-\vec\sigma_1|}=\int\frac{d^3k}{(2\pi)^3}\frac{\tilde r_{\bar a}(\tau,\vec k)}{|\vec k|^2}e^{i\vec k\cdot\vec\sigma},\\
&\int d\sigma_1^u\theta(\sigma^u,\sigma_1^u)\int d\sigma_2^u\theta(\sigma_2^u,\sigma_1^u)e^{ik^u\sigma_2^u}=\frac{e^{ik^u\sigma^u}}{k_u^2}, \quad \int d\sigma_1^r\theta(\sigma_1^r,\sigma^u)e^{ik^r\sigma_1^r}=-\frac{e^{ik^r\sigma^r}}{ik_r},
\end{aligned}
$$

since

$$\frac{\partial^2}{(\partial\sigma^u)^2}\Big(\frac{e^{-ik^u\sigma^u}}{k_u^2}\Big) = e^{ik^u\sigma^u} = \frac{\partial^2}{(\partial\sigma^u)^2}\int_{-\infty}^{\sigma^u} d\sigma_1^u \int_{\sigma_1^u}^{\infty} d\sigma_2^u\, e^{ik^u\sigma_2^u},$$

$$\int_{-\infty}^{\sigma^u} d\sigma_1^u \int_{-\infty}^{\sigma_1^u} d\sigma_2^u \int_{\sigma^v}^{\infty} d\sigma_1^v \int_{\sigma_1^v}^{\infty} d\sigma_2^v \int d^3\sigma_3\, \frac{\partial^2_{3r}\partial^2_{3s}\partial^2_{3w}\partial^2_{3t} r_{\bar c}(\tau, \vec\sigma_3)}{4\pi|\vec\sigma_2 - \vec\sigma_3|}|_{\vec\sigma_2 = (\sigma_2^u\sigma_2^v\sigma^{k\neq u,v})} =$$
$$= \int \frac{d^3k}{(2\pi)^3}\, \frac{k_r^2k_s^2k_w^2k_t^2}{k_u^2k_v^2|\vec k|^2} \tilde r_{\bar c}(\tau, \vec k) e^{i\vec k\cdot\vec\sigma}. \tag{C3}$$

Let us consider Eqs.(5.2). Their Fourier transform is

$$\partial_\tau \tilde r_{\bar a}(\tau, \vec k) = \sum_{\bar b} A_{\bar a\bar b}(\vec k) \tilde\pi_{\bar b}(\tau, \vec k), \tag{C4}$$

where the matrix $A_{\bar a\bar b}(\vec k)$ is

$$A_{\bar a\bar b}(\vec k) = A_{\bar b\bar a}(\vec k) = A_{\bar a\bar b}(-\vec k) = \frac{24\pi G}{c^3}\Big[\delta_{\bar a\bar b} + \frac{1}{4}\sum_{r,s,u,v} \gamma_{\bar ar}\gamma_{\bar bs}$$
$$(1-\delta_{uv})[1 - 2(\delta_{ur}+\delta_{vr})][1 - 2(\delta_{us}+\delta_{vs})]\Big(\frac{k^rk^s}{k^uk^v}\Big)^2\Big],$$
$$det\,\Big(A_{\bar a\bar b}(\vec k)\Big) = \frac{192\pi^2G^2}{c^6}\frac{|\vec k|^6}{(k^1k^2k^3)^2}. \tag{C5}$$

Eq.(C2) implies that Eq.(C4) is well defined, even if the matrix $A_{\bar a\bar b}(\vec k)$ diverges for $k^r \rightarrow 0$, if $\tilde\pi_{\bar a}(\tau, \vec k)$ *vanishes for* $\vec k \rightarrow 0$ *at least as* $(k^1k^2k^3)^{2+\epsilon}$, $\epsilon > 0$.

For $\vec k \neq 0$ its inverse is

$$A^{-1}_{\bar a\bar b}(\vec k) = A^{-1}_{\bar b\bar a}(\vec k) = A^{-1}_{\bar a\bar b}(-\vec k) =$$
$$= \frac{1}{det\,A(\vec k)}\sum_{\bar c\bar d} \epsilon_{\bar a\bar c}\epsilon_{\bar b\bar d}A_{\bar c\bar d}(\vec k) = \frac{c^3}{8\pi G}\frac{1}{|\vec k|^6}\Big[(k^1k^2k^3)^2\delta_{\bar a\bar b} +$$
$$+ \frac{1}{2}\sum_{r,s,t,\bar c,\bar d} \epsilon_{\bar a\bar c}\gamma_{\bar cr}\epsilon_{\bar b\bar d}\gamma_{\bar ds}(2\delta_{tr}-1)(2\delta_{ts}-1)(k^rk^sk^t)^2\Big]. \tag{C6}$$

The Fourier transform of Eq.(5.5) is

$$\partial_\tau \tilde\pi_{\bar a}(\tau, \vec k) = \sum_{\bar b} B_{\bar a\bar b}(\vec k) \tilde r_{\bar b}(\tau, \vec k), \tag{C7}$$

where $B_{\bar a\bar b}(\vec k)$ is the matrix

$$
\begin{aligned}
B_{\bar a\bar b}(\vec k) &= B_{\bar b\bar a}(\vec k) = B_{\bar a\bar b}(-\vec k) =\\
&= \frac{c^3}{24\pi G}\Big[- \delta_{\bar a\bar b} \sum_r k_r^2 + \sum_{rs} \gamma_{\bar ar}\gamma_{\bar bs}\, \Big(2\delta_{rs} k_r^2 - \frac{k_r^2 k_s^2}{2|\vec k|^2}\Big)\Big],\\
&\qquad det\, \Big(B_{\bar a\bar b}(\vec k)\Big) = \frac{c^6\, (k^1 k^2 k^3)^2}{192\pi^2 G^2 |\vec k|^2},
\end{aligned}
\tag{C8}
$$

satisfying

$$
\begin{aligned}
&\sum_{\bar b} A_{\bar a\bar b}(\vec k) B_{\bar b\bar c}(\vec k) = - |\vec k|^2 \delta_{\bar a\bar c} = \sum_{\bar b} B_{\bar a\bar b}(\vec k) A_{\bar b\bar c}(\vec k),\\
&\qquad \Rightarrow \qquad A^{-1}(\vec k) = - |\vec k|^2\, B(\vec k).
\end{aligned}
\tag{C9}
$$

But this implies that the Fourier transform of Eq.(5.7) reduces to the wave equation

$$
\begin{aligned}
\ddot{\tilde r}_{\bar a}(\tau, \vec k) &\overset{\circ}{=} \sum_{\bar b} \Big(- |\vec k|^2 \delta_{\bar a\bar b} + 2 \sum_r \gamma_{\bar ar}\gamma_{\bar br}\, k_r^2 - \frac{1}{2} \sum_{rs} \gamma_{\bar ar} \frac{k_r^2 k_s^2}{|\vec k|^2}\Big) \tilde r_{\bar b}(\tau, \vec k) +\\
&+ \frac{1}{8} \sum_{\bar c} \Big(\sum_{rs} \sum_{tw} \sum_{uv} \Big[- 2\gamma_{\bar ar}\gamma_{\bar cs} + 4\gamma_{\bar ar}\gamma_{\bar ct} (\delta_{ts} - \frac{1}{3}) - \gamma_{\bar ar}\gamma_{\bar cw} (\delta_{ts} - \frac{1}{3})\Big] (1 - \delta_{uv})\\
&\quad [1 - 2(\delta_{ur} + \delta_{vr})][1 - 2(\delta_{us} + \delta_{vs})] \frac{k_t^2 k_w^2 k_r^2 k_s^2}{k_u^2 k_v^2 |\vec k|^2}\Big) \tilde r_{\bar c}(\tau, \vec k) =\\
&= - |\vec k|^2 \tilde r_{\bar a}(\tau, \vec k),
\end{aligned}
\tag{C10}
$$

whose solutions

$$
\begin{aligned}
r_{\bar a}(\tau, \vec \sigma) &= \int \frac{d^3k}{(2\pi)^3} \Big(C_{\bar a}(\vec k) e^{-i(|\vec k|\, \tau - \vec k \cdot \vec \sigma)} + C^*_{\bar a}(\vec k) e^{i(|\vec k|\tau - \vec k\cdot \vec \sigma)}\Big),\\
&\qquad \tilde r_{\bar a}(\tau, \vec k) = C_{\bar a}(\vec k)\, e^{-i\, |\vec k|\, \tau} + C^*_{\bar a}(-\vec k)\, e^{i\, |\vec k|\, \tau},
\end{aligned}
\tag{C11}
$$

have the arbitrary functions $C_{\bar a}(\vec k)$ satisfying Eq.(C2). Then Eqs.(5.4) imply the following form of the momenta restricted to the solutions

$$
\begin{aligned}
\pi_{\bar a}(\tau, \vec \sigma) &= - i \int \frac{d^3k}{(2\pi)^3} \sum_{\bar b} A^{-1}_{\bar a\bar b}(\vec k)\, |\vec k|\, \Big[C_{\bar b}(\vec k)\, e^{-i(|\vec k|\, \tau - \vec k\cdot \vec \sigma)} - C^*_{\bar a}(\vec k) e^{i(|\vec k|\tau - \vec k\cdot \vec \sigma)}\Big],\\
\tilde \pi_{\bar a}(\tau, \vec k) &= \sum_{\bar b} A^{-1}_{\bar a\bar b}(\vec k)\, \partial_\tau \tilde r_{\bar b}(\tau, \vec k) =\\
&= \sum_{\bar b} A^{-1}_{\bar a\bar b}(\vec k)\, \Big[- i |\vec k| C_{\bar b}(\vec k) e^{-i|\vec k|\tau} + i |\vec k| C^*_{\bar b}(-\vec k) e^{i|\vec k|\tau}\Big].
\end{aligned}
\tag{C12}
$$

To have $\tilde \pi_{\bar a}(\tau, \vec k)$ vanishing for $\vec k \rightarrow 0$ at least as $(k^1 k^2 k^3)^{2+\epsilon}$, $\epsilon > 0$, we must require that the functions $C_{\bar a}(\vec k)$ *vanish at least as* $(k^1\, k^2\, k^3)^\epsilon$ for $k^r \rightarrow 0$.

The Fourier transform of the ADM energy (3.1) is

$$
\begin{aligned}
\hat{E}_{ADM} &= \frac{12\pi G}{c^3} \int \frac{d^3k}{(2\pi)^3} \sum_{\bar{a}} \tilde{\pi}_{\bar{a}}(\tau, \vec{k}) \tilde{\pi}_{\bar{a}}(\tau, -\vec{k}) + \\
&+ \frac{c^3}{48\pi G} \int \frac{d^3k}{(2\pi)^3} \sum_{\bar{a}r} k_r^2 \tilde{r}_{\bar{a}}(\tau, \vec{k}) \tilde{r}_{\bar{a}}(\tau, -\vec{k}) + \\
&+ \frac{3\pi G}{c^3} \int \frac{d^3k}{(2\pi)^3} \sum_{\bar{a}\bar{b}} \Big[\sum_{r,s,u,v} \gamma_{\bar{a}r} \gamma_{\bar{b}s} (1 - \delta_{uv}) [1 - 2(\delta_{ur} + \delta_{vr})] \\
&\quad [1 - 2(\delta_{us} + \delta_{vs})] \Big(\frac{k^r k^s}{k^u k^v} \Big)^2 \tilde{\pi}_{\bar{a}}(\tau, \vec{k}) \tilde{\pi}_{\bar{b}}(\tau, -\vec{k}) \Big] - \\
&- \frac{c^3}{24\pi G} \int \frac{d^3k}{(2\pi)^3} \sum_{\bar{a}\bar{b}r} \gamma_{\bar{a}r} \gamma_{\bar{b}r} k_r^2 \tilde{r}_{\bar{a}}(\tau, \vec{k}) \tilde{r}_{\bar{b}}(\tau, -\vec{k}) + \\
&+ \frac{c^3}{96\pi G} \int \frac{d^3k}{(2\pi)^3} \sum_{\bar{a}\bar{b}ru} \gamma_{\bar{a}r} \gamma_{\bar{b}u} \frac{k_r^2 k_u^2}{|\vec{k}|^2} \tilde{r}_{\bar{a}}(\tau, \vec{k}) \tilde{r}_{\bar{b}}(\tau, -\vec{k}). \qquad \text{(C13)}
\end{aligned}
$$

As a check we can recover the Fourier transforms (C4) and (C7) of the Hamilton equations by using this form of the weak ADM energy and the Poisson brackets (C1).

The Fourier transform of the shift functions (3.5) is

$$
\tilde{n}_r(\tau, \vec{k}) = -i\sqrt{3} \frac{2\pi G}{c^3} \sum_{\bar{b}vuc} \gamma_{\bar{b}v} (1 - \delta_{uc}) [1 - 2(\delta_{uv} + \delta_{cv})] [1 - 2(\delta_{ur} + \delta_{cr})] \frac{k_r k_v^2}{k_u^2 k_c^2} \tilde{\pi}_{\bar{b}}(\tau, \vec{k}), \quad \text{(C14)}
$$

while the Fourier transform of the extrinsic curvature is

$$
{}^3\tilde{\hat{K}}_{rs}(\tau, \vec{k}) = \epsilon\sqrt{3} \frac{4\pi G}{c^3} \sum_{\bar{a}} \Big[2\delta_{rs} \gamma_{\bar{a}s} - (1 - \delta_{rs}) \sum_w \gamma_{\bar{a}w} [1 - 2(\delta_{rw} + \delta_{sw})] \frac{k_w^2}{k_r k_s} \Big] \tilde{\pi}_{\bar{a}}(\tau, \vec{k}), \quad \text{(C15)}
$$

The following Fourier transforms are used in the study of the geodesic deviation equation

$$
\begin{aligned}
{}^4\tilde{\hat{\Gamma}}^r_{\tau s}(\tau, \vec{k}) &= ik_s \tilde{n}_r(\tau, \vec{k}) + \epsilon\, {}^3\tilde{\hat{K}}_{rs}(\tau, \vec{k}), \\
{}^4\tilde{\hat{R}}^r{}_{\tau s \tau}(\tau, \vec{k}) &= -\epsilon \partial_\tau\, {}^3\tilde{\hat{K}}_{rs}(\tau, \vec{k}). \qquad \text{(C16)}
\end{aligned}
$$

The Fourier transforms of the exponent $q(\tau, \vec{\sigma})$ ($\sqrt{{}^4\hat{g}} \approx -\epsilon(1 + 3q)$) of the conformal factor of the 3-metric, given in Eq.(2.2), and of the inverse 3-metric (3.7) are

$$
\tilde{q}(\tau, \vec{k}) = \frac{1}{2\sqrt{3}} \sum_{\bar{a}u} \gamma_{\bar{a}u} \frac{k_u^2}{|\vec{k}|^2} \tilde{r}_{\bar{a}}(\tau, \vec{k}), \qquad \text{(C17)}
$$

and

$$
{}^3\tilde{\hat{\gamma}}^{rs}(\tau, \vec{k}) = \delta^{rs} \Big(1 - \frac{2}{\sqrt{3}} \sum_{\bar{a}} \Big[\gamma_{\bar{a}r} \tilde{r}_{\bar{d}}(\tau, \vec{k}) + \frac{1}{2} \sum_u \gamma_{\bar{a}u} \frac{k_u^2}{|\vec{k}|^2} \tilde{r}_{\bar{a}}(\tau, \vec{k}) \Big] \Big). \qquad \text{(C18)}
$$

APPENDIX D: LINEARIZED 3- AND 4-TENSORS.

1. Linearized 3-Tensors on the WSW Hyper-Surfaces Σ_τ

¿From Eqs.(3.8) and (A7) we get the following results for the linearized Riemannian structure of the WSW hyper-surfaces $\Sigma^{(WSW)}_\tau$.

The Christoffel symbols are

$$\begin{aligned}
{}^3\hat{\Gamma}^r_{uv}(\tau,\vec{\sigma}) &= \frac{1}{\sqrt{3}}\sum_{\bar{a}}\Big(-\delta_{uv}\Big[\gamma_{\bar{a}u}\partial_r r_{\bar{a}}(\tau,\vec{\sigma}) - \\
&- \frac{1}{2}\sum_w \gamma_{\bar{a}w}\partial_r \int d^3\sigma_1\, \frac{\partial^2_{1w} r_{\bar{a}}(\tau,\vec{\sigma}_1)}{4\pi|\vec{\sigma}-\vec{\sigma}_1|}\Big] + \\
&+ \delta_{ru}\Big[\gamma_{\bar{a}u}\partial_v r_{\bar{a}}(\tau,\vec{\sigma}) - \\
&- \frac{1}{2}\sum_w \gamma_{\bar{a}w}\partial_v \int d^3\sigma_1\, \frac{\partial^2_{1w} r_{\bar{a}}(\tau,\vec{\sigma}_1)}{4\pi|\vec{\sigma}-\vec{\sigma}_1|}\Big] + \\
&+ \delta_{rv}\Big[\gamma_{\bar{a}v}\partial_u r_{\bar{a}}(\tau,\vec{\sigma}) - \\
&- \frac{1}{2}\sum_w \gamma_{\bar{a}w}\partial_u \int d^3\sigma_1\, \frac{\partial^2_{1w} r_{\bar{a}}(\tau,\vec{\sigma}_1)}{4\pi|\vec{\sigma}-\vec{\sigma}_1|}\Big]\Big) + O(r^2_{\bar{a}}),
\end{aligned}$$

$$\begin{aligned}
\sum_v {}^3\hat{\Gamma}^v_{uv}(\tau,\vec{\sigma}) &= -\frac{\sqrt{3}}{2}\sum_{\bar{a}w}\gamma_{\bar{a}w}\partial_u \int d^3\sigma_1\, \frac{\partial^2_{1w} r_{\bar{a}}(\tau,\vec{\sigma}_1)}{4\pi|\vec{\sigma}-\vec{\sigma}_1|} + O(r^2_{\bar{a}}),\\
\sum_u \partial_{(s}\, {}^3\hat{\Gamma}^u_{r)u}(\tau,\vec{\sigma}) &= -\frac{\sqrt{3}}{2}\sum_{\bar{a}w}\gamma_{\bar{a}w}\partial_r\partial_s \int d^3\sigma_1\, \frac{\partial^2_{1w} r_{\bar{a}}(\tau,\vec{\sigma}_1)}{4\pi|\vec{\sigma}-\vec{\sigma}_1|} + O(r^2_{\bar{a}}),\\
\sum_r \partial_r\, {}^3\hat{\Gamma}^r_{uv}(\tau,\vec{\sigma}) &= \frac{1}{\sqrt{3}}\sum_{\bar{a}}\Big(\Big[-\delta_{uv}\Big(\gamma_{\bar{a}u}\triangle + \frac{1}{2}\sum_w \gamma_{\bar{a}w}\partial^2_w\Big) + (\gamma_{\bar{a}u}+\gamma_{\bar{a}v})\partial_u\partial_v\Big] r_{\bar{a}}(\tau,\vec{\sigma}) - \\
&- \sum_w \gamma_{\bar{a}w}\partial_u\partial_v \int d^3\sigma_1\, \frac{\partial^2_{1w} r_{\bar{a}}(\tau,\vec{\sigma}_1)}{4\pi|\vec{\sigma}-\vec{\sigma}_1|}\Big) + O(r^2_{\bar{a}}),\\
\sum_{ur}\partial_r\, {}^3\hat{\Gamma}^u_{ru}(\tau,\vec{\sigma}) &= \sum_{ru}\partial_r\, {}^3\hat{\Gamma}^r_{uu}(\tau,\vec{\sigma}) = \frac{\sqrt{3}}{2}\sum_w \gamma_{\bar{a}w}\partial^2_w r_{\bar{a}}(\tau,\vec{\sigma}) + O(r^2_{\bar{a}}),
\end{aligned} \tag{D1}$$

while the spin connection and the field strength are

$$\begin{aligned}
{}^3\hat{\omega}_{r(a)}(\tau,\vec{\sigma}) &= \frac{1}{\sqrt{3}}\sum_u \epsilon_{(a)(r)(u)}\sum_{\bar{a}}\Big[\gamma_{\bar{a}r}\partial_u r_{\bar{a}}(\tau,\vec{\sigma}) - \\
&- \frac{1}{2}\sum_v \gamma_{\bar{a}v}\frac{\partial}{\partial\sigma^u}\int d^3\sigma_1 \frac{\partial^2_{1u} r_{\bar{a}}(\tau,\vec{\sigma}_1)}{4\pi|\vec{\sigma}-\vec{\sigma}_1|}\Big] + O(r^2_{\bar{a}}),
\end{aligned} \tag{D2}$$

$$\begin{aligned}
{}^3\hat{\Omega}_{rs(a)}(\tau,\vec{\sigma}) &= \partial_r\, {}^3\hat{\omega}_{s(a)}(\tau,\vec{\sigma}) - \partial_s\, {}^3\hat{\omega}_{r(a)}(\tau,\vec{\sigma}) + O(r^2_{\bar{a}}) = \\
&= \frac{1}{\sqrt{3}}\sum_{\bar{a}u}\Big(\epsilon_{(a)(s)(u)}\Big[\gamma_{\bar{a}s}\partial_u\partial_r r_{\bar{a}}(\tau,\vec{\sigma}) -
\end{aligned}$$

$$
\begin{aligned}
&- \frac{1}{2} \sum_v \gamma_{\bar{a}v} \partial_u \partial_r \int d^3\sigma_1 \, \frac{\partial^2_{1v} r_{\bar{a}}(\tau, \vec{\sigma}_1)}{4\pi |\vec{\sigma} - \vec{\sigma}_1|} \Big] - \\
&- \epsilon_{(a)(r)(u)} \Big[\gamma_{\bar{a}r} \partial_u \partial_s r_{\bar{a}}(\tau, \vec{\sigma}) - \\
&- \frac{1}{2} \sum_v \gamma_{\bar{a}v} \partial_u \partial_s \int d^3\sigma_1 \, \frac{\partial^2_{1v} r_{\bar{a}}(\tau, \vec{\sigma}_1)}{4\pi |\vec{\sigma} - \vec{\sigma}_1|} \Big] \Big) + O(r^2_{\bar{a}}).
\end{aligned}
\qquad \text{(D3)}
$$

Finally the Riemann and Ricci tensors and the curvature scalars are

$$
\begin{aligned}
{}^3\hat{R}_{rsuv}(\tau, \vec{\sigma}) &= \epsilon_{(r)(s)(a)} \, {}^3\hat{\Omega}_{uv(a)}(\tau, \vec{\sigma}) + O(r^2_{\bar{a}}) = \\
&= \frac{1}{\sqrt{3}} \sum_{\bar{a}} \Big(\Big[\gamma_{\bar{a}r} (\delta_{rv} \partial_s \partial_u - \delta_{ru} \partial_s \partial_v) - \gamma_{\bar{a}s} (\delta_{sv} \partial_r \partial_u - \delta_{su} \partial_r \partial_v) \Big] r_{\bar{a}}(\tau, \vec{\sigma}) - \\
&- \frac{1}{2} \sum_t \gamma_{\bar{a}t} \Big[\delta_{rv} \partial_s \partial_u - \delta_{ru} \partial_s \partial_v - (\delta_{sv} \partial_r \partial_u - \delta_{su} \partial_r \partial_v) \Big] \\
&\int d^3\sigma_1 \, \frac{\partial^2_{1t} r_{\bar{a}}(\tau, \vec{\sigma}_1)}{4\pi |\vec{\sigma} - \vec{\sigma}_1|} \Big) + O(r^2_{\bar{a}}), \\
{}^3\hat{R}_{rs}(\tau, \vec{\sigma}) &= \sum_u {}^3\hat{R}_{urus}(\tau, \vec{\sigma}) + O(r^2_{\bar{a}}) = \\
&= \frac{1}{\sqrt{3}} \sum_{\bar{a}} \Big(\Big[(\gamma_{\bar{a}r} + \gamma_{\bar{a}s}) \partial_r \partial_s - \delta_{rs} (\gamma_{\bar{a}r} \triangle + \frac{1}{2} \sum_t \gamma_{\bar{a}t} \partial^2_t) \Big] r_{\bar{a}}(\tau, \vec{\sigma}) - \\
&- \frac{1}{2} \int d^3\sigma_1 \, \sum_t \gamma_{\bar{a}t} \partial_r \partial_s d^3\sigma_1 \, \frac{\partial^2_{1t} r_{\bar{a}}(\tau, \vec{\sigma}_1)}{4\pi |\vec{\sigma} - \vec{\sigma}_1|} \Big) + O(r^2_{\bar{a}}), \\
{}^3\hat{R}(\tau, \vec{\sigma}) &= \sum_r {}^3\hat{R}_{rr}(\tau, \vec{\sigma}) + O(r^2_{\bar{a}}) = \frac{1}{\sqrt{3}} \sum_{\bar{a}v} \partial^2_v r_{\bar{a}}(\tau, \vec{\sigma}) + O(r^2_{\bar{a}}).
\end{aligned}
\qquad \text{(D4)}
$$

2. The 4-Christoffel Symbols, the 4-Riemann Tensor and Einstein Equations.

In this Subsection we will give the linearized form of the main 4-tensors of our space-time.

By using the parametrization of the 4-metric ${}^4g_{AB}$ given in footnote 6 and Eqs.(4.1), with the ${}^3\hat{\Gamma}^u_{rs}$'s of Eq.(D1), we get the following 4-Christoffel symbols $[N = -\epsilon + n,\ N^r = n_r + O(r^2_{\bar{a}})]$

$$
\begin{aligned}
{}^4\hat{\Gamma}^\tau_{\tau\tau}(\tau, \vec{\sigma}) &= \Big(\frac{1}{N} [\partial_\tau N + N^r \partial_r N - N^r N^s \, {}^3\hat{K}_{rs}] \Big)(\tau, \vec{\sigma}) = \\
&= 0 + O(r^2_{\bar{a}}),
\end{aligned}
$$

$$
{}^4\hat{\Gamma}^\tau_{r\tau}(\tau, \vec{\sigma}) = \Big(\frac{1}{N} [\partial_r N - {}^3\hat{K}_{rs} N^s] \Big)(\tau, \vec{\sigma}) = 0 + O(r^2_{\bar{a}}),
$$

$$
\begin{aligned}
{}^4\hat{\Gamma}^\tau_{rs}(\tau, \vec{\sigma}) &= -\Big(\frac{1}{N} \, {}^3\hat{K}_{rs} \Big)(\tau, \vec{\sigma}) = \\
&= \frac{4\pi G}{c^3} {}^3G_{o(a)(b)(c)(d)} \delta_{(a)r} \delta_{(b)s} \sum_u \delta_{(c)u} \, {}^3\check{\tilde{\pi}}^u_{(d)}(\tau, \vec{\sigma}) + O(r^2_{\bar{a}}) =
\end{aligned}
$$

$$
\begin{aligned}
&= \sqrt{3}\frac{4\pi G}{c^3}\sum_{\bar{a}}\Big[2\delta_{rs}\gamma_{\bar{a}r}\pi_{\bar{a}}(\tau,\vec{\sigma}) + \\
&+ [1-\delta_{rs}]\sum_w \gamma_{\bar{a}w}[1-2(\delta_{rw}+\delta_{sw})] \\
&\quad \frac{\partial^2}{\partial(\sigma^w)^2}\int_{\sigma^r}^{\infty} d\sigma_1^r \int_{\sigma^s}^{\infty} d\sigma_1^s\ \pi_{\bar{a}}(\tau,\sigma_1^r\sigma_1^s\sigma^{k\neq r,s})\Big] + O(r_{\bar{a}}^2),
\end{aligned}
$$

$$
\begin{aligned}
{}^4\hat{\Gamma}^u_{\tau\tau}(\tau,\vec{\sigma}) &= \Big(\partial_\tau N^u - \frac{N^u}{N}\partial_\tau N + ({}^3\hat{g}^{uv} - \frac{N^uN^v}{N^2})N\partial_v N + \\
&+ N^u{}_{|v}N^v - 2N({}^3\hat{g}^{uv} - \frac{N^uN^v}{N^2})^3\hat{K}_{vr}N^r\Big)(\tau,\vec{\sigma}) = \\
&= \partial_\tau n_u(\tau,\vec{\sigma}) + O(r_{\bar{a}}^2) = \\
&= \sqrt{3}\frac{2\pi G}{c^3}\sum_{\bar{a}t}\gamma_{\bar{a}t}\sum_{mn}(1-\delta_{mn})[1-2(\delta^u_m+\delta^u_n)][1-2(\delta_{tm}+\delta_{tn})] \\
&\quad \frac{\partial}{\partial\sigma^u}\int_{-\infty}^{\sigma^m} d\sigma_1^m \int_{-\infty}^{\sigma^n} d\sigma_1^n \int_{\sigma_1^m}^{\infty} d\sigma_2^m \int_{\sigma_1^n}^{\infty} d\sigma_2^n \frac{\partial^2\partial_\tau\pi_{\bar{a}}(\tau,\sigma_2^m\sigma_2^n\sigma_2^{k\neq m,n})}{(\partial\sigma_2^t)^2}|_{\sigma_2^k=\sigma^k} + O(r_{\bar{a}}^2),
\end{aligned}
$$

$$
\begin{aligned}
{}^4\hat{\Gamma}^u_{r\tau}(\tau,\vec{\sigma}) &= \Big(N^u{}_{|r} - \frac{N^u}{N}\partial_r N - N({}^3\hat{g}^{uv} - \frac{N^uN^v}{N^2})^3\hat{K}_{vr}\Big)(\tau,\vec{\sigma}) = \\
&= \partial_r n_u(\tau,\vec{\sigma}) + \epsilon\,{}^3\hat{K}_{ur}(\tau,\vec{\sigma}) + O(r_{\bar{a}}^2) = \\
&= \partial_r n_u(\tau,\vec{\sigma}) + \frac{4\pi G}{c^3}{}^3G_{o(a)(b)(c)(d)}\delta^u_{(}a)\delta_{(b)r}\sum_w \delta_{(c)w}{}^3\check{\tilde{\pi}}^w_{(d)}(\tau,\vec{\sigma}) + O(r_{\bar{a}}^2) = \\
&= \sqrt{3}\frac{4\pi G}{c^3}\sum_{\bar{a}}\Big[2\delta^u_r\gamma_{\bar{a}r}\pi_{\bar{a}}(\tau,\vec{\sigma}) + [1-\delta^u_r]\sum_t\gamma_{\bar{a}t}[1-2(\delta^u_t+\delta_{rt})] \\
&\quad \frac{\partial^2}{\partial(\sigma^t)^2}\int_{\sigma^u}^{\infty} d\sigma_1^u \int_{\sigma^r}^{\infty} d\sigma_1^r\ \pi_{\bar{a}}(\tau,\sigma_1^u\sigma_1^r\sigma^{k\neq u,r}) + \\
&+ \frac{1}{2}\sum_{\bar{a}t}\gamma_{\bar{a}t}\sum_{mn}(1-\delta_{mn})[1-2(\delta^u_m+\delta^u_n)][1-2(\delta_{tm}+\delta_{tn})] \\
&\quad \frac{\partial^2}{\partial\sigma^r\partial\sigma^u}\int_{-\infty}^{\sigma^m} d\sigma_1^m \int_{-\infty}^{\sigma^n} d\sigma_1^n \int_{\sigma_1^m}^{\infty} d\sigma_2^m \int_{\sigma_1^n}^{\infty} d\sigma_2^n \frac{\partial^2\pi_{\bar{a}}(\tau,\sigma_2^m\sigma_2^n\sigma_2^{k\neq m,n})}{(\partial\sigma_2^t)^2}|_{\sigma_2^k=\sigma^k}\Big] + \\
&+ O(r_{\bar{a}}^2),
\end{aligned}
$$

$$
\begin{aligned}
{}^4\hat{\Gamma}^u_{rs}(\tau,\vec{\sigma}) &= \Big({}^3\hat{\Gamma}^u_{rs} + \frac{N^u}{N}\,{}^3\hat{K}_{rs}\Big)(\tau,\vec{\sigma}) = \\
&= {}^3\hat{\Gamma}^u_{rs}(\tau,\vec{\sigma}) + O(r_{\bar{a}}^2) = \\
&= \frac{1}{\sqrt{3}}\sum_{\bar{a}}\Big(-\delta_{uv}\Big[\gamma_{\bar{a}u}\partial_r r_{\bar{a}}(\tau,\vec{\sigma}) - \frac{1}{2}\sum_w\gamma_{\bar{a}w}\partial_r\int d^3\sigma_1\,\frac{\partial^2_{1w}r_{\bar{a}}(\tau,\vec{\sigma}_1)}{4\pi|\vec{\sigma}-\vec{\sigma}_1|}\Big] + \\
&+ \delta_{ru}\Big[\gamma_{\bar{a}u}\partial_v r_{\bar{a}}(\tau,\vec{\sigma}) - \frac{1}{2}\sum_w\gamma_{\bar{a}w}\partial_v\int d^3\sigma_1\,\frac{\partial^2_{1w}r_{\bar{a}}(\tau,\vec{\sigma}_1)}{4\pi|\vec{\sigma}-\vec{\sigma}_1|}\Big] + \\
&+ \delta_{rv}\Big[\gamma_{\bar{a}v}\partial_u r_{\bar{a}}(\tau,\vec{\sigma}) - \frac{1}{2}\sum_w\gamma_{\bar{a}w}\partial_u\int d^3\sigma_1\,\frac{\partial^2_{1w}r_{\bar{a}}(\tau,\vec{\sigma}_1)}{4\pi|\vec{\sigma}-\vec{\sigma}_1|}\Big]\Big) + O(r_{\bar{a}}^2). \qquad \text{(D5)}
\end{aligned}
$$

Since $\varphi_{(a)}(\tau,\vec{\sigma}) = 0$ in our gauge, the space-time spin connection ${}^4\omega_A{}^{(\alpha)}{}_{(\beta)} =$

${}^4E^{(\alpha)}_B\,[\partial_A\,{}^4E^B_{(\beta)} + {}^4\Gamma^B_{AC}\,{}^4E^C_{(\beta)}]$ can be evaluated with the formula

$$ {}^4\overset{\circ}{\omega}_A{}^{(\alpha)}{}_{(\beta)} = {}^4_{(\Sigma)}\check{\tilde{E}}^{(\alpha)}_B\,[\partial_A\,{}^4_{(\Sigma)}\check{\tilde{E}}^B_{(\beta)} + {}^4\Gamma^B_{AC}\,{}^4_{(\Sigma)}\check{\tilde{E}}^C_{(\beta)}], \tag{D6} $$

by using Eqs.(3.9) and (D5).

For the 4-Riemann tensor ${}^4R^A{}_{BCD} = \frac{1}{2}\,{}^4g^{AE}\,(\partial_B\partial_D\,{}^4g_{EC} + \partial_E\partial_C\,{}^4g_{BD} - \partial_E\partial_D\,{}^4g_{BC} - \partial_B\partial_C\,{}^4g_{ED}) + {}^4g^{AL}\,{}^4g_{EF}\,({}^4\Gamma^E_{LC}\,{}^4\Gamma^F_{BD} - {}^4\Gamma^E_{LD}\,{}^4\Gamma^F_{BC})$ we get

$$ {}^4\hat{R}^A{}_{BCD} = \partial_C\,{}^4\hat{\Gamma}^A_{BD} - \partial_D\,{}^4\hat{\Gamma}^A_{CB} + O(r^2_{\bar a}), $$

$$
\begin{aligned}
{}^4\hat{R}^\tau{}_{\tau CD}(\tau,\vec\sigma) &= 0 + O(r^2_{\bar a}),\\
{}^4\hat{R}^\tau{}_{r\tau s}(\tau,\vec\sigma) &= \partial_\tau\,{}^4\hat{\Gamma}^\tau_{rs} + O(r^2_{\bar a}) = \epsilon\partial_\tau\,{}^3\hat{K}_{rs} + O(r^2_{\bar a}) =\\
&= \sqrt{3}\frac{4\pi G}{c^3}\sum_{\bar a}\Big[2\delta_{rs}\gamma_{\bar ar}\pi_{\bar a}(\tau,\vec\sigma) +\\
&+ [1-\delta_{rs}]\sum_w \gamma_{\bar aw}[1-2(\delta_{rw}+\delta_{sw})]\\
&\quad \frac{\partial^2}{\partial(\sigma^w)^2}\int_{\sigma^r}^\infty d\sigma_1^r \int_{\sigma^s}^\infty d\sigma_1^s\;\partial_\tau\pi_{\bar a}(\tau,\sigma_1^r\sigma_1^s\sigma^{k\neq r,s})\Big] + O(r^2_{\bar a}),\\
{}^4\hat{R}^\tau{}_{ruv}(\tau,\vec\sigma) &= \partial_u\,{}^4\hat{\Gamma}^\tau_{rv} - \partial_v\,{}^4\hat{\Gamma}^\tau_{ur} = +\epsilon(\partial_u\,{}^3\hat{K}_{rv} - \partial_v\,{}^3\hat{K}_{ru}) =\\
&= \sqrt{3}\frac{4\pi G}{c^3}\sum_{\bar a}\Big[2\gamma_{\bar ar}(\delta_{rv}\partial_u - \delta_{ru}\partial_v)\pi_{\bar a}(\tau,\vec\sigma) + \sum_t \gamma_{\bar at}\\
&\quad \Big((1-\delta_{rv})[1-2(\delta_{rt}+\delta_{vt})]\frac{\partial^3}{\partial\sigma^u\partial(\sigma^t)^2}\int_{\sigma^r}^\infty dw^r\int_{\sigma^v}^\infty dw^v\;\pi_{\bar a}(\tau,w^rw^v\sigma^{k\neq r,v}) -\\
&- (1-\delta_{ru})[1-2(\delta_{rt}+\delta_{ut})]\frac{\partial^3}{\partial\sigma^v\partial(\sigma^t)^2}\int_{\sigma^r}^\infty dw^r\int_{\sigma^u}^\infty dw^u\;\pi_{\bar a}(\tau,w^rw^u\sigma^{k\neq r,u})\Big)\Big] +\\
&+ O(r^2_{\bar a}),\\
{}^4\hat{R}^r{}_{\tau s\tau}(\tau,\vec\sigma) &= -\epsilon\partial_\tau\,{}^3\hat{K}_{rs}(\tau,\vec\sigma) + O(r^2_{\bar a}) =\\
&= -\sqrt{3}\frac{4\pi G}{c^3}\sum_{\bar a}\Big[2\delta_{rs}\gamma_{\bar ar}\partial_\tau\pi_{\bar a}(\tau,\vec\sigma) +\\
&+ [1-\delta_{rs}]\sum_w \gamma_{\bar aw}[1-2(\delta_{rw}+\delta_{sw})]\\
&\quad \frac{\partial^2}{\partial(\sigma^w)^2}\int_{\sigma^r}^\infty d\sigma_1^r \int_{\sigma^s}^\infty d\sigma_1^s\;\partial_\tau\pi_{\bar a}(\tau,\sigma_1^r\sigma_1^s\sigma^{k\neq r,s})\Big] + O(r^2_{\bar a}),\\
{}^4\hat{R}^r{}_{\tau uv}(\tau,\vec\sigma) &= {}^4\hat{R}^\tau{}_{ruv}(\tau,\vec\sigma),\\
{}^4\hat{R}^r{}_{suv}(\tau,\vec\sigma) &= \partial_u\,{}^4\hat{\Gamma}^r_{sv} - \partial_v\,{}^4\hat{\Gamma}^r_{us} + O(r^2_{\bar a}) =\\
&= \frac{1}{\sqrt{3}}\sum_{\bar a}\Big(\delta_{su}\Big[\gamma_{\bar ar}\partial_v\partial_r r_{\bar a}(\tau,\vec\sigma) - \frac{1}{2}\sum_w \gamma_{\bar aw}\frac{\partial^2}{\partial\sigma^v\partial\sigma^r}\int d^3\sigma_1\frac{(\partial^2_{1w}r_{\bar a})(\tau,\vec\sigma)}{4\pi|\vec\sigma-\vec\sigma_1|}\Big] -\\
&- \delta_{sv}\Big[\gamma_{\bar as}\partial_u\partial_r r_{\bar a}(\tau,\vec\sigma) - \frac{1}{2}\sum_w \gamma_{\bar aw}\frac{\partial^2}{\partial\sigma^u\partial\sigma^r}\int d^3\sigma_1\frac{(\partial^2_{1w}r_{\bar a})(\tau,\vec\sigma)}{4\pi|\vec\sigma-\vec\sigma_1|}\Big] +\\
&+ \delta_{rv}\Big[\gamma_{\bar av}\partial_u\partial_s r_{\bar a}(\tau,\vec\sigma) - \frac{1}{2}\sum_w \gamma_{\bar aw}\frac{\partial^2}{\partial\sigma^u\partial\sigma^s}\int d^3\sigma_1\frac{(\partial^2_{1w}r_{\bar a})(\tau,\vec\sigma)}{4\pi|\vec\sigma-\vec\sigma_1|}\Big] -
\end{aligned}
$$

$$- \delta_{ru}\Big[\gamma_{\bar{a}u}\partial_v\partial_s r_{\bar{a}}(\tau,\vec{\sigma}) - \frac{1}{2}\sum_w \gamma_{\bar{a}w}\frac{\partial^2}{\partial\sigma^v\partial\sigma^s}\int d^3\sigma_1 \frac{(\partial^2_{1w} r_{\bar{a}})(\tau,\vec{\sigma})}{4\pi|\vec{\sigma}-\vec{\sigma}_1|}\Big]\Big) + O(r^2_{\bar{a}}). \quad \text{(D7)}$$

Then we get the following linearization of the Ricci tensor and of the curvature scalar

$$\begin{aligned}
{}^4\hat{R}_{AB} &= {}^4\hat{R}^E{}_{AEB} = {}^4\hat{R}^\tau{}_{A\tau B} + {}^4\hat{R}^r{}_{ArB},\\
\\
{}^4\hat{R}_{\tau\tau} &= \sum_r {}^4\hat{R}^r{}_{\tau r\tau} = -\epsilon \sum_r \partial_\tau\, {}^3\hat{K}_{rr} + O(r^2_{\bar{a}}) \overset{\circ}{=} 0 + O(r^2_{\bar{a}}),\\
{}^4\hat{R}_{\tau u} &= \sum_r {}^4\hat{R}^r{}_{\tau ru} = \epsilon \sum_r (\partial_r\, {}^3\hat{K}_{ru} - \partial_u\, {}^3\hat{K}_{rr}) + O(r^2_{\bar{a}}) =\\
&= \epsilon \sum_r \partial_r\, {}^3\hat{K}_{ru} + O(r^2_{\bar{a}}) \overset{\circ}{=} 0 + O(r^2_{\bar{a}}),\\
{}^4\hat{R}_{rs} &= {}^4\hat{R}^\tau{}_{r\tau s} + \sum_u {}^4\hat{R}^u{}_{rus} =\\
&= \epsilon\, \partial_\tau\, {}^3\hat{K}_{rs} + \sum_u (\partial_u\, {}^3\hat{\Gamma}^u_{rs} - \partial_{(s}\, {}^3\hat{\Gamma}^u_{r)u}) + O(r^2_{\bar{a}}),\\
\\
{}^4\hat{R} &= {}^4\eta^{AB}\, {}^4\hat{R}_{AB} + O(r^2_{\bar{a}}) = \epsilon({}^4\hat{R}_{\tau\tau} - \sum_r {}^4\hat{R}_{rr}) + O(r^2_{\bar{a}}) =\\
&= -\epsilon \sum_{ur} (\partial_u\, {}^3\hat{\Gamma}^u_{rr} - \partial_r\, {}^3\hat{\Gamma}^u_{ru}) + O(r^2_{\bar{a}}) = 0 + O(r^2_{\bar{a}}). \qquad \text{(D8)}
\end{aligned}$$

We see that the Einstein's equations ${}^4\hat{R}_{\tau A} \overset{\circ}{=} 0$ (corresponding to the super-hamiltonian and super-momentum constraints) are satisfied as a consequence of the results of Section II.

The spatial Einstein equations ${}^4\hat{R}_{rs} \overset{\circ}{=} 0$ are independent from the shift functions n_r and, after having used the Hamilton equations (5.6) to eliminate $\partial_\tau \pi_{\bar{a}}(\tau,\vec{\sigma})$, it can be checked with a long but straightforward calculation that they are satisfied

$$\begin{aligned}
{}^4\hat{R}_{rs}(\tau,\vec{\sigma}) &= \epsilon\, \partial_\tau\, {}^3\hat{K}_{rs} + \sum_u (\partial_u\, {}^3\hat{\Gamma}^u_{rs} - \partial_{(s}\, {}^3\hat{\Gamma}^u_{r)u})(\tau,\vec{\sigma}) + O(r^2_{\bar{a}}) =\\
&= \sqrt{3}\frac{4\pi G}{c^3} \sum_{\bar{a}} \Big[2\delta_{rs}\gamma_{\bar{a}s}\partial_\tau \pi_{\bar{a}}(\tau,\vec{\sigma}) +\\
&+ (1-\delta_{rs}) \sum_w \gamma_{\bar{a}w}[1 - 2(\delta_{rw} + \delta_{sw})]\\
&\quad \partial^2_w \int_{\sigma^r}^\infty d\sigma^r_1 \int_{\sigma^s}^\infty d\sigma^s_1\ \partial_\tau\, \pi_{\bar{a}}(\tau, \sigma^r_1 \sigma^s_1 \sigma^{k\neq r,s})\Big] +\\
&+ \frac{1}{\sqrt{3}} \sum_{\bar{a}} \Big(\Big[-\delta_{rs}(\gamma_{\bar{a}r}\triangle + \frac{1}{2}\sum_w \gamma_{\bar{a}w}\partial^2_w) + (\gamma_{\bar{a}r} + \gamma_{\bar{a}s})\partial_r\partial_s\Big] r_{\bar{a}}(\tau,\vec{\sigma}) -\\
&- \sum_w \gamma_{\bar{a}w}\partial_r\partial_s \int d^3\sigma_1 \frac{\partial^2_{1w} r_{\bar{a}}(\tau,\vec{\sigma}_1)}{4\pi|\vec{\sigma}-\vec{\sigma}_1|}\Big) +\\
&+ \frac{\sqrt{3}}{2} \sum_{\bar{a}w} \gamma_{\bar{a}w}\partial_r\partial_s \int d^3\sigma_1 \frac{\partial^2_{1w} r_{\bar{a}}(\tau,\vec{\sigma}_1)}{4\pi|\vec{\sigma}-\vec{\sigma}_1|} \overset{\circ}{=} 0 + O(r^2_{\bar{a}}). \qquad \text{(D9)}
\end{aligned}$$

Then we can obtain the field strength in our gauge by using the equation ${}^4\overset{\circ}{\Omega}_{AB(\alpha)(\beta)} = {}^4_{(\Sigma)}\check{\tilde{E}}^C_{(\alpha)}\, {}^4_{(\Sigma)}\check{\tilde{E}}^D_{(\beta)}\, {}^4R_{CDAB} = \partial_A\, {}^4\overset{\circ}{\omega}_{B(\alpha)(\beta)} - \partial_B\, {}^4\overset{\circ}{\omega}_{A(\alpha)(\beta)} + {}^4\overset{\circ}{\omega}_{A(\alpha)(\gamma)}\ {}^4\overset{\circ}{\omega}{}^{(\gamma)}_{B(\beta)} - {}^4\overset{\circ}{\omega}_{B(\alpha)(\gamma)}\ {}^4\overset{\circ}{\omega}{}^{(\gamma)}_{A(\beta)}$.

3. The 4-Weyl and 4-Bel-Robinson Tensors and the Kretschmann Invariant.

In this Section we will study the radiation gauge form of others 4-tensors.

Let us evaluate the 4-Weyl tensor ${}^4C_{ABCD} = {}^4R_{ABCD} + \frac{1}{2}({}^4R_{AC}\, {}^4g_{BD} - {}^4R_{BC}\, {}^4g_{AD} + {}^4R_{BD}\, {}^4g_{AC} - {}^4R_{AD}\, {}^4g_{BC}) + \frac{1}{6}({}^4g_{AC}\, {}^4g_{BD} - {}^4g_{AD}\, {}^4g_{BC})\, {}^4R \stackrel{\circ}{=} {}^4R_{ABCD}$. We get

$$\begin{aligned}
{}^4C_{rstu} &= {}^4R_{rstu} + \frac{1}{2}(-\epsilon {}^4R_{rt}\delta_{su} + \epsilon {}^4R_{st}\delta_{ru} - \epsilon {}^4R_{su}\delta_{rt} + \epsilon {}^4R_{ru}\delta_{st}) = \\
&= -\epsilon \frac{1}{\sqrt{3}} \sum_{\bar a} \Big(\delta_{st} \Big[\gamma_{\bar a r} \partial_u \partial_r r_{\bar a}(\tau, \vec \sigma) - \frac{1}{2} \sum_w \gamma_{\bar a w} \frac{\partial^2}{\partial \sigma^u \partial \sigma^r} \int d^3\sigma_1 \frac{(\partial^2_{1w} r_{\bar a})(\tau, \vec \sigma)}{4\pi |\vec \sigma - \vec \sigma_1|} \Big] - \\
&- \delta_{su} \Big[\gamma_{\bar a s} \partial_t \partial_r r_{\bar a}(\tau, \vec \sigma) - \frac{1}{2} \sum_w \gamma_{\bar a w} \frac{\partial^2}{\partial \sigma^t \partial \sigma^r} \int d^3\sigma_1 \frac{(\partial^2_{1w} r_{\bar a})(\tau, \vec \sigma)}{4\pi |\vec \sigma - \vec \sigma_1|} \Big] + \\
&+ \delta_{ru} \Big[\gamma_{\bar a u} \partial_t \partial_s r_{\bar a}(\tau, \vec \sigma) - \frac{1}{2} \sum_w \gamma_{\bar a w} \frac{\partial^2}{\partial \sigma^t \partial \sigma^s} \int d^3\sigma_1 \frac{(\partial^2_{1w} r_{\bar a})(\tau, \vec \sigma)}{4\pi |\vec \sigma - \vec \sigma_1|} \Big] - \\
&- \delta_{rt} \Big[\gamma_{\bar a t} \partial_u \partial_s r_{\bar a}(\tau, \vec \sigma) - \frac{1}{2} \sum_w \gamma_{\bar a w} \frac{\partial^2}{\partial \sigma^u \partial \sigma^s} \int d^3\sigma_1 \frac{(\partial^2_{1w} r_{\bar a})(\tau, \vec \sigma)}{4\pi |\vec \sigma - \vec \sigma_1|} \Big] \Big) + \\
&+ \frac{1}{2} \Big(-\epsilon \delta_{su} \sqrt{3} \frac{4\pi G}{c^3} \sum_{\bar a} \Big[2\delta_{rt} \gamma_{\bar a t} \partial_\tau \pi_{\bar a}(\tau, \vec \sigma) + \\
&+ (1 - \delta_{rt}) \sum_w \gamma_{\bar a w} [1 - 2(\delta_{rw} + \delta_{tw})] \\
&\quad \partial^2_w \int_{\sigma^r}^\infty d\sigma^r_1 \int_{\sigma^t}^\infty d\sigma^t_1\ \partial_\tau\, \pi_{\bar a}(\tau, \sigma^r_1 \sigma^s_1 \sigma^{k \neq r,t}) \Big] + \\
&+ \frac{1}{\sqrt{3}} \sum_{\bar a} \Big(\Big[-\delta_{rt} (\gamma_{\bar a r} \triangle + \frac{1}{2} \sum_w \gamma_{\bar a w} \partial^2_w) + (\gamma_{\bar a r} + \gamma_{\bar a t}) \partial_r \partial_t \Big] r_{\bar a}(\tau, \vec \sigma) - \\
&- \sum_w \gamma_{\bar a w} \partial_r \partial_t \int d^3\sigma_1 \frac{\partial^2_{1w} r_{\bar a}(\tau, \vec \sigma_1)}{4\pi |\vec \sigma - \vec \sigma_1|} \Big) + \\
&+ \frac{\sqrt{3}}{2} \sum_{\bar a w} \gamma_{\bar a w} \partial_r \partial_t \int d^3\sigma_1 \frac{\partial^2_{1w} r_{\bar a}(\tau, \vec \sigma_1)}{4\pi |\vec \sigma - \vec \sigma_1|} + \\
&+ \epsilon \delta_{ru} \sqrt{3} \frac{4\pi G}{c^3} \sum_{\bar a} \Big[2\delta_{st} \gamma_{\bar a t} \partial_\tau \pi_{\bar a}(\tau, \vec \sigma) + \\
&+ (1 - \delta_{st}) \sum_w \gamma_{\bar a w} [1 - 2(\delta_{sw} + \delta_{tw})] \times \\
&\quad \partial^2_w \int_{\sigma^s}^\infty d\sigma^s_1 \int_{\sigma^t}^\infty d\sigma^t_1\ \partial_\tau\, \pi_{\bar a}(\tau, \sigma^s_1 \sigma^t_1 \sigma^{k \neq s,t}) \Big] + \\
&+ \frac{1}{\sqrt{3}} \sum_{\bar a} \Big(\Big[-\delta_{st} (\gamma_{\bar a s} \triangle + \frac{1}{2} \sum_w \gamma_{\bar a w} \partial^2_w) + (\gamma_{\bar a s} + \gamma_{\bar a s}) \partial_s \partial_t \Big] r_{\bar a}(\tau, \vec \sigma) - \\
&- \sum_w \gamma_{\bar a w} \partial_s \partial_t \int d^3\sigma_1 \frac{\partial^2_{1w} r_{\bar a}(\tau, \vec \sigma_1)}{4\pi |\vec \sigma - \vec \sigma_1|} \Big) + \\
&+ \frac{\sqrt{3}}{2} \sum_{\bar a w} \gamma_{\bar a w} \partial_s \partial_t \int d^3\sigma_1 \frac{\partial^2_{1w} r_{\bar a}(\tau, \vec \sigma_1)}{4\pi |\vec \sigma - \vec \sigma_1|} - \\
&- \epsilon \delta_{rt} \sqrt{3} \frac{4\pi G}{c^3} \sum_{\bar a} \Big[2\delta_{su} \gamma_{\bar a u} \partial_\tau \pi_{\bar a}(\tau, \vec \sigma) +
\end{aligned}$$

$$
\begin{aligned}
&+ (1-\delta_{su}) \sum_w \gamma_{\bar{a}w}[1-2(\delta_{sw}+\delta_{uw})] \\
&\quad \partial^2_w \int_{\sigma^s}^{\infty} d\sigma^s_1 \int_{\sigma^u}^{\infty} d\sigma^u_1 \;\partial_\tau\, \pi_{\bar{a}}(\tau, \sigma^s_1 \sigma^u_1 \sigma^{k\neq s,u})\Big] + \\
&+ \frac{1}{\sqrt{3}} \sum_{\bar{a}} \Big(\Big[-\delta_{su}(\gamma_{\bar{a}s}\triangle + \frac{1}{2}\sum_w \gamma_{\bar{a}w}\partial^2_w) + (\gamma_{\bar{a}s}+\gamma_{\bar{a}u})\partial_s\partial_u\Big] r_{\bar{a}}(\tau,\vec{\sigma}) - \\
&- \sum_w \gamma_{\bar{a}w}\partial_s\partial_u \int d^3\sigma_1 \frac{\partial^2_{1w} r_{\bar{a}}(\tau,\vec{\sigma}_1)}{4\pi|\vec{\sigma}-\vec{\sigma}_1|}\Big) + \\
&+ \frac{\sqrt{3}}{2} \sum_{\bar{a}w} \gamma_{\bar{a}w}\partial_s\partial_u \int d^3\sigma_1 \frac{\partial^2_{1w} r_{\bar{a}}(\tau,\vec{\sigma}_1)}{4\pi|\vec{\sigma}-\vec{\sigma}_1|} + \\
&+ \epsilon\delta_{st}\sqrt{3}\frac{4\pi G}{c^3} \sum_{\bar{a}} \Big[2\delta_{rs}\gamma_{\bar{a}u}\partial_\tau \pi_{\bar{a}}(\tau,\vec{\sigma}) + \\
&+ (1-\delta_{ru}) \sum_w \gamma_{\bar{a}w}[1-2(\delta_{rw}+\delta_{uw})] \\
&\quad \partial^2_w \int_{\sigma^r}^{\infty} d\sigma^r_1 \int_{\sigma^u}^{\infty} d\sigma^u_1 \;\partial_\tau\, \pi_{\bar{a}}(\tau, \sigma^r_1 \sigma^u_1 \sigma^{k\neq r,u})\Big] + \\
&+ \frac{1}{\sqrt{3}} \sum_{\bar{a}} \Big(\Big[-\delta_{ru}(\gamma_{\bar{a}r}\triangle + \frac{1}{2}\sum_w \gamma_{\bar{a}w}\partial^2_w) + (\gamma_{\bar{a}r}+\gamma_{\bar{a}s})\partial_r\partial_u\Big] r_{\bar{a}}(\tau,\vec{\sigma}) - \\
&- \sum_w \gamma_{\bar{a}w}\partial_r\partial_u \int d^3\sigma_1 \frac{\partial^2_{1w} r_{\bar{a}}(\tau,\vec{\sigma}_1)}{4\pi|\vec{\sigma}-\vec{\sigma}_1|}\Big) + \\
&+ \frac{\sqrt{3}}{2} \sum_{\bar{a}w} \gamma_{\bar{a}w}\partial_r\partial_u \int d^3\sigma_1 \frac{\partial^2_{1w} r_{\bar{a}}(\tau,\vec{\sigma}_1)}{4\pi|\vec{\sigma}-\vec{\sigma}_1|}\Big) + O(r^2_{\bar{a}}),
\end{aligned}
$$

$$
\begin{aligned}
{}^4C_{\tau ruv} &= {}^4R_{\tau ruv} + O(r^2_{\bar{a}}) = \\
&= \epsilon\sqrt{3}\frac{4\pi G}{c^3} \sum_{\bar{a}} \Big[2\gamma_{\bar{a}r}(\delta_{rv}\partial_u - \delta_{ru}\partial_v)\pi_{\bar{a}}(\tau,\vec{\sigma}) + \sum_t \gamma_{\bar{a}t} \\
&\quad \Big((1-\delta_{rv})[1-2(\delta_{rt}+\delta_{vt})]\frac{\partial^3}{\partial\sigma^u\partial(\sigma^t)^2} \int_{\sigma^r}^{\infty} dw^r \int_{\sigma^v}^{\infty} dw^v\, \pi_{\bar{a}}(\tau, w^r w^v \sigma^{k\neq r,v}) - \\
&- (1-\delta_{ru})[1-2(\delta_{rt}+\delta_{ut})]\frac{\partial^3}{\partial\sigma^v\partial(\sigma^t)^2} \int_{\sigma^r}^{\infty} dw^r \int_{\sigma^u}^{\infty} dw^u\, \pi_{\bar{a}}(\tau, w^r w^u \sigma^{k\neq r,u})\Big)\Big] + \\
&+ O(r^2_{\bar{a}}),
\end{aligned}
$$

$$
\begin{aligned}
{}^4C_{\tau r\tau s} &= {}^4R_{\tau r\tau s} + \frac{1}{2}\epsilon\, {}^4R_{rs} = \\
&= \epsilon\sqrt{3}\frac{4\pi G}{c^3} \sum_{\bar{a}} \Big[2\delta_{rs}\gamma_{\bar{a}r}\pi_{\bar{a}}(\tau,\vec{\sigma}) + \\
&+ [1-\delta_{rs}] \sum_w \gamma_{\bar{a}w}[1-2(\delta_{rw}+\delta_{sw})] \\
&\quad \frac{\partial^2}{\partial(\sigma^w)^2} \int_{\sigma^r}^{\infty} d\sigma^r_1 \int_{\sigma^s}^{\infty} d\sigma^s_1 \;\partial_\tau \pi_{\bar{a}}(\tau, \sigma^r_1 \sigma^s_1 \sigma^{k\neq r,s})\Big] + \\
&+ \frac{\epsilon}{2}\sqrt{3}\frac{4\pi G}{c^3} \sum_{\bar{a}} \Big[2\delta_{rs}\gamma_{\bar{a}s}\partial_\tau \pi_{\bar{a}}(\tau,\vec{\sigma}) +
\end{aligned}
$$

$$
\begin{aligned}
&+ (1-\delta_{rs}) \sum_w \gamma_{\bar{a}w} [1 - 2(\delta_{rw} + \delta_{sw})] \\
&\quad \partial_w^2 \int_{\sigma^r}^{\infty} d\sigma_1^r \int_{\sigma^s}^{\infty} d\sigma_1^s\, \partial_\tau\, \pi_{\bar{a}}(\tau, \sigma_1^r \sigma_1^s \sigma^{k\not= r,s}) \Big] + \\
&+ \frac{1}{\sqrt{3}} \sum_{\bar{a}} \Big(\Big[-\delta_{rs}(\gamma_{\bar{a}r} \triangle + \frac{1}{2} \sum_w \gamma_{\bar{a}w} \partial_w^2) + (\gamma_{\bar{a}r} + \gamma_{\bar{a}s}) \partial_r \partial_s \Big] r_{\bar{a}}(\tau, \vec{\sigma}) - \\
&- \sum_w \gamma_{\bar{a}w} \partial_r \partial_s \int d^3\sigma_1 \frac{\partial_{1w}^2 r_{\bar{a}}(\tau, \vec{\sigma}_1)}{4\pi |\vec{\sigma} - \vec{\sigma}_1|} \Big) + \\
&+ \frac{\sqrt{3}}{2} \sum_{\bar{a}w} \gamma_{\bar{a}w} \partial_r \partial_s \int d^3\sigma_1 \frac{\partial_{1w}^2 r_{\bar{a}}(\tau, \vec{\sigma}_1)}{4\pi |\vec{\sigma} - \vec{\sigma}_1|} + O(r_{\bar{a}}^2),
\end{aligned}
$$

$$
\begin{aligned}
&{}^4C_{\tau s\tau\tau} = {}^4R_{\tau s\tau\tau} = 0, \\
&{}^4C_{\tau\tau\tau\tau} = {}^4R_{\tau\tau\tau\tau} = 0.
\end{aligned}
\qquad \text{(D10)}
$$

On the solutions of the Hamilton equations the Weyl tensor ${}^4\hat{C}^A{}_{BCD}(\tau, \vec{\sigma})$ coincides with the Riemann tensor.

Let us consider the congruence of time-like observers whose 4-velocities are given by the unit normals to $\Sigma_\tau^{(WSW)}$, viz. ${}^4_{(\Sigma)}\check{\tilde{E}}^A_{(o)}(\tau, \vec{\sigma}) = \epsilon l^A(\tau, \vec{\sigma}) = -(1; \delta^{rs} n_s(\tau, \vec{\sigma}))$, ${}^4_{(\Sigma)}\check{\tilde{E}}^{(o)}_A(\tau, \vec{\sigma}) = l_A(\tau, \vec{\sigma}) = -\epsilon(1; 0)$ [see Eqs.(2.2)]. The *electric and magnetic parts of the Weyl tensor* with respect to these time-like observers are [25]

$$
\begin{aligned}
{}^4\hat{E}_{AB}(\tau, \vec{\sigma}) &= {}^4\hat{E}_{BA}(\tau, \vec{\sigma}) = [{}^4\hat{C}_{ACBD}\, l^C\, l^D](\tau, \vec{\sigma}), \\
&\qquad [{}^4\hat{E}_{AB}\, l^B](\tau, \vec{\sigma}) = -\Big[{}^4\hat{E}_{A\tau} + {}^4\hat{E}_{Ar} n_r \Big](\tau, \vec{\sigma}) = 0, \quad {}^4\hat{E}^A{}_A(\tau, \vec{\sigma}) = 0, \\
{}^4\hat{E}_{rs}(\tau, \vec{\sigma}) &= [{}^4\hat{C}_{r\tau s\tau} + ({}^4\hat{C}_{r\tau su} + {}^4\hat{C}_{s\tau ru}) n_u + {}^4\hat{C}_{rusv} n_u n_v](\tau, \vec{\sigma}) = \\
&= {}^4\hat{C}_{r\tau s\tau} + O(r_{\bar{a}}^2), \\
{}^4\hat{E}_{r\tau}(\tau, \vec{\sigma}) &= [{}^4\hat{C}_{r\tau\tau u} n_u + {}^4\hat{C}_{ru\tau v} n_u n_v](\tau, \vec{\sigma}) = -[{}^4\hat{E}_{rs} n_s](\tau, \vec{\sigma}) = \\
&= 0 + O(r_{\bar{a}}^2), \\
{}^4\hat{E}_{\tau\tau}(\tau, \vec{\sigma}) &= [{}^4\hat{C}_{\tau r\tau s} n_r n_s](\tau, \vec{\sigma}) = -[{}^4\hat{E}_{\tau s} n_s](\tau, \vec{\sigma}) = \\
&= 0 + O(r_{\bar{a}}^2),
\end{aligned}
$$

$$
\begin{aligned}
{}^4\hat{H}_{AB}(\tau, \vec{\sigma}) &= {}^4\hat{H}_{BA}(\tau, \vec{\sigma}) = \frac{1}{2} \epsilon_{BE}{}^{CD}\, [{}^4\hat{C}_{AFCD}\, l^E\, l^F](\tau, \vec{\sigma}) = [*{}^4\hat{C}_{ABCD} l^C l^D](\tau, \vec{\sigma}), \\
&\qquad [{}^4\hat{H}_{AB}\, l^B](\tau, \vec{\sigma}) = -\Big[{}^4\hat{H}_{A\tau} + {}^4H_{Ar} n_r \Big](\tau, \vec{\sigma}) = 0, \quad {}^4\hat{H}^A{}_A(\tau, \vec{\sigma}) = 0, \\
{}^4\hat{H}_{rs}(\tau, \vec{\sigma}) &= [*{}^4\hat{C}_{r\tau s\tau} + (*{}^4\hat{C}_{r\tau su} + *{}^4C_{s\tau ru}) n_u + *{}^4\hat{C}_{rusv} n_u n_v](\tau, \vec{\sigma}) = \\
&= *{}^4\hat{C}_{r\tau s\tau} + O(r_{\bar{a}}^2),
\end{aligned}
$$

---

[25] $*{}^4\hat{C}_{ABCD} = \frac{1}{2} \epsilon_{AB}{}^{EF}\, {}^4\hat{C}_{EFCD} = \frac{1}{2} \epsilon_{CD}{}^{EF}\, {}^4\hat{C}_{ABEF}$ is, due to the Lanczos identity, the unique Hodge dual of the Weyl tensor [47] (the Riemann tensor has different left and right duals).

$$\begin{aligned}
{}^4\hat{H}_{r\tau}(\tau,\vec{\sigma}) &= [*{}^4\hat{C}_{r\tau\tau u} n_u + *{}^4\hat{C}_{ru\tau v} n_u n_v](\tau,\vec{\sigma}) = -[{}^4\hat{H}_{rs} n_s](\tau,\vec{\sigma}) = \\
&= 0 + O(r_{\bar{a}}^2),\\
{}^4\hat{H}_{\tau\tau}(\tau,\vec{\sigma}) &= [*{}^4\hat{C}_{\tau r\tau s} n_r n_s](\tau,\vec{\sigma}) = -[{}^4\hat{H}_{\tau s} n_s](\tau,\vec{\sigma}) = \\
&= 0 + O(r_{\bar{a}}^2).
\end{aligned} \qquad \text{(D11)}$$

Both ${}^4\hat{E}_{AB}$ and ${}^4\hat{H}_{AB}$ have five independent components. See Ref. [47] for an analogous decomposition of the Riemann tensor.

The four eigenvalues Λ_α, $\alpha = 1,..,4$, of the Weyl tensor are

$$\begin{aligned}
&{}^4\hat{C}_{ABCD}\,{}^4\hat{C}^{ABCD} = O(r_{\bar{a}}^2),\\
&{}^4\hat{C}_{AB}{}^{CD}\,\epsilon_{CDEF}\,{}^4\hat{C}^{ABEF} = O(r_{\bar{a}}^2),\\
&{}^4\hat{C}^{AB}{}_{CD}\,{}^4\hat{C}^{CD}{}_{EF}\,{}^4\hat{C}^{EF}{}_{AB} = O(r_{\bar{a}}^3),\\
&{}^4\hat{C}^{ABCD}\,\epsilon_{CDEF}\,{}^4\hat{C}^{EF}{}_{UV}\,{}^4\hat{C}^{UV}{}_{AB} = O(r_{\bar{a}}^3).
\end{aligned} \qquad \text{(D12)}$$

As shown in Ref. [15], the 4-coordinate system $\sigma^A = \{\tau, \vec{\sigma}\}$, corresponding to our completely fixed 3-orthogonal gauge, is identifiable by means of 4 gauge-fixing constraints $\sigma^A - F^{\bar{A}}[\Lambda_\alpha(\tau,\vec{\sigma})] \approx 0$, where the $F^{\bar{A}}$ are 4 suitable scalar functions of the 4 eigenvalues of the Weyl tensor such that in the linearized theory we have $F^{\bar{A}} = F^{\bar{A}}_{(L)} + O(r_{\bar{a}}^2)$ with $F^{\bar{A}}_{(L)} = O(r_{\bar{a}})$.

The *Kretschmann invariant and pseudo-invariant* are respectively $[*{}^4\hat{R}^{ABCD} = \frac{1}{2}\epsilon^{ABEF}\,{}^4\hat{R}_{EF}{}^{CD}]$

$$\begin{aligned}
&\hat{\mathbf{R}}\cdot\hat{\mathbf{R}} = {}^4\hat{R}_{ABCD}\,{}^4\hat{R}^{ABCD} = O(r_{\bar{a}}^2),\\
&*\hat{\mathbf{R}}\cdot\hat{\mathbf{R}} = *{}^4\hat{R}^{ABCD}\,{}^4\hat{R}_{ABCD} = O(r_{\bar{a}}^2).
\end{aligned} \qquad \text{(D13)}$$

In Ref. [16] it is suggested that in presence of matter a coordinate-independent characterization of gravito-magnetism is $*\hat{\mathbf{R}}\cdot\hat{\mathbf{R}} \neq 0$.

The *Bel-Robinson and Bel tensors* are respectively

$$\begin{aligned}
{}^4\hat{\mathcal{T}}_{ABCD} &= {}^4\hat{C}_{AECF}\,{}^4\hat{C}_B{}^E{}_D{}^F + {}^4\hat{C}_{AEDF}\,{}^4\hat{C}_B{}^E{}_C{}^F -\\
&- \frac{1}{2}{}^4g_{AB}\,{}^4\hat{C}_{EFCG}\,{}^4\hat{C}^{EF}{}_D{}^G - \frac{1}{2}{}^4g_{CD}\,{}^4\hat{C}_{AEFG}\,{}^4\hat{C}_B{}^{EFG} +\\
&+ \frac{1}{8}{}^4g_{AB}\,{}^4g_{CD}\,{}^4\hat{C}_{EFGH}\,{}^4\hat{C}^{EFGH} = O(r_{\bar{a}}^2),\\
{}^4\hat{\mathcal{B}} &= {}^4\hat{R}_{AECF}\,{}^4\hat{R}_B{}^E{}_D{}^F + {}^4\hat{R}_{AEDF}\,{}^4\hat{R}_B{}^E{}_C{}^F -\\
&- \frac{1}{2}{}^4g_{AB}\,{}^4\hat{R}_{EFCG}\,{}^4\hat{R}^{EF}{}_D{}^G - \frac{1}{2}{}^4g_{CD}\,{}^4\hat{R}_{AEFG}\,{}^4\hat{R}_B{}^{EFG} +\\
&+ \frac{1}{8}{}^4g_{AB}\,{}^4g_{CD}\,{}^4\hat{R}_{EFGH}\,{}^4\hat{R}^{EFGH} = O(r_{\bar{a}}^2),\\
\\
&{}^4\hat{\mathcal{B}}_{ABCD} = {}^4\hat{\mathcal{T}}_{ABCD} + {}^4\hat{\mathcal{M}}_{ABCD} + {}^4\hat{\mathcal{Q}}_{ABCD},
\end{aligned} \qquad \text{(D14)}$$

where ${}^4\hat{\mathcal{Q}}_{ABCD} = \frac{1}{6}{}^4\hat{R}\,[{}^4\hat{C}_{ACBD} + {}^4\hat{C}_{ADBC}]$ is the matter-gravity coupling tensor and ${}^4\hat{\mathcal{M}}_{ABCD}$ is the pure matter gravitational super-energy tensor (see Ref. [47] for its expression).

REFERENCES

[1] L.Lusanna, *The Rest-Frame Instant Form of Metric Gravity*, Gen.Rel.Grav. **33**, 1579 (2001)(gr-qc/0101048).

[2] L.Lusanna and S.Russo, *A New Parametrization for Tetrad Gravity*, Gen.Rel.Grav. **34**, 189 (2002)(gr-qc/0102074).

[3] R.De Pietri, L.Lusanna, L.Martucci and S.Russo, *Dirac's Observables for the Rest-Frame Instant Form of Tetrad Gravity in a Completely Fixed 3-Orthogonal Gauge*, Gen.Rel.Grav. **34**, 877 (2002) (gr-qc/0105084).

[4] L. Lusanna, *The N- and 1-Time Classical Descriptions of N-Body Relativistic Kinematics and the Electromagnetic Interaction*, Int. J. Mod. Phys. **A12**, 645 (1997).

[5] L.Lusanna, *Towards a Unified Description of the Four Interactions in Terms of Dirac-Bergmann Observables*, invited contribution to the book *Quantum Field Theory: a 20th Century Profile* of the Indian National Science Academy, ed. A.N.Mitra, foreward F.J.Dyson (Hindustan Book Agency, New Delhi, 2000) (hep-th/9907081).

[6] R.Arnowitt, S.Deser and C.W.Misner, Phys.Rev. **117**, 1595 (1960); in *Gravitation: an Introduction to Current Research*, ed.L.Witten (Wiley, New York, 1962).

[7] D.Christodoulou and S.Klainerman, *The Global Nonlinear Stability of the Minkowski Space* (Princeton, Princeton, 1993).

[8] J.Frauendiener, *Triads and the Witten Equation*, Class.Quantum Grav. **8**, 1881 (1991).

[9] A.Sen, *On the Existence of Neutrino "Zero-Modes" in Vacuum Space-Times*, J.Math.Phys. **22**, 1781 (1981); *Gravity as a Spin System*, Phys.Lett. **119B**, 89 (1982).
E.Witten, *A New Proof of the Positive Energy Theorem*, Commun.Math.Phys. **80**, 381 (1981).

[10] A.Lichnerowicz, *Propagateurs, Commutateurs et Anticommutateurs en Relativite Generale*, in Les Houches 1963, *Relativity, Groups and Topology*, eds. C.DeWitt and B.DeWitt (Gordon and Breach, New York, 1964).
C.Moreno, *On the Spaces of Positive and Negative Frequency Solutions of the Klein-Gordon Equation in Curved Space-Times*, Rep.Math.Phys. **17**, 333 (1980).

[11] N.D.Birrell and P.C.W.Davies, *Quantum Fields in Curved Space* (Cambridge University Press, Cambridge, 1982).
P.C.W.Davis, *Particles do not Exist*, in *Quantum Theory of Gravity. Essays in Honor of the 60th Birthday of Bryce DeWitt.*, ed. S.Christensen (Hilger, Bristol, 1984).

[12] S.Shanmugadhasan, *Canonical Formalism for Degenerate Lagrangians*, J.Math.Phys. **14**, 677 (1973).
L.Lusanna, *The Shanmugadhasan Canonical Transformation, Function Groups and the Second Noether Theorem*, Int.J.Mod.Phys. **A8**, 4193 (1993).

[13] M.Pauri and M.Vallisneri, *Ephemeral Point-Events: is there a Last Remnant of Physical Objectivity?*, essay for the 70th birthday of R.Torretti, 2002(gr-qc/0203014).

[14] L.Lusanna, *Spacetime, General Covariance, Dirac-Bergmann Observables and Non-Inertial Frames*, talk at the 25th Johns Hopkins Workshop *2001: A Relativistic Spacetime Odyssey* (gr-qc/0205039).

[15] L.Lusanna and M.Pauri, *General Covariance and the Objectivity of Space-Time Point-Events: The Role of Gravitational and Gauge Degrees of Freedom in General Relativity* (gr-qc/0301040).

[16] I.Ciufolini and J.A.Wheeler, *Gravitation and Inertia* (Princeton Univ.Press, Princeton, 1995).

[17] Y.Choquet-Bruhat, A.Fischer and J.E.Marsden, *Maximal Hypersurfaces and Positivity of Mass*, LXVII E.Fermi Summer School of Physics *Isolated Gravitating Systems in General Relativity*, ed. J.Ehlers (North-Holland, Amsterdam, 1979).

[18] A.Lichnerowicz, *L'inte'gration des e'quations de la gravitation relativiste et le proble'me des n corps*, J.Math.Pure Appl. **23**, 37 (1944).

Y.Choquet-Bruhat, *The Cauchy Problem* in *Gravitation: An Introduction to Current Research*, ed.L.Witten (Wiley, New York, 1962).

[19] J.W.York jr, *Gravitational Degrees of Freedom and the Initial Value Problem*, Phys.Rev.Lett. **26**, 1656 (1971).

Role of Conformal Three Geometry in the Dynamics of Gravitation, **28**, 1082 (1972).

Mapping onto Solutions of the Gravitational Initial Value Problem, J.Math.Phys. **13**, 125 (1972).

Conformally Invariant Orthogonal Decompositions of Symmetric Tensors on Riemannian Manifolds and the Initial Value Problem of General Relativity, **14**, 456 (1972).

Covariant Decompositions of Symmetric Tensors in the Theory of Gravitation, Ann.Ins.H.Poincaré **XXI**, 318 (1974).

N.O'Murchadha and J.W.York jr, *Existence and Uniqueness of Solutions of the Hamiltonian Constraint of General Relativity on Compact Manifolds*, J.Math.Phys. **14**, 1551 (1972).

Initial Value Problem of General Relativity. I. General Formulation and Physical Interpretation, Phys.Rev. **D10**, 428 (1974).

Kinematics and Dynamics of General Relativity, in *Sources of Gravitational Radiation*, Battelle-Seattle Workshop 1978, ed.L.L.Smarr (Cambridge Univ.Press, Cambridge, 1979).

A.Qadir and J.A.Wheeler, *York's Cosmic Time Versus Proper Time*, in *From SU(3) to Gravity*, Y.Ne'eman's festschrift, eds. E.Gotsma and G.Tauber (Cambridge Univ.Press, Cambridge, 1985).

[20] P.A.M.Dirac, *Lectures on Quantum Mechanics*, Belfer Graduate School of Science, Monographs Series (Yeshiva University, New York, N.Y., 1964).

[21] M.H.Soffel, *Relativity in Astrometry, Celestial Mechanics and Geodesy* (Springer, Berlin, 1989).

[22] C.W.Misner, K.S.Thorne and J.A.Wheeler, *Gravitation* (Freeman, New York, 1973).

[23] H.Stephani, *General Relativity* (Cambridge Univ.Press, Cambridge, 1996).

[24] B.Schutz, *A First Course in General Relativity* (Cambridge Univ. Press, Cambridge, 1990).

[25] C.R.Chester, *Techniques in Partial Differential Equations* (McGraw-Hill Kogakusha, Tokyo, 1971).

[26] P.Havas, *Simultaneity, Cnventionalism, General Covariance and the Special Theory of Relativity*, Gen.Rel.Grav. **19**, 435 (1987).

R.Anderson, I.Vetharaniam and G.E.Stedman, *Conventionality of Synchronization, Gauge Dependence and Test Theories of Relativity*, Phys.Rep. **295**, 93 (1998).

[27] L.Landau and E.Lifschitz, *The Classical Theory of Fields* (Addison-Wesley, Cambridge, 1951).

[28] S.Persides and D.Papadopoulos, *A Covariant Formulation of the Landau-Lifshitz Complex*, Gen.Rel.Grav. **11**, 233 (1979).

[29] R.M.Wald, *General Relativity* (University of Chicago, Chicago, 1984), pp.292-294.

[30] L.Lusanna and S.Russo, *Tetrad Gravity: II) Dirac's Observables*, Firenze Univ. preprint (gr-qc/9807073).

[31] B.Mashhoon, F.Gronwald, F.H.Hehl and D.S.Theiss, *On Measuring Gravitomagnetism via Spaceborn Clocks: a Gravitomagnetic Clock Effect* (gr-qc/9804008).
B.Mashhoon, F.Gronwald and H.I.M.Lichtenegger, *Gravitomagnetism and the Clock Effect* (gr-qc/9912027).

[32] C.M.Will, *Theory and Experiment in Gravitational Physics*, rev.ed. (Cambridge Univ.Press, Cambridge, 1993).
S.G.Turyshev, *Relativistic Navigation: A Theoretical Foundation*, NASA/JPL No 96-013 (gr-qc/9606063).

[33] L.Blanchet and T.Damour, *Post-Newtonian Generation of Gravitational Waves*, Ann.Inst.H.Poincaré **50**, 377 (1989).
L.Blanchet, T.Damour and G.Schäfer, *Post-Newtonian Hydrodynamics and Post-Newtonian Gravitational Wave Generation for Numerical Relativity*, Mon.Not.R.Astr.Soc. **242**, 289 (1990).
T.Damour, M.Soffel and C.Xu, *General Relativistic Celestial Mechanics*: *I. Method and Definition of Reference Systems*, Phys.Rev. **D43**, 3273 (1991); *II. Translational Equations of Motion*, Phys.Rev. **D45**, 1017 (1992); *III. Rotational Equations of Motion*, Phys.Rev.**D47**, 3124 (1993); *IV. Theory of Satellite Motion*, Phys.Rev.**D49**, 618 (1994).

[34] T.Damour, *Selected Themes in Relativistic Gravity*, in *Relativistic Gravitation and Gravitational Radiation*, Les Houches 1995, eds. J.A.Marck and J.P.Lasota (Cambridge Univ.Press, Cambridge, 1997).

[35] R.DePietri, L.Lusanna and M.Pauri, *Standard and Generalized Newtonian Gravities as 'Gauge' Theories of the Extended Galilei Group: I. The Standard Theory*, Class. Quantum Grav. **12**, 219 (1995).

[36] D.Alba, L.Lusanna and M.Pauri, *Centers of Mass and Rotational Kinematics for the Relativistic N-Body Problem in the Rest-Frame Instant Form*, J.Math.Phys. **43**, 1677 (2002) (hep-th/0102087).

[37] L.Lusanna and Materassi, *A Canonical Decomposition in Collective and Relative Variables of a Klein-Gordon Field in the Rest-Frame Wigner-Covariant Instant Form*, Int.J.Mod.Phys. **A15**, 2821 (2000)(hep-th/9904202).

[38] Ph.DrozVincent, *Relativistic Systems of Interacting Particles*, Phys.Scr. **2**, 129 (1970); *Hamiltonian Systems of Relativistic Particles*, Rep.Math.Phys. **8**, 79 (1975); *Two Body Relativistic Systems*, Ann.Inst.H.Poincaré **27**, 407 (1977); *Action at a Distance and Relativistic Wave Equations for Spinless Quarks*, Phys.Rev. **D19**, 702 (1979).
A.Komar, Phys.Rev. **D18**, *Constraint Formalism of Classical Mechanics* 1881, *Interacting Relativistic Particles* 1887 and *Spacetime Orbits for Interacting Relativistic Particles: Syntactic versus Semantic Observables* 3617 (1978).
I.T.Todorov, *Dynamics of Relativistic Point Particles as a Problem with Constraints*, Dubna preprint JINR E2-10125 (unpublished); *Sur la quantification d'un syste'me me'canique avec des contraintes de deuxie'me classe*, Ann.Int.H.Poincaré **28**, 207 (1978).

V.A.Rizov, H.Sazdjian and I.T.Todorov, *On the Relativistic Quantum Mechanics of Two Interacting Spinless Particles*, Ann.Phys.(N.Y.) **165**, 59 (1985).

[39] H.Crater and L.Lusanna, *The Rest-Frame Darwin Potential from the Lienard-Wiechert Solution in the Radiation Gauge*, Ann.Phys.(N.Y.) **289**, 87 (2001)(hep-th/0001046).
D.Alba, H.Crater and L.Lusanna, *The Semiclassical Relativistic Darwin Potential for Spinning Particles in the Rest-Frame Instant Form: Two-Body Bound States with Spin 1/2 Constituents*, Int.J.Mod.Phys. **A16**, 3365 (2001) (hep-th/0103109).

[40] A.Barducci, R.Casalbuoni and L.Lusanna, *Energy-Momentum Tensor of Extended Relativistic Systems*, Nuovo Cim. **54A**, 340 (1979).

[41] R.J.Jantzen, P.Carini and D.Bini, *The Many Faces of Gravito-Magnetism*, Ann.Phys.(N.Y.) **215**, 1 (1992)(gr-qc/0106043).
Relative Observer Kinematics in General Relativity, Class.Quantum Grav. **12**, 2549 (1995).
The Intrinsic Derivatives and Centrifugal Forces in General Relativity. 1 Theoretical Foundations, Int.J.Mod.Phys. **D6**, 1 and *2 Applications to Circular Orbits in some Familiar Stationary Axisymmetric Space-Times*, **D6**,143 (1997)(gr-qc/0106013 and 0106014).
The Inertial Forces - Test Particle Motion Game, 1998 (gr-qc/9710051).
D.Bini, A.Merloni and R.T.Jantzen, *Adapted Frames for Space-Time Splittings with an Additional Observer Family*, Nuovo Cimento **113B**, 611 (1998).
D.Bini and R.T.Jantzen, *Circular Holonomy, Clock Effects and Gravito-Magnetism: Still Going around in Circles after All These Years..*, Proc. of the 9th ICRA Workshop on *Fermi and Astrophysics*, 2001, eds. R.Ruffini and C.Sigismondi (World Scientific, 2002)(gr-qc/0202085).

[42] R.A.d'Inverno and J.Stachel, *Conformal Two Structure as the Gravitational Degrees of Freedom in General Relativity*, J.Math.Phys. **19**, 2447 (1978).
R.d'Inverno, *2+2 Formalism and Applications*, in *Relativistic Gravitation and Gravitational Radiation*, Les Houches 1995, eds. J.A.Marck and J.P.Lasota (Cambridge Univ.Press, Cambridge, 1997).
R.d'Inverno and J.Smallwood, *Canonical 2+2 Formulation of the Initial Value Problem in General Relativity*, Phys.Rev. **D22**, 1233 (1980).
J.Smallwood, *Orthogonal 2+2 Decomposition of Space-Time*, J.Math.Phys. **24**, 599 (1983).
C.G.Torre, *Null Surface Geometrodynamics*, Class.Quantum Grav. **3**, 773 (1986).
S.A.Hayward, *Dual-Null Dynamics of the Einstein Field*, Class.Quantum Grav. **10**, 779 (1993).

[43] J.Stewart, *Advanced General Relativity* (Cambridge Univ. Press, Cambridge, 1993).

[44] J.Agresti, R.DePietri and L.Lusanna, in preparation.

[45] L.Lusanna and D. Nowak-Szczepaniak, *The Rest-Frame Instant Form of Relativistic Perfect Fluids with Equation of State $\rho = \rho(\eta, s)$ and of Non-Dissipative Elastic Materials*, Int.J.Mod.Phys. **A15**, 4943 (2000) (hep-th/0003095).

[46] D.Alba and L.Lusanna, *Eulerian Coordinates for Relativistic Fluids: Hamiltonian Rest-Frame Instant Form, Relative Variables, Rotational Kinematics* (hep-th/0209032).

[47] J.M.M.Senovilla, *Superenergy Tensors*, Class.Quantum Grav, **17**, 2799 (2000).